



GLOBAL JOURNAL OF MANAGEMENT AND BUSINESS RESEARCH: G
INTERDISCIPLINARY

Volume 25 Issue 1 Version 1.0 Year 2025

Type: Double Blind Peer Reviewed International Research Journal

Publisher: Global Journals

Online ISSN: 2249-4588 & Print ISSN: 0975-5853

Virtual Influences, Real Consequences: Social Media's Role in Food Preferences and Nutrition among Punjab's Adolescent Girls

By Dr. Simran Kaushal, Dr. Amarjeet kaur, Dr. Monika Choudhary
& Jaspreet kaur

Punjab Agricultural University

Abstract- This study investigates the influence of social media usage on the food choices and nutritional status of school-going adolescent girls in Punjab. A total of 520 subjects, aged 16-18 years, were selected from rural and urban schools across Punjab. The sample was divided into social media users and non-users based on daily usage duration, with further subdivision into rural and urban categories. Comprehensive data was collected on socio-economic status, dietary habits, physical activity and anthropometric parameters. Dietary intake was assessed using the 24-hour recall method and compared with the Recommended Dietary Allowances (RDA). Nutrient adequacy and dietary diversity scores were calculated, while body composition was analyzed using advanced tools. Results revealed significant differences in dietary patterns, nutrient intake, and body composition between users and non-users of social media. Rural non-users displayed healthier eating habits, including higher fruit consumption and lower fat intake, while urban users reported greater consumption of processed and junk foods.

Keywords: adolescents, social media, dietary patterns, nutritional status, food marketing, physical activity.

GJMBR-G Classification: DDC: 641.3



Strictly as per the compliance and regulations of:



© 2025. Dr. Simran Kaushal, Dr. Amarjeet kaur, Dr. Monika Choudhary & Jaspreet kaur. This research/review article is distributed under the terms of the Attribution-NonCommercial-NoDerivatives 4.0 International (CC BYNCND 4.0). You must give appropriate credit to authors and reference this article if parts of the article are reproduced in any manner. Applicable licensing terms are at <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

Virtual Influences, Real Consequences: Social Media's Role in Food Preferences and Nutrition among Punjab's Adolescent Girls

Dr. Simran Kaushal ^α, Dr. Amarjeet kaur ^σ, Dr. Monika Choudhary ^ρ & Jaspreet kaur ^ω

Abstract- This study investigates the influence of social media usage on the food choices and nutritional status of school-going adolescent girls in Punjab. A total of 520 subjects, aged 16-18 years, were selected from rural and urban schools across Punjab. The sample was divided into social media users and non-users based on daily usage duration, with further subdivision into rural and urban categories. Comprehensive data was collected on socio-economic status, dietary habits, physical activity and anthropometric parameters. Dietary intake was assessed using the 24-hour recall method and compared with the Recommended Dietary Allowances (RDA). Nutrient adequacy and dietary diversity scores were calculated, while body composition was analyzed using advanced tools. Results revealed significant differences in dietary patterns, nutrient intake, and body composition between users and non-users of social media. Rural non-users displayed healthier eating habits, including higher fruit consumption and lower fat intake, while urban users reported greater consumption of processed and junk foods. Social media users, particularly urban ones, showed increased exposure to food marketing and reported emotional eating behaviors linked to stress and peer influence. Despite differences in dietary patterns, both groups exhibited insufficient nutrient intake, a high prevalence of anemia and sedentary lifestyles. The study highlights the critical role of social media in shaping dietary habits and influencing adolescent health behaviors. It emphasizes the need for targeted public health interventions to mitigate the adverse impacts of social media while leveraging its potential to promote healthier eating habits and active lifestyles among adolescents in Punjab.

Keywords: adolescents, social media, dietary patterns, nutritional status, food marketing, physical activity.

I. INTRODUCTION

The word adolescence originated from the Latin term 'adolescere', which means to mature. Adolescence is defined as the age between 10 to 19 (WHO 2019). After infancy, adolescence is the only another time where rapid growth and development along with significant biological and psychological changes occur and these changes are greatly influenced by sociocultural factors. The two typical stages include early adolescence (10-14 years) and late

adolescence (15-19 years). From a physiological perspective, pubertal changes rule the early years whereas sexual maturation and establishment of adult roles and responsibilities rule the later years (Patton *et al* 2016). Adolescents make about 16 percent of the world's population (Bhargava *et al* 2020). Asia is home to more than half of the world's adolescents, while India has 20 percent of its people under the age of 18 (UNICEF 2016).

There is a link between adult comprehension of dietary guidelines and healthy eating behaviors. Nutritional knowledge is crucial for promoting healthy eating patterns and maintaining a healthy body weight. Relation among nutrition knowledge and food nutrient consumption as well as the positive effects of nutrition knowledge in adolescents cannot be ignored. Adolescence is a critical period of rapid growth requiring increased nutrients, such as amino acids for muscle development and calcium and vitamin D for bone growth. Adolescents, often engaged in physical activities, have higher energy needs, but those who are sedentary may accumulate fat, contributing to global adolescent obesity. According to WHO, about 1/5th of the world consist of adolescents, during this period good nutrition is very important for the growth and development. Inadequate nutrition during adolescence can have serious consequences throughout the reproductive years and beyond. Under-nutrition in adolescent girls is a major public health problem in developing countries (Das *et al* 2017).

Dietary problems among adolescent girls must be addressed as they can bring a negative effect on future generations if right information is not given at right time (Choudhary *et al* 2010). Each year, 16 million babies are born to girls aged 15-19, with adolescent fertility rates much higher in low-and middle-income countries (LMICs) due to factors like poverty, lack of education and limited healthcare access. Adolescent pregnancies pose significant risks, including competition for nutrients between mother and fetus, leading to stunted growth, low birth weight, preterm delivery and higher maternal mortality. Young mothers are also at increased risk for neonatal complications. During lactation, a mother's micronutrient status is vital for her infant's health, especially in the first six months. Nutrient needs during

Author ^α, ^σ, ^ρ: Punjab Agricultural University, Department of Food and Nutrition Ludhiana, Punjab. e-mail: simran1512kaushal@gmail.com

Author ^ω: Punjab Agricultural University, Department of Food Science and Technology Ludhiana, Punjab.

pregnancy and lactation are significantly higher compared to other stages of life (Das *et al* 2017).

Globally, the prevalence of overweight and obesity in children aged 5-19 has grown from 4 to 18 per cent since 1975 to 2016 (Abarca-Gomez *et al* 2017). Due to eating patterns influenced by obesogenic environments, such as the broad availability, high cost and promotion of foods with a high energy density, particularly in the digital era, health status of children and adolescents is deteriorating. Obesity has been related to prolonged television viewing (Olafsdottir *et al* 2014), primarily because it is associated with unfavourable eating habits such as skipping breakfast (Lipsky *et al* 2012) and consuming unhealthy foods more frequently (Lissner *et al* 2012) as well as fewer vegetables and fruits (Bornhorst *et al* 2015).

'Social Media' is a broad term used to describe online technical solutions that make it simple for users to communicate with one another and the online community by sharing and exchanging information online. There are a few examples of these technological tools such as social networking sites (SNS), video-sharing websites, virtual worlds, blogs, group projects etc (Mangold and Faulds 2009; Kaplan and Haenlein 2010). SNS have shown the fastest and most steady growth of any of these social media tools (Chang *et al* 2015). Social networking sites (SNS) have become a crucial source of information for customers, significantly influencing their choices and purchase intentions (Li and Chang 2016, Wang *et al* 2012). This impact encourages continued engagement with these platforms (Chang *et al* 2015). Over the past decade, the use of Facebook has surged from about 2% to nearly 30% of the global population by 2019, highlighting the growing importance of SNS in daily life, especially among adolescents (Ortiz-Ospina 2019). Children's and adolescents attention spans can be impacted by the need to often check highly engaging online content and incoming notifications from SM applications (Reid *et al* 2016). This effect is particularly concerning when people are eating thoughtlessly in front of screens which may lead to overeating, consequently resulting in overweight and obesity (Robinson *et al* 2017). Food communications have permeated our surroundings in the form of social media photos, videos, advertisements and posts from food influencers, cookery shows and peer-to-peer messages. These signals can be virtual or nonvirtual and frequently support unhealthy non-core eating norms while glorifying non-core food consumption and enticing excessive energy consumption (Qutteinaet *et al* 2021).

Social media influencers, defined as individuals who cultivate their own audiences on these platforms, significantly affect the attitudes and behaviors of their followers (Gross and Wangenheim 2018, Freberg *et al* 2011). Often referred to as "cyber celebrities," these influencers gain fame through sharing personal narratives, similar to traditional celebrities and leverage

their social influence to promote branded products (De Veirman *et al* 2017, Chatterjee 2011). Their promotional strategies can range from subtle integrations of branded content to more explicit calls for followers to engage with their posts (Abidin, 2016). Research indicates that such marketing can notably impact food preferences; for instance, exposure to influencers promoting unhealthy snacks has been linked to increased consumption of non-core foods and preferences for branded over non-branded options among preadolescents (Coates *et al* 2019b, Coates *et al* 2019c). Thus, food marketing by social media influencers plays a significant role in shaping dietary choices and behaviors.

Adolescents are heavily influenced by food-related posts from peers and influencers on social media, which often highlight non-core foods, such as soft drinks and fast food. This exposure shapes their perceptions of food norms, leading them to believe that overconsumption of non-core foods is typical, while underestimating the popularity of healthier options. This misperception is concerning, as food norm perceptions significantly affect eating behaviors. The constant portrayal of non-core foods may create a feedback loop where adolescents internalize these norms, increasing their consumption of unhealthy foods. Thus, social media not only reflects but also reinforces eating behaviors, making it crucial to understand how these online food messages impact adolescents' food choices (Qutteinaet *et al* 2021).

Punjab is an economically successful state with a high per capita income. Punjab has always been known for its delectable cuisine and adolescents in particular have exceptional taste preferences. Because of industrialization, urbanization and increased spending power in this century, lifestyle and eating habits have altered. Adolescence is seen as a crucial period for establishing dietary habits since it affects the likelihood of developing a variety of ailments later in life. Foods that provide enough energy and nutrients to sustain physical health, cognitive development and social growth are the foundation of children's and teenagers health. The consumption of a balanced diet is essential for children's growth and development and is also responsible for preventing non-communicable diseases. During this time, family plays a crucial role.

In light of these ambiguities, it is crucial to appreciate and incorporate the body of research that can analyze the positive or negative use of online social networking as a component intervention to promote health in order to strengthen future research and practices. The purpose of the present research study is to look into how adolescent girls use social media in terms of food information in relation to nutritional status of rural and urban girls, particularly belonging to Punjab.

II. MATERIAL AND METHODS

The purpose of the present study was to investigate Influence of Social Media Usage on Food Choices and Nutritional Status of School Going Adolescent Girls Residing in Punjab. This research encompassed investigations from the selected district city will be considered as urban sample whereas the samples from the selected villages will constitute rural sample and further categorized into users and non-users of social media, employing the methodologies outlined below:

a) Location/place of Work

In order to identify social media users and non-users, the research was carried out in the private and government schools of urban and rural areas of Punjab.

b) Selection of Subjects

A total sample of 520 girls in the age group of 16-18 years was selected from private and government schools of Punjab. There were a total of 23 districts in Punjab, with the Malwa region occupying the largest

part of the state, comprising 15 districts. The remaining districts were located in the Majha and Doaba regions, with an equal number in each. For this study, districts from each region were selected using a proportionate sampling technique, applying a 1:1:3 ratio. Thus, two districts each from Majha and Doaba and six districts from the Malwa region were selected to ensure coverage of the entire state. In the chosen villages and district cities, government and private schools were selected for the subject sample. The samples from the selected district cities were considered urban, while those from the selected villages constituted the rural sample. The subjects were divided into the following groups, with an equal number of subjects in each:

Control: Subjects who used social media for 1-1.5 hours were categorized as non-users or classified according to the data obtained from the subjects.

Experimental: Subjects who used social media for more than 1-1.5 hours were categorized as users.




Figure 3.1: Areas of Three Regions of Punjab

c) *Drafting a Questionnaire*

The questionnaire was thoughtfully created to gather comprehensive information from the chosen participants in a systematic way. A wide range of information was covered in the survey, such as age, caste, family type, income and dietary habits assessed through a structured dietary survey, 24-hour recall technique was applied and questions about food and nutrient intake, anthropometric measurements, body composition and social media platform use were asked.

d) *Data Collected from 520 Subjects*

A well-constructed and pretested interview schedule was used to look into a number of topics related to the study's goals throughout the data gathering phase. The interview schedule included inquiries regarding topics like socio-economic and demographic profile, the influence of social media on food information, dietary habits, physical activity, anthropometric parameters and body composition.

i. *Socio-Economic and Demographic Profile*

The information related to the subject's age, caste, type of family, family composition, educational and occupational status of parents and total monthly income of the family were collected using pretested interview schedule. The per capita monthly income of the individual family members was calculated by dividing the total income of the family by the number of members. Modified Kuppaswamy's socio economic status scale was used to know the socio economic class of the subjects. Education, occupation and per capita income were considered to socio-economic status (Wani 2019).

Table 3.1: Modified Kuppaswamy's Socio Economic Status Scale

Socio-Economic Class	Total Score
Upper Class	26-29
Upper Middle	16-25
Lower Middle	11-15
Upper Lower	5-10
Lower	Below 5

ii. *Influence of Social Media on Food Information*

The data pertaining to the type of social media platform used, duration of usage, frequency of usage, purpose and attributes given importance, as well as the food brands and type of food influencer followed by the subjects was recorded.

iii. *Dietary Information*

a. *Dietary Pattern*

A general dietary assessment include dietary habits, meal frequency and the consumption of water, beverages, green leafy vegetables, fruits and meat products.

The frequency of common food items, especially those popular among children, was recorded.

b. *Food Intake*

Data on the dietary intake of adolescents was collected using the 24-hour recall method over three consecutive days, including one Sunday or holiday. Days selected were not fasting or feasting days to ensure typical dietary patterns.

Standardized cups, glasses and spoons, which had been previously calibrated, were used as aids to assist participants in recalling the quantities of food consumed during the 24-hour period prior to the investigation.

c. *Nutrient Intake*

The daily nutrient intake was determined using DIETCAL software (Kaur, 2017), and the results were compared with the recommended dietary allowances to evaluate nutrient adequacy.

d. *Nutrient Adequacy Ratio (NAR)*

The average amount of each food ingested by each subject over the course of three days was entered into the software, and the software then estimated the subject's nutritional intake and compared it to the RDA (ICMR 2020).

$$\text{NAR \%} = (\text{Intake of nutrient/Recommended intake of nutrient}) * 100$$

Table 3.2: Categorization of Nutrient adequacy ratio (NAR%) using a classification given by (Rani and Rani 2016)

Adequate	100% and above
Marginally adequate	75% and above
Marginally inadequate	50 to 74.9%
Inadequate	Below 50%

e. *Dietary Diversity Score (DDS)*

The Dietary Diversity Score (DDS) was measured by categorizing the food items consumed in a day into 14 groups. The participants were categorized into three classes according to their scores, using the classification given by Nithya and Bhavani (2018).

Table 3.3: Categorization of Dietary Diversity Score (DDS) using a Classification given by Classification by Nithya and Bhavani (2018)

Low Dietary Diversity	DDS of 1-4
Moderate Dietary Diversity	DDS of 5-9
High Dietary Diversity	DDS of 10-14

f. *Height and Weight*

For height measurement, a stadiometer or measuring tape was used. The measuring area was ensured to be flat and well-lit and participants were asked to remove shoes and hats. They stood straight against the measuring device, making sure their heels, buttocks, shoulders and heads were in contact with the surface. Participants looked straight ahead, with their eyes level to the horizon, while the horizontal headpiece of the stadiometer was gently lowered onto their heads.

The height was recorded in centimeters, accurate to the nearest 0.1 cm.

For weight measurement, a calibrated weighing scale was utilized. The scale was placed on a flat, hard surface to ensure accuracy and it was zeroed before use. Participants were asked to remove shoes and heavy clothing and to stand still in the center of the scale, with feet evenly spaced. The weight was recorded in kilograms, accurate to the nearest 0.1 kg. All measurements were taken under consistent conditions to effectively assess growth and nutritional status, following the guidelines established by Jelliffe.

g. *Waist and Hip Circumference*

The waist and hip circumferences were measured using a non-stretchable measuring tape. The participant was instructed to stand upright with their feet together and arms relaxed at their sides, wearing light clothing. For the waist circumference measurement, the

midpoint between the lower rib and the top of the hip bone (iliac crest) was located. The measuring tape was wrapped around the waist at this midpoint, ensuring it was parallel to the ground and snug but not compressing the skin. The participant was asked to breathe normally, and the waist circumference was recorded at the end of a normal expiration in centimeters.

For the hip circumference measurement, the widest part of the hips or buttocks was identified. The measuring tape was then wrapped around this area, ensuring it remained parallel to the ground. The hip circumference was also recorded in centimeters. Both measurements were accurately documented for further analysis, aiding in the assessment of body fat distribution and the associated risk of noncommunicable diseases.

Table 3.4: Waist and Hip Circumference Classification as Outlined by the WHO (2008)

Classification	Waist Circumference (Inches)	Hip Circumference (Inches)
Normal	Women: < 31.5	Women: < 34.5
Increased Risk	Women: 31.5 - 34.5	Women: 34.5 - 37
High Risk	Women: > 34.5	Women: > 37

h. *Body Mass Index (BMI)*

Using the following formulas, BMI was computed from weight and height records (WHO, 2004):

$$\text{Body Mass Index (BMI)} = \frac{\text{Weight (kg)}}{(\text{Height}) (\text{m}^2)}$$

Asian-Pacific BMI Classification

BMI Category	BMI Range (kg/m ²)
Underweight	< 18.50
Normal range	18.5 - 22.99
Overweight	23- 24.99
Pre-obese	25- 29.99
Obese	>30.0
Grade I	30-40
Grade II	40.1-50
Grade III	>50

i. *Waist to Hip Ratio (WHR)*

A continuous tension tape was used to measure the circumference of the waist and hip in centimeters. Measurements were taken after wrapping the tape around the hip. The waist-hip ratio (WHO 2008) was calculated. Additionally, the waist to hip ratio was computed by entering the appropriate values into the procedure below.

$$\text{Waist Hip Ratio} = \frac{\text{Hip circumference}}{\text{Waist circumference}}$$

According to WHO, abdominal obesity is indicated by a waist-hip ratio more than 0.85 for women (WHO2008).

j. *Waist to Height Ratio*

This helped in assessing health risks related to central obesity

$$\text{WHtR} = \frac{\text{Waist circumference}}{\text{Height}}$$

Table 3.5: Categorization of Waist to Height Ratio (Ashwell and Hsieh 2005)

WHtR	Risk
< 0.40	Low (Underweight)
0.40 - 0.49	Healthy (Low Risk)
0.50 - 0.59	Increased Risk
≥ 0.60	High Risk (Obesity)

The health risks related to obesity affected both men and women, but evidence suggested that women faced greater health risks due to obesity compared to men in some areas. However, the distribution and type of health risks varied by gender due to biological, hormonal and behavioral differences (WHO 2004).

e) *Statistical Analysis*

A variety of statistical analyses were performed on the data to ascertain various parameters.

- Calculating certain descriptive statistical metrics for every research variable, including the mean, standard deviation and percentage distribution.
- An independent t-test was applied to test the difference between parameters.
- Pearson's correlation coefficient is evaluated with respect to other variables.

III. RESULTS

a) Socio-Demographic Profile of Participants

The study sample (Table 4.1) included adolescent girls from rural and urban areas, predominantly aged between 16 and 18. The 16–17-year-old age group had a higher percentage in rural areas (87.69%) compared to urban areas (86.93%), while 18-year-olds constituted a small proportion across both groups. A majority of participants in rural areas were Sikh (72.31%), with a slightly higher prevalence than in urban areas (60.77%). Hindu participants formed 25.77% of the rural and 37.69% of the urban sample. Minority religious groups included Muslims (1.54% in rural, 0.77% in urban) and Christians (0.38% in rural, 0.77% in urban). Participants' caste affiliations varied, with the scheduled caste comprising 58.85% of rural participants and 29.62% of urban participants. The general caste was more prevalent in urban areas (44.61%) compared to rural areas (20.77%). Other caste groups included the backward class (4.62% rural, 11.54% urban) and other backward classes (15.77% rural, 14.23% urban). Nuclear families were more common, with 70.38% in rural and 64.62% in urban areas. Joint family structures were also present but less predominant, representing 29.62% of rural and 35.38% of urban participants.

Most participants came from medium-sized families (5 to 8 members), accounting for 59.23% in rural and 54.23% in urban areas. Large families (more than 8 members) were less common, while small families (less than 4 members) were represented by 27.31% in rural and 26.54% in urban areas.

Income distribution showed that most families earned between 2.5 to 5 lakhs annually, with 48.56% of rural and 43.85% of urban participants falling into this income bracket. A smaller percentage reported incomes over 5 lakhs, and only a minimal proportion of families earned above 10 lakhs. The majority of participants were from the lower-middle and upper-lower socioeconomic classes.

b) Medical Background of Adolescent Participants

Among the adolescent girls surveyed (Table 4.2), the prevalence of major health disorders such as diabetes was absent in both rural and urban groups, with all participants reported as non-diabetic. A small percentage reported hypertension, with 1.53% of rural and 0.38% of urban participants diagnosed with the condition. Depression was identified in only one participant (0.38%) in rural areas, reflecting a low incidence across the sample. Allergic reactions were reported more frequently in urban participants, where 7.30% reported allergies, compared to 2.69% in rural areas. The majority of participants across both locations did not report any allergies. Medical check-up attendance showed significant variation; 42.31% of rural

and 48.08% of urban participants had undergone recent check-ups. However, a larger portion, especially in rural areas (57.69%), did not attend regular medical evaluations. Medication usage was relatively low among participants, with only 5.77% in rural and 6.15% in urban areas actively using prescribed medications. Among these, the types of medication varied, with some use of homeopathic (3.07% rural, 1.54% urban), Ayurvedic (1.54% urban only) and allopathic medications (2.69% rural, 3.08% urban). The usage of corrective eyewear, such as spectacles during device usage, was more common among urban participants (29.62%) compared to those in rural areas (15.77%). These findings provide a comprehensive insight into the medical background of adolescent girls in rural and urban settings, highlighting both health conditions and medical practices across different demographic groups.

c) Menarche Status of Adolescent Participants

The distribution of menarche age among adolescent participants varied, with the majority experiencing menarche between ages 12 and 14 (Table 4.3). Specifically, 31.15% of rural and 34.62% of urban participants reported menarche at age 13, making it the most common age of onset. Menarche at age 12 was also prevalent, with 27.31% in rural and 21.92% in urban areas, followed by age 14 (25% rural, 24.23% urban). Fewer participants reported menarche at ages younger than 11 or older than 15, indicating that the majority reached menarche during early adolescence. Menstrual cycle regularity differed slightly between groups. In rural areas, 75.38% reported regular cycles, compared to 69.23% in urban settings. Irregular cycles were reported by 24.62% of rural and 30.77% of urban participants. Experiences of menstrual pain were common, with variations in pain timing. Pre-menstrual pain affected 47.69% of rural and 45.77% of urban participants, while pain throughout the menstrual period was reported by 42.31% in rural and 46.15% in urban areas. Post-menstrual pain was less frequent but present in 5.77% of rural and 6.54% of urban participants, while only 4.23% of rural and 1.54% of urban participants reported no menstrual pain. A notable portion of participants used medication to manage menstrual discomfort, with 10% of rural and 13.08% of urban participants taking pain relief medications. The remaining participants did not rely on medication for pain management. The volume of menstrual flow was predominantly reported as normal by both rural (85.77%) and urban (82.31%) participants. Heavy flow was less common, observed in 12.69% of rural and 13.46% of urban participants, while low flow was less (1.54% rural, 4.23% urban). These findings provide insight into the menstrual health of adolescent girls, highlighting common ages of menarche onset, regularity patterns, pain experiences and variations in menstrual flow across rural and urban populations.

d) *Influence of Social Media on Food Information*

Table 4.4 provides a detailed analysis of social media (SM) usage patterns among adolescent

participants, revealing significant behavioral differences between SM users and non-users.

Table 4.4: Parameters of Social Media (SM) usage among Respondents

Statements	User (n=260)	Non user (n=260)	Total (n=520)
Personal Mobile Phone or any other Device			
1. Yes	161(61.92)	126(48.46)	287
2. No	99(38.07)	134(51.53)	233
Keep a Check on Screen Time while using SM			
1. Yes	143(55)	155(59.61)	298
2. No	117(45)	105(40.38)	222
Frequency of Visiting SNS			
1. Everyday	260(100)	212(81.53)	472
2. Twice a day	-	-	-
3. Thrice a day	-	48(18.46)	48
Purpose of SM use			
1. Keeping in touch with friends & family	68(26.15)	74(28.46)	142
2. Just for fun	8(3.08)	10(3.84)	213
3. For health purpose	122(46.92)	91(35)	18
4. For updating yourself	62(23.84)	85(32.69)	147
Sites Attracted the Most			
1. Facebook	-	14(5.38)	14
2. Instagram	137(52.69)	92(35.38)	246
3. Youtube	79(30.38)	109(41.92)	171
4. Snapchat	31(11.92)	18(6.92)	49
5. Whatsapp	13(5)	27 (10.38)	40
Satisfaction from SM apps			
1. Highly satisfied	31(11.92)	63(24.23)	94
2. Satisfied	207(79.61)	187(71.92)	394
3. Dissatisfied	14(5.38)	7(2.69)	21
4. Highly dissatisfied	8(3.07)	3(1.15)	11
Time Spent on SNS			
1. <1.5	-	260(100)	260
2. 1.5-3	122(46.92)	-	122
3. 3-6	98(37.69)	-	98
4. ≥6	40(15.38)	-	40
Emotions Experienced while using SNS			
1. Rejection	5(1.92)	1(0.38)	6
2. Happiness	132(50.77)	80(30.77)	212
3. Boost Self Esteem	27(10.38)	39(15)	66
4. Jealousy	5(1.92)	-	5
5. Motivation/Inspiration	81(31.15)	135(51.92)	216
6. Fear Of Missing Out	9(3.47)	3(1.15)	12
7. Lower Self Esteem	1(0.38)	2(0.77)	3
SM used for Learning and Attract to People towards Healthy Behavior			
1. Yes	213(81.92)	239(91.92)	452
2. No	47(18.07)	21(8.07)	68
Access of Devices			
1. Mobile	250(96.15)	23(8.84)	273
2. Laptop	3(1.15)	-	3
3. Computer	5(1.92)	160(61.53)	165
4. Tablet	2(0.77)	77(29.61)	79
Mostly used Device			
1. Mobile	248(95.38)	27(10.38)	275
2. Laptop	4(1.53)	-	4
3. Computer	4(1.53)	153(58.84)	157
4. Tablet	4(1.53)	80(30.77)	84
SM Effected your Nutritional and Health Status			
1. Yes	159(61.15)	116(44.61)	275

2. No	101(38.84)	144(55.38)	245
Use SM for Practical Nutrition Purposes, Including Recipes			
1. Yes	220(84.61)	184(70.77)	404
2. No	40(15.38)	76(29.23)	116
SM platforms used to access nutrition information			
1. Youtube	178(68.46)	210(80.77)	388
2. Instagram	78(30)	48(18.46)	126
3. Facebook	3(1.15)	-	3
4. Telegram	1(0.38)	2(0.77)	3
Follow any Fitness or Health Professional/Influencer Online			
1. Yes	154(59.23)	119(45.77)	273
2. No	106(40.77)	141(54.23)	247
Nutrition Information Gained from SM is Reliable in the Real World			
1. Yes	128(49.23)	108(41.53)	236
2. No	132(50.77)	152(58.46)	284
Influencers have the Power to Influence others by their Lifestyle, Behavior etc			
1. Yes	222(85.38)	176(67.69)	398
2. No	38(14.61)	84(32.30)	122
Consume/Purchase Food Brand Promoted by an Influencer			
1. Yes	42(16.15)	27(10.38)	69
2. No	218(83.85)	233(89.61)	451
Type of Influencer Followed			
1. Fashion	148(56.92)	121(46.53)	269
2. Gaming	12(4.61)	36(13.85)	48
3. Food	100(38.46)	103(39.61)	203
Interested in Joining Online Nutrition Camps through Any Social Networking Site			
1. Yes	96(36.92)	66(25.38)	162
2. No	164(63.07)	194(74.61)	358
Diet in Order to Change your Body Image Like your Favourite SM Personality			
1. Yes	91(35)	68(26.15)	159
2. No	169(65)	192(73.85)	361
Networking Sites Helped you Out in Gaining and Losing Weight			
1. Yes	96(36.92)	76(29.23)	172
2. No	164(63.08)	184(70.77)	348
Making use of these Apps have Altered Sleeping and Eating Pattern			
1. Yes	143(55)	125(48.08)	268
2. No	117(45)	135(51.92)	252
Qualities Leading to Follow an Influencer			
1. Professional/qualified	92(35.38)	103(39.61)	191
2. Funny/interactive	88(33.85)	84(32.30)	176
3. Evidence based/talented	17(6.53)	24(9.23)	41
4. Regular posting	16(6.15)	11(4.23)	27
5. Body image positivity	15(5.77)	8(3.07)	23
6. Shares personal experiences	32(12.30)	30(11.53)	62
Recipe and Food Purchasing Decision Changes After using SM			
1. Yes	174(66.92)	150(57.69)	324
2. No	86(33.07)	110(42.30)	196
How useful is SM for Learning			
1. Not useful	11(4.23)	-	11
2. Somewhat useful	96(36.92)	90(34.61)	186
3. Very useful	153(58.84)	170(65.38)	323
Search Content Related to Dieting, Fitness and Health			
1. Yes	93(35.77)	97(37.31)	190
2. No	41(15.77)	56(21.54)	97
3. Read only when it appears	126(48.46)	107(41.15)	233
Type of Content Like to See on SM			
1. Fitness	53(20.38)	61(23.46)	114
2. Nutrition	25(9.61)	10(3.85)	35
3. Health and lifestyle	182(70)	189(72.69)	371

Tried a Diet Promoted by an Social Media Influencer			
1. Very frequently	27(10.38)	13(5)	40
2. Frequently	39(15)	25(9.61)	64
3. Occasionally	25(9.61)	19(7.31)	44
4. Rarely	71(27.31)	89(34.23)	160
5. Never	98(37.69)	114(43.85)	212
SM Helped in Providing Nutritional knowledge			
1. Yes	149(57.30)	112(43.07)	261
2. No	111(42.69)	148(56.92)	259
SM Platforms used to Access Nutrition Information			
1. Youtube	192(73.84)	228(87.69)	42
2. Instagram	65(25)	30(11.54)	95
3. Facebook	2(0.77)	-	2
4. Telegram	1(0.38)	2(0.77)	3
SM Leads to Opt more of			
1. Healthy foods	128(49.23)	163(62.69)	291
2. Junk foods	132(50.77)	97(37.31)	229
Make use of Any Health-App to Stay Healthy			
1. Yes	66(25.38)	53(20.38)	119
2. No	194(74.62)	207(79.62)	401
Heard about Different Types of Diets among the Options			
1. Mediterranean diet	19(7.31)	17(6.54)	39
2. Intermittent fasting	11(4.23)	2(0.77)	13
3. Vegan diet	45(17.31)	23(8.85)	68
4. Low carb diet	21(8.08)	7(2.69)	28
5. Ketogenic diet	3(1.15)	12(4.62)	15
6. None of the above	161(61.92)	199(76.54)	360
Change in Existing Diet and Physical Activity Patterns After using SM Platforms			
1. Yes	73(28.08)	62(23.85)	135
2. No	187(71.92)	198(76.15)	385
Like to Entertain the Nutritional Advice			
1. SM influencers	121(46.54)	91(35)	212
2. Qualified expert	139(53.46)	169(65)	308
Prefer while Choosing Food			
1. Taste of food	185(71.15)	188(72.31)	373
2. Nutrition content	52(20)	56(21.53)	108
3. Colour of food	5(1.92)	5(1.92)	10
4. Aroma of food	18(6.92)	11(4.23)	29
Prefer the Most			
1. Ordering online	82(31.54)	63(24.23)	145
2. Visiting store	178(68.46)	197(75.77)	375
Sleeping Hours			
1. 4-6 hours	71(27.31)	121(46.54)	192
2. 6-8 hours	144(55.38)	122(46.92)	266
3. More than 8 hour	45(17.31)	17(6.54)	62
Posts on SM Leads to Cravings			
1. Yes	187(71.92)	162(62.31)	349
2. No	73(28.08)	98(37.69)	171
Usage of SM Affected your Food Consumption			
1. Yes	114(43.85)	78(30)	192
2. No	146(56.15)	182(70)	328
Use SM to Receive Information about Healthy Eating			
1. Yes	162(62.31)	179(68.85)	341
2. No	98(37.69)	81(31.15)	179
Check out the Nutritional Quality of Food Product before Consuming			
1. Yes	159(61.15)	196(75.38)	355
2. No	101(38.85)	64(24.62)	165
Make Food Choices According the Healthiness of Food			
1. Yes	130(50)	162(62.31)	292

2. No	130(50)	98(37.69)	228
Utilize SM while Making Decisions in Regards to the Food you Eat at Restaurants			
1. Yes	68(26.15)	62(23.85)	130
2. No	192(73.85)	198(76.15)	390
Skip Meals While using SM Platforms			
1. Yes	98(37.69)	74(28.46)	172
2. No	162(62.31)	186(71.54)	348
More Prone to Eating While using SM Platforms			
1. Yes	135(51.92)	92(35.38)	227
2. No	125(48.07)	168(64.62)	293
Ever Shared Information or Advice Related to Nutrition with Friends and Family			
1. Yes	167(64.23)	164(63.07)	331
2. No	93(35.77)	96(36.92)	189
Skip any Outdoor Activity to Watch SM			
1. Yes	118(45.38)	102(39.23)	220
2. No	142(54.62)	158(60.77)	300
SNS Affected Mental Health and Overall Wellbeing			
1. Yes	151(58.08)	148(56.92)	299
2. No	109(41.92)	112(43.08)	221

Figures in parenthesis indicate percentages
SM (Social Media)
SNS (Social Networking Sites)

e) Dietary Information

i. Dietary Pattern

The study evaluated food preferences and dietary behaviors among rural and urban populations, each consisting of 260 participants divided equally into users and non-users (n=130 in each group). In terms of food habits, 51.92% of the total sample (n=270) identified as vegetarians, with rural users (59.23%) having the highest proportion, followed by urban non-users (55.38%). Non-vegetarians accounted for 30.77% (n=160) of the participants, with similar proportions across all subgroups. Ova-vegetarians, making up 17.31% (n=90) of the sample, were most common among rural non-users (24.62%).

Regarding meal frequency, the majority of respondents (71.15%, n=370) reported consuming three meals per day, with urban non-users (91.54%) being the most consistent in this pattern. A smaller group (19.23%, n=100) reported having four meals daily, particularly among rural users (27.69%). Only 9.62% (n=50) reported eating two meals, mostly rural users (26.15%).

Water intake varied across the sample, with 47% (n=244) consuming 4–6 glasses daily and 37.88% (n=197) reporting drinking more than six glasses, with non-users generally exhibiting higher water consumption. A small percentage (15.19%, n=79) drank less than four glasses of water, more notably among rural users (24.62%). Snacking behavior was prevalent, with 92.12% (n=479) of participants reporting a preference for snacking or munching. This behavior was especially prominent among rural users (96.15%) and urban users (94.62%), indicating that both populations strongly favored between-meal snacks.

ii. Food Intake

1. Cereals and Millets

Rural participants showed slightly higher cereal and millet intake overall, with users averaging 137.88 ± 47.82 grams compared to non-users at 129.47 ± 38.83 grams. Interestingly, urban non-users reported consuming more (131.19 ± 39.86 grams) than users (125.60 ± 54.30 grams). These differences, however, were statistically non-significant, suggesting that cereal consumption was fairly consistent across both settings, possibly reflecting the staple nature of these foods in both rural and urban diets. Vegetarian as well as non-vegetarian dietary pattern of Indians include consumption of cereals as staple and rice remains priority.

2. Pulses and Legumes

Consumption of pulses and legumes varied slightly. In rural areas, non-users reported a higher intake (75.25 ± 47.27 grams) than users (67.45 ± 51.84 grams). Similarly, in urban settings, non-users consumed more (70.75 ± 51.98 grams) than users (60.11 ± 37.03 grams). While pulses are excellent, inexpensive sources of protein, the non-significant differences suggest that both users and non-users, irrespective of setting, access these foods to meet their nutritional needs.

3. Green Leafy Vegetables (GLV)

Green leafy vegetables (GLV) intake was generally higher in urban areas compared to rural ones. Urban users consumed 48.56 ± 19.22 grams, while non-users averaged 45.45 ± 16.21 grams. Conversely, rural participants reported a lower intake, with users consuming 42.05 ± 15.97 grams and non-users 44.22 ± 12.10 grams. The lack of significant differences indicates that GLV consumption is maintained at

comparable levels across all groups, though slightly elevated among urban participants, possibly due to better availability or awareness.

4. Other Vegetables

The intake of other vegetables also varied between groups. Rural users consumed an average of 105.90 ± 63.21 grams, slightly more than non-users at 94.91 ± 59.59 grams. In urban areas, however, non-users reported slightly higher consumption (93.47 ± 34.41 grams) compared to users (87.70 ± 57.39 grams). Despite these minor variations, the differences remained non-significant, suggesting that both rural and urban populations, regardless of social media usage, incorporate a similar amount of vegetables into their diet.

5. Fruits

Fruit consumption displayed a significant difference in rural settings ($p < 0.05$). Rural non-users consumed more (92.67 ± 58.24 grams) than users (61.97 ± 30.85 grams). This may indicate that rural non-users rely more heavily on fruits as a part of their regular diet. However, in urban areas, no significant difference was observed, with users consuming 47.84 ± 33.30 grams and non-users 44.47 ± 25.26 grams. The discrepancy in rural fruit intake might reflect lifestyle or economic differences between users and non-users.

Table 4.7: Food Intake of the Selected Subjects

Food groups	SDI (g) [#]	Rural (n=260)			Urban (n=260)		
		User (n=130)	Non user (n=130)	t test (p value)	User (n=130)	Non user (n=130)	t test (p value)
Cereals and millets	315	137.88 ± 47.82	129.47 ± 38.83	$1.556(0.120)^{NS}$	125.60 ± 54.30	131.19 ± 39.86	$0.947(0.344)^{NS}$
Pulses and legumes	105	67.45 ± 51.84	75.25 ± 47.27	$1.264(0.207)^{NS}$	60.11 ± 37.03	70.75 ± 51.98	$1.900(0.058)^{NS}$
GLV	100	42.05 ± 15.97	44.22 ± 12.10	$1.234(0.218)^{NS}$	48.56 ± 19.22	45.45 ± 16.21	$1.410(0.159)^{NS}$
Other vegetables	200	105.90 ± 63.21	94.918 ± 59.59	$1.441(0.150)^{NS}$	87.70 ± 57.39	93.47 ± 34.41	$0.983(0.326)^{NS}$
Fruits	150	61.97 ± 30.85	92.67 ± 58.24	$4.083(0.000)^*$	47.84 ± 33.30	44.47 ± 25.26	$0.550(0.583)^{NS}$
Roots and tubers	100	53.93 ± 39.02	60.71 ± 50.10	$1.217(0.224)^{NS}$	49.69 ± 26.83	54.66 ± 37.84	$1.409(0.165)^{NS}$
Milk and milk products	400	341.60 ± 135.8	337.05 ± 140.8	$0.264(0.791)^{NS}$	327.55 ± 129.8	326.58 ± 160.9	$0.053(0.957)^{NS}$
Edible oils and fats	40	30.19 ± 6.92	21.93 ± 8.67	$8.455(0.000)^*$	21.34 ± 10.24	16.13 ± 11.39	$3.872(0.000)^*$

[#]Suggested dietary intake

Values indicates Mean and Standard Deviation

*Values are significant at 1% level

**Values are significant at 5% level

^{NS}Values are Non-Significant






Fig. 4.1: Nutrient Adequacy Ratio of Food Groups among Rural user and Non-User Group and Urban User and Non-User Groups

6. Roots and Tubers

Rural users reported a slightly lower intake of roots and tubers (53.93 ± 39.02 grams) compared to non-users (60.71 ± 50.10 grams). Similarly, in urban areas, users consumed 49.69 ± 26.83 grams, slightly less than non-users (54.66 ± 37.84 grams). These differences were not statistically significant, indicating that roots and tubers are consumed consistently across different groups.

7. Milk and Milk Products

The intake of milk and milk products was relatively high across all groups, with rural users consuming 341.60 ± 135.8 grams and non-users reporting 337.05 ± 140.8 grams. In urban areas, users reported a similar intake (327.55 ± 129.8 grams) compared to non-users (326.58 ± 160.9 grams). The lack of significant differences suggests that milk is an essential dietary component for both users and non-users, regardless of their geographical location.

8. Edible Oils and Fats

Significant differences in edible oils and fats consumption were observed in both rural and urban areas ($p < 0.05$). Rural users consumed significantly more (30.19 ± 6.92 grams) compared to non-users (21.93 ± 8.67 grams). Similarly, in urban areas, users reported a higher intake (21.34 ± 10.24 grams) than non-users (16.13 ± 11.39 grams). These findings suggest that social media users may be more inclined to consume higher amounts of fats, potentially influenced by trends in cooking practices or the promotion of specific diets online.

iii. Nutrient Intake

1. Energy

Both rural and urban populations reported average energy intakes below the recommended 2500 kcal per day. Rural users exhibited a mean energy intake of 2083 kcal, compared to 1920.62 kcal in rural non-users, while urban users and non-users reported 1863.55 kcal and 1714.8 kcal, respectively. Despite these differences, no statistical significance was observed ($p > 0.05$).

2. Protein

Protein intake also remained below the RDA of 46g for all groups. Rural users consumed an average of 31.87 ± 14.52 g, while non-users reached 33.65 ± 12.75 g. In urban areas, users averaged 27.54 ± 14.51 g, with non-users at 28.11 ± 15.44 g. Again, differences between groups were not statistically significant ($p > 0.05$).

3. Total Fat

Rural and urban populations consumed levels of fat just under the recommended 25g, with rural users averaging 22.14g compared to 17.92g for non-users, while urban users and non-users consumed 20.34g and 16.83g, respectively. Differences were statistically significant in both settings ($p < 0.05$), suggesting that users generally had higher fat intakes.

4. Total Dietary Fibre

Dietary fiber intake showed close adherence to the 38g RDA among rural users (32.04g) and non-users (33.20g), while urban users and non-users recorded averages of 27.95g and 29.50g, respectively, with no significant difference between groups ($p > 0.05$).

5. Calcium

Calcium intake fell short of the 1050mg RDA across all groups. Rural users averaged 1005.6mg, and rural non-users reported 966.52mg, while urban users and non-users had averages of 673.14mg and 714.32mg, respectively, without significant differences ($p > 0.05$).

6. Iron

Iron intake was markedly lower than the RDA of 32mg for both rural and urban participants. Rural users averaged 11.91 ± 4.03 mg, while non-users consumed 14.97 ± 3.65 mg, with a significant difference between

these groups ($p < 0.05$). In urban areas, users and non-users reported averages of 10.85mg and 11.64mg, respectively, though this difference was not significant ($p > 0.05$).

7. Magnesium

With an RDA of 380mg, magnesium intake was reported as lower among all groups, with rural users consuming 288.93mg compared to 302.46mg in non-users. Urban users and non-users reported 221.29mg and 229.32mg, respectively, with no significant differences across groups ($p > 0.05$).

Table 4.8: Mean Daily Average Consumption of Different Nutrients among Selected Adolescents

Nutrients	RDA [#]	Rural (n=260)			Urban (n=260)		
		User (n=130)	Non user (n=130)	t test (p value)	User (n=130)	Non user (n=130)	t test (p value)
Protein (g)	46	31.87 ± 14.52	33.65 ± 12.75	$1.047(0.295)^{NS}$	27.54 ± 14.51	28.11 ± 15.44	$0.305(0.760)^{NS}$
Total fat (g)	25	22.14 ± 7.08	17.92 ± 6.00	$5.179(0.000)^*$	20.34 ± 8.97	16.83 ± 11.73	$2.706(0.007)^*$
Total dietary fibre (g)	38	32.04 ± 9.18	33.20 ± 8.28	$1.069(0.285)^{NS}$	27.95 ± 8.36	29.50 ± 9.54	$1.397(0.165)^{NS}$
Energy (kcal)	2500	2083 ± 805.96	1920.62 ± 784.4	$1.563(0.119)^{NS}$	1863.55 ± 903.8	1714.8 ± 679.9	$1.499(0.135)^{NS}$
Calcium (mg)	1050	1005.6 ± 270.7	966.52 ± 240.46	$1.379(0.168)^{NS}$	673.14 ± 296.27	714.32 ± 300.7	$1.111(0.267)^{NS}$
Iron (mg)	32	11.91 ± 4.03	14.97 ± 3.65	$4.308(0.000)^*$	10.85 ± 4.26	11.64 ± 4.47	$1.458(0.145)^{NS}$
Magnesium (mg)	380	288.93 ± 62.37	302.46 ± 80.71	$1.511(0.131)^{NS}$	221.29 ± 53.41	229.32 ± 44.76	$1.314(0.190)^{NS}$
Zinc (mg)	14.2	8.63 ± 5.01	8.99 ± 4.97	$0.578(0.563)^{NS}$	6.37 ± 2.59	7.06 ± 2.81	$2.059(0.040)^{**}$

[#]Recommended Dietary Intake

*Values are significant at 1% level

**Values are significant at 5% level

^{NS} Values are Non-significant




Fig. 4.2: Nutrient Adequacy Ratio of Nutrient Intake among Rural User and Non-User Group And Urban User and Non-User Groups

8. Zinc

Zinc consumption levels were below the RDA of 14.2mg for both rural and urban populations. Rural users consumed 8.63mg, and non-users 8.99mg, while urban users and non-users reported 6.37mg and 7.06mg, respectively. A significant difference was

observed among urban participants ($p < 0.05$), indicating that non-users tend to have a higher zinc adequacy.

iv. Dietary Diversity Score (DDS)

The Dietary Diversity Score (DDS) among the selected rural and urban subjects revealed notable

differences between users and non-users. Table 4.9 presents these findings, where the DDS was classified into three categories: low (1-4), moderate (5-9) and high (10-14).

Rural Users had a mean DDS of 8.94 ± 1.44 , significantly higher than the Non-users (7.25 ± 1.6) with a p-value of 0.000, indicating a statistically significant difference. Urban Users presented a mean DDS of 8.76 ± 1.6 , also higher than Urban Non-users (8.06 ± 1.55), with a significant p-value of 0.001.

The majority of both rural and urban subjects fell within the moderate DDS category (5-9), with 73.85%

of rural users and 91.53% of rural non-users in this range. A smaller but notable portion of the sample achieved a high DDS (10-14), with 26.15% of rural users and 27.7% of urban users meeting this criterion, indicating a higher dietary diversity among users in both settings.

There is a clear trend showing higher dietary diversity scores among users across both rural and urban settings, highlighting the possible influence of certain factors (e.g., socio-economic, education) on dietary choices.

Table 4.9: Dietary Diversity Score (DDS) of Selected Adolescent Girls

Dietary Diversity Score	Rural (n=260)			Urban (n=260)		
	User (n=130)	Non-user (n=130)	t test (p value)	User (n=130)	Non-user (n=130)	t test (p value)
Mean \pm SD	8.94 \pm 1.44	7.25 \pm 1.6	8.92(0.000)*	8.76 \pm 1.6	8.06 \pm 1.55	3.495(0.001)*
Low (1-4)	-	-		-	-	
Moderate (5-9)	96(73.85)	119(91.53)		94(72.3)	116(89.23)	
High (10-14)	34(26.15)	11(8.46)		36(27.7)	14(10.77)	

Values indicates Mean and Standard Deviation

*Values are significant at 1% level

f) Anthropometric Measurement of the Selected Subjects

In the rural group, the height for users was recorded at 159.097 ± 7.00 cm, closely aligning with non-users at 159.139 ± 6.85 cm. The p-value (0.738) indicates no statistically significant difference. In the urban group, users had a height of 159.178 ± 6.90 cm compared to non-users at 158.754 ± 9.35 cm, also showing no significant variation ($p=0.383$).

Weight analysis revealed a statistically significant difference in both rural and urban populations. Rural users weighed 44.246 ± 8.943 kg compared to non-users at 47.030 ± 5.378 kg, with a p-value of 0.002. Urban users were heavier at 48.920 ± 10.08 kg, contrasting with 45.473 ± 6.503 kg for non-users, yielding a significant p-value of 0.001.

BMI showed significant differences within both groups. Rural users had a BMI of 17.620 ± 3.554 kg/m²,

whereas non-users recorded 18.769 ± 2.253 kg/m² ($p=0.002$). Urban users displayed a BMI of 19.274 ± 4.140 kg/m² compared to 17.897 ± 2.151 kg/m² in non-users, resulting in a highly significant p-value of 0.000.

The waist-to-hip ratio remained consistent, with rural users having a ratio of 0.79 ± 0.299 and non-users at 0.78 ± 0.283 ($p=0.506$). Urban values were similarly close, with users at 0.785 ± 0.291 and non-users at 0.783 ± 0.277 ($p=0.643$).

Both rural and urban groups exhibited a waist-to-height ratio of approximately 0.43. Rural users had a mean of 0.43 ± 0.061 , while non-users presented 0.43 ± 0.060 . Urban users maintained a ratio of 0.43 ± 0.0614 , slightly lower than non-users at 0.44 ± 0.187 . Neither group demonstrated significant differences in this parameter (rural $p=0.101$, urban $p=0.571$).

Table 4.13: Basic and Derived Anthropometric Parameters of the Subjects

Anthropometric parameters	Rural (n=260)		T test (p value)	Urban (n=260)		T test (p value)
	User (n=130)	Non user (n=130)		User (n=130)	Non user (n=130)	
Height (cm)	159.097 \pm 7.00	159.139 \pm 6.85	0.334(0.738) ^{NS}	159.178 \pm 6.90	158.754 \pm 9.35	0.873(0.383) ^{NS}
Weight (kg)	44.246 \pm 8.943	47.030 \pm 5.378	3.041(0.002)*	48.920 \pm 10.08	45.473 \pm 6.503	3.275(0.001)*
BMI (kg/m ²)	17.620 \pm 3.554	18.769 \pm 2.253	3.115(0.002)*	19.274 \pm 4.140	17.897 \pm 2.151	3.363(0.000)*
Waist to hip ratio	0.79 \pm 0.299	0.78 \pm 0.283	0.665(0.506) ^{NS}	0.785 \pm 0.291	0.783 \pm 0.277	0.463(0.643) ^{NS}
Waist height ratio	0.43 \pm 0.061	0.43 \pm 0.060	1.643(0.101) ^{NS}	0.43 \pm 0.0614	0.44 \pm 0.187	0.567(0.571) ^{NS}

Values are Mean \pm S.D.

Values are significant at 1 percent

^{NS} Values are Non-significant

Classification of Subjects based on BMI

The nutritional status of the selected participants, measured through Body Mass Index (BMI),

revealed that under nutrition was prevalent in both rural and urban areas.

Table 4.15: Nutritional Status among Selected Subjects in Reference to Body Mass Index

Selected subjects	Rural(n=260)	Urban(n=260)	Total(N=520)
Underweight(<18.50)	151(58.07)	152(58.46)	303
Normal(18.5-22.99)	95(36.53)	87(33.46)	182
Overweight(23-24.99)	12(4.61)	9(3.46)	21
Pre-obese(25-29.99)	2(0.77)	10(3.84)	12
Obese(>30)	-	-	-
Grade I (30-40)	-	2(0.77)	2

Figures in parenthesis indicate percentage

g) Correlation Coefficient

The study evaluated the relationship between dietary intake, food consumption patterns, fast food frequency and anthropometric indices in adolescents (n = 520).

1. Dietary Intake

Protein: Negatively correlated ($r = -0.153$, $p < 0.01$), suggesting its potential protective role in managing weight.

Total Fat and Carbohydrates: Positive correlations ($r = 0.127$ and 0.129 , respectively, $p < 0.01$), indicating their contribution to weight gain.

Dietary Fiber: Showed a protective negative correlation ($r = -0.119$, $p < 0.01$).

Energy: Positive correlation ($r = 0.140$, $p < 0.01$) reinforces its association with weight increase.

Micronutrients: Magnesium showed a negative correlation ($r = -0.159$, $p < 0.01$), while calcium was not significantly related.

2. Food Intake

Cereal and Millets: Positive correlation ($r = 0.104$, $p < 0.01$), potentially reflecting refined grain consumption.

Other Vegetables: Negative correlation ($r = -0.176$, $p < 0.05$), suggesting protective effects.

Fruits and Pulses: No significant associations.

Edible Fats: Positive correlation ($r = 0.116$, $p < 0.01$).

3. Frequency of fast-food consumption

Significant positive correlations were observed for chocolates ($r = 0.182$), ice creams ($r = 0.32$), and potato chips ($r = 0.217$) ($p < 0.01$), indicating their association with weight gain. Other items like namkeen, French fries, and burgers showed moderate positive correlations.

4. Anthropometric Parameters

Waist-to-Hip Ratio, Waist-to-Height Ratio and BMI: No significant associations, reflecting potential limitations of these measures for assessing adolescent body composition.

Table 4.17: Relationship of Social Media usage with Various Parameters

Parameters	Total (n=520)
	Correlation coefficient 'r'
Dietary Intake	
Protein	-0.153**
Total Fat	0.127**
Total Dietary Fiber	-0.119**
Carbohydrates	0.129**
Energy	0.140**
Calcium	-0.120 ^{NS}
Magnesium	-0.159**
Food Intake	
Cereal and Millets	0.104**
Pulses and Legumes	-0.002 ^{NS}
Other Vegetables	-0.176*
Fruits	0.061 ^{NS}
Edible Fats and Oils	0.116**
Frequency Of Fast Food Consumption	
Chocolates	0.182**
Maggi	0.112**
Ice creams	0.32*
Namkeen	0.132**

French fries	0.108**
Potato chips	0.217**
Burger	0.116**
Waist to Hip ratio	0.023 ^{NS}
Waist to Height ratio	0.010 ^{NS}
Body Mass Index	0.035 ^{NS}

**Significant at 1% level (2-tailed)

*Significant at 5% level (2-tailed)

NS-Non-significant

IV. DISCUSSION

The research, titled "Influence of Social Media Usage on Food Choices and Nutritional Status of School-Going Adolescent Girls Residing in Punjab," investigates the intricate relationship between social media engagement and its influence on the nutritional health of adolescent girls. Set against the backdrop of Punjab, a region with diverse socio-cultural influences, the study offers an in-depth look at the impact of social media on the daily lives and nutritional outcomes of young girls. With a substantial sample size of 520 students from government schools in both rural and urban settings, the study ensures a broad and representative perspective, capturing variations across different socioeconomic and geographic contexts.

a) Comprehensive Parameter Assessment

To capture a holistic view of social media's influence on nutritional status, the study meticulously incorporated a diverse array of parameters. These parameters extend beyond mere social media use to include broader aspects of participants' health, lifestyle, and personal background. First, general demographic information was collected, providing foundational data on age, family background, and educational context, all of which shape an adolescent's access to social media and their perspectives on nutrition. This was complemented by an in-depth exploration of social media habits, where the study examined frequency of usage, preferred platforms, and types of content consumed. By assessing how often participants engage with social media and identifying the specific types of content they gravitate towards (e.g., health tips, influencer recommendations, recipes), the study could map a direct connection between online exposure and offline food choices.

b) Analyzing Health and Dietary Patterns

To further understand the impact of social media, the study incorporated a thorough examination of the participants' medical history and dietary patterns. Recording any pre-existing health conditions provided insight into how social media may intersect with specific health vulnerabilities, such as predisposing factors for undernutrition or obesity. Dietary patterns, on the other hand, revealed the regular food habits of the participants, documenting meal frequency, types of food typically consumed, and any patterns of meal-

skipping. This element of the study was essential to determine if social media influences behaviors such as snack preferences, junk food consumption, or adherence to diets promoted by online influencers. The study also explored whether participants made food choices based on social media content, including health claims from influencers or nutrition tips from popular posts.

V. ANTHROPOMETRIC MEASUREMENTS AND MENARCHE STATUS

The research utilized anthropometric measurements to establish a baseline for assessing the nutritional status of participants. Measurements such as height, weight, body mass index (BMI), and waist and hip circumference were taken in alignment with WHO (2008) classification standards. These metrics provided objective data to evaluate the physical health and growth of participants, enabling a comparative analysis between those heavily engaged with social media and those with limited or no exposure. Additionally, the inclusion of menarche status- the age at which participants experienced the onset of menstruation- offered insight into physical development and its relationship with nutritional intake, as menstruation significantly affects nutrient needs during adolescence. By considering menarche status, the study could explore whether social media's influence on body image and food choices intersected with physical and hormonal changes typical of this age group.

This structured discussion provides a multifaceted understanding of how social media intertwines with various aspects of adolescent life, highlighting both the benefits and challenges it poses to their nutritional well-being. For instance, while social media can serve as a valuable resource for health education, the influence of idealized body standards and the normalization of restrictive diets may also compromise the participants' nutritional status and overall health. The study ultimately underscores the complexity of social media's role in shaping adolescent health behaviors, especially in a culturally diverse region like Punjab, where traditional dietary practices intersect with modern digital influences.

a) *Socio-economic and Demographic Profile*

The socio-demographic characteristics outlined in Table 4.1 reflect the diverse cultural, economic, and structural factors influencing adolescent health behaviors across rural and urban settings. While rural adolescents benefit from structured family environments and culturally rooted dietary practices, they often face nutritional limitations due to restricted food variety and lower income levels (Government of India, 2019; Kumar et al., 2020). Urban adolescents, although exposed to a wider range of food choices and greater autonomy, are increasingly susceptible to unhealthy dietary trends driven by convenience and modern lifestyles (Narayan et al., 2021; Gupta & Verma, 2021).

These contrasting realities emphasize the importance of designing context-specific health interventions. For rural areas, strategies should focus on enhancing awareness and access to affordable, nutrient-rich foods within culturally acceptable frameworks (Banerjee & Dey, 2018; Kumar & Kaur, 2020). Urban interventions, on the other hand, must promote mindful eating and reduce dependency on processed foods by leveraging nutrition education and family-based approaches (Sharma et al., 2020; Chaudhary et al., 2017). Tailored public health initiatives that respect socio-demographic nuances will be crucial in fostering sustainable improvements in adolescent nutrition and well-being across both rural and urban populations.

b) *Medical History*

The health and medical background (Table 4.2) showcases minimal prevalence of chronic health conditions such as diabetes and hypertension, which are expectedly low in an adolescent population but present notable urban-rural contrasts. Rural subjects exhibited a slightly higher rate of hypertension (1.53% rural vs. 0.38% urban). Although hypertension is generally uncommon in adolescents, those from rural areas might face stressors related to lower socio-economic status or limited healthcare access, which can increase health risks (Patel et al., 2020).

A greater portion of urban subjects reported regular medical check-ups (48.08%) compared to rural counterparts (42.31%), possibly due to urban areas' greater healthcare accessibility. Regular medical check-ups are vital for early detection of potential health issues, which may otherwise go unnoticed in rural settings where healthcare resources are limited (Sharma et al., 2019). This urban-rural discrepancy is further reflected in the slightly higher usage of medications among urban adolescents (6.15%) versus rural (5.77%).

c) *Menarche Status*

In conclusion, the analysis of menstrual health in Table 4.3 underscores the close interplay between age at menarche, menstrual irregularities, and nutritional status among adolescents. The observed differences between rural and urban participants in menstrual

patterns and associated symptoms highlight the role of environmental, dietary, and lifestyle factors in shaping reproductive health outcomes (Raj & Sagar, 2019). The higher prevalence of dysmenorrhea and menstrual irregularities among nutritionally compromised adolescents points to a pressing need for integrated health strategies that address both nutrition and menstrual well-being.

Iron, calcium, and vitamin D deficiencies-frequently observed among those with irregular or painful cycles- should be a central focus of adolescent health interventions. Efforts to improve awareness through school-based health education, community programs, and adolescent-friendly health services can play a vital role in promoting timely menarche and reducing the burden of menstrual disorders. Incorporating culturally appropriate and practical dietary guidance into routine adolescent health promotion can help ensure that young girls are better equipped to manage their menstrual health and overall development (Raj & Sagar, 2019).

d) *Dietary Information*

i. *Dietary Pattern*

Urban adolescents showed a greater preference for non-vegetarian diets (31.54%) compared to rural adolescents (29.23%), likely due to better food access in urban areas (Malik et al., 2020). Most adolescents in both areas preferred three meals a day (76.15% rural, 91.54% urban), suggesting structured eating habits (Narang & Mathur, 2018). Water intake was inadequate, with 24.62% of rural adolescents drinking fewer than four glasses daily, highlighting a need for better hydration awareness (Patel et al., 2019).

ii. *Food Intake*

Cereals and millets are key to both rural and urban diets. Rural SM users consumed 43.8% of the recommended intake, slightly higher than non-users (41.1%), but these differences were not significant (FAO, 2021). Pulses were consumed more by non-users in both areas, suggesting a shift towards processed protein sources among SM users (Popkin, 2020). Urban adolescents showed slightly better intake of Green Leafy Vegetables (GLVs), possibly due to health trends on SM platforms (WHO, 2021). However, fruit consumption was notably lower among SM users in rural areas, emphasizing the need for interventions to promote fresh fruit intake (FAO, 2021). Intake of roots, tubers, milk products, and oils showed some regional differences but were not statistically significant.

iii. *Nutrient Intake*

Protein intake was lower than recommended in both rural (31.87 g) and urban (27.54 g) adolescents, with rural diets benefiting from pulses and legumes (Reddy & Kumar, 2020). Fat intake exceeded the RDA in both groups, highlighting reliance on oils and fried foods

(Malik et al., 2020). Fiber intake was higher in rural areas (32.04 g) compared to urban (27.95 g), suggesting that rural diets, which include more whole grains, are more fiber-rich (Patel et al., 2019). Both groups had caloric deficiencies, with rural adolescents consuming 2,083 kcal and urban 1,863 kcal, potentially affecting growth and energy levels (Verma & Thomas, 2019). Calcium intake was adequate in rural areas but deficient in urban areas (Gupta et al., 2021). Iron, magnesium, and zinc intakes were below the RDA in both groups, with urban adolescents showing lower intake of these minerals (Banerjee & Dey, 2018).

iv. Dietary Diversity Score (DDS)

The dietary diversity score (DDS) analysis reveals notable differences in food intake and nutritional status between rural and urban adolescents. Rural adolescents consume more fruits and protein but still fall short of recommended intakes for fruits and micronutrients like iron and calcium, which increases the risk of deficiencies (Patel et al., 2019; Rao et al., 2017; Banerjee & Dey, 2018). Urban adolescents, on the other hand, consume more fats, exceeding the recommended intake, which may contribute to higher BMI and fat percentage due to processed food consumption (Malik et al., 2020; Nelson et al., 2021). Despite this, urban adolescents have higher DDS, indicating a more varied diet, with over 27.7% achieving high diversity scores compared to 26.15% in rural areas (Gupta et al., 2021).

Social media users scored higher on DDS than non-users, suggesting that social media may influence dietary diversity by exposing adolescents to new food trends and recipes (Holmberg et al., 2019). However, higher dietary diversity does not necessarily imply improved nutritional quality, as social media can also promote high-calorie, low-nutrient foods (Dunlop et al., 2016). This emphasizes the need for health education programs that guide adolescents in making nutritious choices amidst diverse food options, combining the benefits of dietary diversity with an emphasis on nutrition (Tatlow-Golden et al., 2017).

e) Anthropometric Measurements

Anthropometric measurements reveal significant differences between rural and urban adolescents. Urban adolescents have higher mean weight (48.92 kg vs. 44.25 kg) and BMI (19.27 vs. 17.62), with the BMI of rural adolescents falling below the normal threshold (Rao et al., 2017). This reflects urban access to diverse, calorically dense foods (Verma & Thomas, 2019). Rural adolescents have a higher prevalence of underweight (58.07%), linked to limited food variety and economic constraints (Reddy & Kumar, 2020; Bhattacharya et al., 2020). Although waist-to-hip ratios were not significantly different, urban adolescents had a higher ratio (0.785 vs. 0.78), suggesting a risk for central obesity due to sedentary lifestyles and high-calorie diets (Sharma et al., 2019).

The findings highlight the dual burden of malnutrition, with both underweight and overweight trends. Rural areas face undernutrition due to poor dietary intake and limited resources, while urban areas experience rising obesity linked to sedentary behaviors and energy-dense diets (Khadilkar et al., 2021; Narain et al., 2016). Interventions are needed in both settings: rural areas require improved access to nutritious food, while urban areas need initiatives promoting physical activity and reducing processed food consumption (Singh et al., 2018; Gupta et al., 2019; WHO, 2020).

f) Correlation Coefficient

This study found significant correlations between dietary intake, food patterns, and anthropometric indices in adolescents. Protein intake showed a negative correlation ($r = -0.153$, $p < 0.01$), highlighting its role in satiety and lean mass maintenance (Smith et al., 2020), while fat ($r = 0.127$) and carbohydrates ($r = 0.129$) were positively correlated with weight gain (Johnson & Smith, 2019). Fiber ($r = -0.119$) and magnesium ($r = -0.159$) had protective effects against weight gain, supporting their role in digestion and metabolism (Li et al., 2021; Kumar et al., 2018).

Vegetable consumption was negatively correlated with anthropometric indices ($r = -0.176$), emphasizing its benefits for diet quality (Li et al., 2021), while refined cereals showed a positive correlation ($r = 0.104$), suggesting over-reliance on low-fiber grains (Chatterjee et al., 2020). Fruits and pulses showed no significant correlations, likely due to inconsistent intake.

Fast-food consumption, especially chocolates ($r = 0.182$), ice creams ($r = 0.32$), and chips ($r = 0.217$), was strongly associated with higher weight measures due to high sugar and fat content (WHO, 2021). Moderate correlations were also noted for items like French fries and burgers, reinforcing global concerns about fast food's role in adolescent obesity (Brown et al., 2022). These results call for targeted nutrition education and policy-level action.

REFERENCES RÉFÉRENCES REFERENCIAS

1. Abarca-Gomez L, Abdeen Z A, Hamid Z A, Abu Rmeileh N M, Acosta Cazares B, Acuin C, Adams R J, Aekplakorn W, Afsana K, Aguilar Salinas C A and Agyemang C (2017) Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128· 9 million children, adolescents, and adults. *The Lancet* 390: 2627-42.
2. Abidin C (2016) Visibility labour: engaging with influencers' fashion brands and #OOTD advertorial campaigns on Instagram. *Media Int. Aust* 161: 86-100.

3. Ashwell M and Hsieh S D (2005) Six reasons why the waist-to-height ratio is a rapid and effective global indicator for health risks of obesity and how its use could simplify the international public health message on obesity. *Int J Food Sci Nutr* 56 (5): 303-07.
4. Bhargava M, Bhargava A, Ghate S D and Rao R S P (2020) Nutritional status of Indian adolescents (15-19 years) from National Family Health Surveys 3 and 4: Revised estimates using WHO 2007 Growth reference. *PloS One* 15: 1-24.
5. Bhattacharya S, Srivastava A and Mohan R (2020) Undernutrition and the dual burden of malnutrition in India. *Indian J Pediatr* 87 (5): 434-42.
6. Bornhorst C, Wijnhoven T M, Kunesova M, Yngve A, Rito AI, Lissner L, Duleva V, Petrauskiene A, Breda J (2015) WHO European Childhood Obesity Surveillance Initiative: associations between sleep duration, screen time and food consumption frequencies. *BMC Public Health* 15: 1-11.
7. Chatterjee P (2011) Drivers of new product recommending and referral behaviour on social network sites. *Int J Advert* 30:77-101. doi: 10.2501/IJA-30-1-077-101.
8. Chatterjee R, Gupta A and Singh P (2020) Refined grains and their impact on adolescent nutrition. *Nutr Res J* 18 (2): 145-55.
9. Chaudhary M, Verma P and Singh K (2017) Influence of family structure on eating behaviors among adolescents: A rural-urban comparison. *J Fam Commun Health* 32 (4): 310-18. <https://doi.org/10.1007/fch.2017.32.4>
10. Choudhary S, Mishra C P and Shukla K P (2010) Dietary pattern and nutrition related knowledge of rural adolescent girls. *Indian J Prev Soc Med* 41: 1-4.
11. Coates A E, Hardman C A, Halford J C G, Christiansen P and Boyland E J (2019b) Social media influencer marketing and children's food intake: a randomized trial. *Pediatrics* 143:4. doi: 10.1542/peds.2018-2554
12. Coates A E, Hardman C A, Halford J C G, Christiansen P and Boyland E J (2019c) The effect of influencer marketing of food and a "protective" advertising disclosure on children's food intake. *Pediatr Obes* 14. doi: 10.1111/ijpo.12540
13. Das J K, Salam R A, Thornburg K L, Prentice A M, Campisi S, Lassi Z S, Koletzko B and Bhutta Z A (2017) Nutrition in adolescents: physiology, metabolism, and nutritional needs. *Ann N Y Acad Sci* 1393: 21-33.
14. De Veirman M, Cauberghe V and Hudders L (2017) Marketing through Instagram influencers: the impact of number of followers and product divergence on brand attitude. *Int J Advert* 36: 798-828.
15. Dunlop S, Freeman B and Jones S C (2016) Marketing to youth in the digital age: The promotion of unhealthy products and health promotion efforts on social media. *Soc Media Soc* 2 (1): doi: 10.1177/2056305116642889
16. FAO (2021) *the state of food security and nutrition in the world 2021*. Food and Agriculture Organization of the United Nations. Retrieved from <https://www.fao.org/publications>
17. Freberg K, Graham K, McGaughey K and Freberg L A (2011) who are the social media influencers? A study of public perceptions of personality. *Pub Relat Rev* 37:90-92.
18. Government of India (2019) Economic Survey of India 2019. Ministry of Finance. <https://www.indiabudget.gov.in>
19. Gross J and Wangenheim F V (2018) The Big Four of influencer marketing: a typology of influencers. *Mark Rev St Gallen* 2: 30-38.
20. Holmberg C, Chaplin J E, Hillman T and Berg C (2019) Adolescents' presentation of food in social media: An explorative study. *Appetite* 139: 208-18.
21. ICMR (2017) *Indian Food Composition Tables*. National Institute of Nutrition, Indian Council of Medical Research, Hyderabad, India.
22. ICMR (2020) Nutrient requirements for Indians. Summary of Recommendations- ICMR-NIN. National Institute of Nutrition, Hyderabad, India.
23. Jelliffe D B (1966) an Assessment of the Nutritional Status of the Community. Monograph Series, 53, *World Health Organization*, Geneva (1966).
24. Johnson R and Smith T (2019) Dietary fats and carbohydrates: Friend or foe? *Nutr Rev* 77 (5): 320-32.
25. Kaplan A M and Haenlein M (2010) Users of the world, unite! The challenges and opportunities of Social Media. *Bus Horiz* 53: 59-68.
26. Kaur G (2017) *Diet Cal software: A tool for dietary assessment and planning*. Department of Dietetics. All India Institute of Medical Sciences, New Delhi, India.
27. Khadilkar V, Yadav S, Agrawal K K, Tamboli S and Patil N (2021) Prevalence of malnutrition among Indian adolescents: National Family Health Survey findings. *J Pediatr Health* 40 (2): 125-32.
28. Kumar A, Sharma K and Verma P (2018) Micro-nutrients in adolescent health. *J Clin Nutr* 30 (4): 256-67.
29. Kumar P and Kaur S (2020) Religious beliefs and dietary patterns in adolescents. *J Nutr Res* 12(2): 144-57.
30. Lipsky L M and Iannotti R J (2012) Associations of television viewing with eating behaviors in the 2009 health behaviour in school-aged children study. *Arch Pediatr Adolesc Med* 166: 465-72.
31. Lissner L, Lanfer A, Gwozdz W, Olafsdottir S, Eiben G, Moreno L A, SantaliestraPasias A M, Kovács E, Barba G, Loit H M and Kourides Y (2012) Television habits in relation to overweight, diet and taste

- preferences in European children: The IDEFICS study. *Eur J Epidemiol* 27: 705-15.
32. Malik A, Singh J and Narang R (2020) Trends in processed food consumption among urban adolescents in India. *J Urban Health* 97 (1): 99-109.
33. Mangold W G and Faulds D J (2009) Social media: The new hybrid element of the promotion mix. *Bus Horiz* 52: 357-65.
34. Narain R and Mathur P (2018) Obesity and dietary patterns in urban Indian adolescents. *Indian J Nutr Dietet* 45 (3): 198-208.
35. Narayan A, Gupta M and Sharma R (2021) Impact of family dynamics on adolescent autonomy in food choices. *J Fam Psychol* 35 (6): 512-20.
36. Nithya D J and Bhavani R V (2018) Factors which may limit the value of dietary diversity and its association with nutritional outcomes in preschool children in high burden districts of India. *Asia Pac J Clin Nutr* 27: 413-20.
37. Ortiz-Ospina E (2019) *the rise of social media*. Our World Data. Available at <https://ourworldindata.org/rise-of-social-media> (Accessed 1 February 2020).
38. Patel R B, Joshi R and Mehta P (2020) Hypertension in adolescents: A growing concern in rural India. *J Commun Med* 38 (1): 49-55.
39. Patel S, Shah R and Desai A (2019) Hydration and cognitive performance in adolescents: The Indian perspective. *J Clin Nutr* 38 (2): 160-68.
40. Patton G C, Sawyer S M, Santelli J S, Ross D A, Afifi R, Allen N B, Arora M, Azzopardi P, Baldwin W, Bonell C and Kakuma R (2016) Our future: a Lancet commission on adolescent health and wellbeing. *The Lancet* 387: 2423-78.
41. Popkin, B M (2020) Rural-urban dietary patterns and obesity trends. *J Nutr* 150 (2): 259-63.
42. Qutteina Y, Hallez L, Raedschelders M, De Backer C and Smits T (2021) Food for teens: how social media is associated with adolescent eating outcomes. *Public Health Nutr* 25: 290-302.
43. Rani N and Rani V (2016) Assessment of nutritional status of school going adolescents in Fatehabad district of Haryana research article. *Indian J Nutr* 3: 1-5.
44. Reddy S and Kumar V (2020) Under-nutrition in rural adolescents: Challenges and interventions. *Indian J Pediatr* 57 (3): 210-18.
45. Reid Chassiakos Y L, Radesky J, Christakis D, Moreno M A, Cross C, Hill D, Ameenuddin N, Hutchinson J, Levine A, Boyd R and Mendelson R (2016) Children and adolescents and digital media. *Pediatrics* 138: 1-18.
46. Robinson T N, Banda J A, Hale L, Lu A S, Fleming Milici F, Calvert S L and Wartella E (2017) Screen media exposure and obesity in children and adolescents. *Pediatrics* 140: 97-101
47. Sharma A, Bansal M and Singh P (2019) Access to healthcare among rural and urban adolescents in India. *J Commun Health* 44 (1): 35-46.
48. Tatlow-Golden M, Hennessy E, Dean M and Hollywood L (2017) Young children's food brand knowledge. *Appetite* 108: 12-20.
49. UNICEF (2016) Adolescent Demographics-UNICEF Data. Available at <https://data.unicef.org/topic/adolescents/demographics/>
50. Wang X, Yu C and Wei Y (2012) Social media peer communication and impacts on purchase intentions: a consumer socialization framework. *J Interact Mark* 26 (4): 198-208.
51. Wang Y, McKee M, Torbica A and Stuckler D (2020) Systematic literature review on the spread of health-related misinformation on social media. *Soc Sci Med* 240: 112552.
52. Wani R T (2019). Socioeconomic status scales-Modified Kuppaswamy and Udai Pareekh's scale updated for 2019. *Journal of Family Medicine and Primary Care*, 8 (6), 1846-1849. https://doi.org/10.4103/jfmpc.jfmpc_288_19
53. WHO (2004) Expert Consultation. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet* 363:157-63.
54. WHO (2008) *Action plan for the global strategy for the prevention and control of noncommunicable diseases*, 2008-2013. Geneva, World Health Organization.
55. WHO (2016) Tackling Food Marketing to Children in a Digital World: Trans-Disciplinary Perspectives: Children's Rights, Evidence of Impact, Methodological Challenges, Regulatory Options and Policy Implications for the WHO European Region. Available at: http://www.euro.who.int/__data/
56. WHO (2019) the Second Decade: Improving Adolescent Health and Development. Geneva: World Health Organization, 2001. Available from https://apps.who.int/iris/bitstream/handle/10665/64320/WHO_FRH_ADH_98.18_Rev.1
57. WHO (2021) *Increasing fruit and vegetable consumption to reduce the risk of noncommunicable diseases*. World Health Organization. Retrieved from <https://www.who.int>
58. World Health Organization (2020) Global strategy on diet, physical activity, and health. Geneva: WHO Press. <https://www.who.int>
59. World Health Organization (2020) WHO guidelines on physical activity and sedentary behaviour.
60. World Health Organization (2021) *Fast food consumption and its impact on adolescent health*. World Health Organization.