



GLOBAL JOURNAL OF MEDICAL RESEARCH: I
SURGERIES AND CARDIOVASCULAR SYSTEM
Volume 14 Issue 5 Version 1.0 Year 2014
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Triple Vessel Coronary Artery Disease in Young Female

By Ali Razaghani & Hafeez-Ul-Hassanvirk

Thomas Jefferson University, United States

Background- Although coronary heart disease (CHD) primarily occurs in patients over the age of 40, younger men and women can be involved. Majority of studies have used an age cut-off of 40 to 45 years to define "young" patients with CHD or acute myocardial infarction (MI). The same age definition will be used in this article. The prevalence of CHD in younger subjects is difficult to establish accurately since it is frequently a silent process. Acute Myocardial infarction in young females is an uncommon occurrence and even if we see cases, very few of them have shown to have greater than one vessel coronary artery disease. When a young female present with acute MI, the presentation is very vague and can be easily missed so, presence or absence of cardiovascular risk factors regardless of age should be the key factor in making a decision to perform coronary angiography and full cardiovascular workup. We report here 31 year old female with multiple cardiovascular risk factors who presented with an atypical chest pain with normal EKG in emergency room and was ultimately diagnosed with triple vessel coronary artery disease. In this paper we will describe a case to describe the importance of early coronary angiography and cardiovascular workup in presence of significant risk factors despite atypical presentation and younger age of patient.

GJMR-I Classification: NLMC Code: WG 595



Strictly as per the compliance and regulations of:



Triple Vessel Coronary Artery Disease in Young Female

Ali Razaghani ^α & Hafeez-Ul-Hassanvirk ^σ

I. BACKGROUND

Although coronary heart disease (CHD) primarily occurs in patients over the age of 40, younger men and women can be involved. Majority of studies have used an age cut-off of 40 to 45 years to define "young" patients with CHD or acute myocardial infarction (MI). The same age definition will be used in this article.

The prevalence of CHD in younger subjects is difficult to establish accurately since it is frequently a silent process.

Acute Myocardial infarction in young females is an uncommon occurrence and even if we see cases, very few of them have shown to have greater than one vessel coronary artery disease. When a young female present with acute MI, the presentation is very vague and can be easily missed so, presence or absence of cardiovascular risk factors regardless of age should be the key factor in making a decision to perform coronary angiography and full cardiovascular workup. We report here 31 year old female with multiple cardiovascular risk factors who presented with an atypical chest pain with normal EKG in emergency room and was ultimately diagnosed with triple vessel coronary artery disease. In this paper we will describe a case to describe the importance of early coronary angiography and cardiovascular workup in presence of significant risk factors despite atypical presentation and younger age of patient.

Conclusion:

II. CASE REPORT

a) Presenting Complaint

31 Y O F p/w chest pain for 2 weeks.

b) History of Present Illness

31 year old Female with PMH of HTN, DM 2(on insulin), HLD, glaucoma, Major depression, Asthma was in her usual state of health 2 week ago when she started having chest pain. It was mid sternal, 10/10, intermittent, sharp/stabbing, and increased by lying down, improved by sitting associated with nausea but no vomiting. She woke up multiple times at night due to this pain. She preferred to sleep in front of fan due to dyspnea at night.

She also had shortness of breath for couple of months on exertion. She has been recently using two pillows for sleep. She used to walk 4 blocks before getting short of breath (for years) but recently her exercise tolerance has decreased to half a block. She denies any palpitations, any previous history of such pain. Due to this pain, she went to the PMD who called 911 and sent her to ED.

III. ED COURSE

By EMS, she received Aspirin 162mg po once. She received tylenol 650mg po once in ED.

ED VS: afebrile; HR 105/min; BP 146/81mmHg; RR 20/min; O2S 99% RA.

She is not compliant with the medications at home. She has a HHA 5days a week from 8am to 1pm. She has recently gained >20lbs in last year,

Past Medical History: Admitted in hospital multiple times for DKA.

Allergy: NKDA.

Hospitalization: multiple times for DKA.

GI: Not significant.

Family History: hypertension in sister, no diabetes in family.

Social History: HHA, quit smoking 2 years ago, nonalco- holic, no illicit drugs.

Gyne-Obstetrics: menses regula, not pregnant.

Allergies:

Home Medications: Lantus insulin q12h, Humalog with meals, Zyrtec 10mg daily, Simvastatin 20mg daily, Tricor 160mg daily, Cozaar 25mg daily, Vitamin D, Os-Ca.

Vitals: O2 Saturation: 98, Pulse Rate: 73*, Respirations: 17*, BP Position: Lying*, Systolic BP: 124*, Diastolic BP: 72*, Pain Level: 0*, POC Blood Glucose: 109, Temperature (F): 99.3*, Body Mass Index: 37.55.

Physical E

General Exam: She was lying in bed, comfortably, fully oriented.

Head Exam: NC/AT.

Eye Exam: PEERLA, EOMI intact.

Neck Exam: supple, no JVD.

Respiratory Exam: CTA BL, no wheeze.

Cardiac Exam: S1 S2 heard, regular, tachycardiac.

Gastrointestinal Exam: ND/NT, BS+

Author α σ: 39 Brandon road upper Darby, PA 19082. United States.
e-mails: alighani152@gmail.com, hafeezvirkmd@gmail.com

Extremities Exam: palpable pulses, no cyanosis, minimal edema.

Neurological Exam: AAOx3

IV. RESULTS

Initial Labs:

Normal

Cardiac Enzymes:

Troponin I: 2.6-->2.7-->2.8

CK-MB: 4.4→3.2→3.0

Cholesterol, Total	287 MG/DL
Cholesterol, HDL	48 MG/DL
Cholesterol, LDL (Calculated)	SEE TEXT
Cholesterol/HDL Ratio	6.0
Triglycerides	611 MG/DL

Diagnostic studies: TTE

Left Ventricle: The left ventricular cavity size is normal. Left ventricular wall motion is normal. Visually estimated left ventricular ejection fraction is 60%.

Right Ventricle: Normal right ventricular size and function.

Left Atrium: Normal left atrial size.

Right Atrium: Normal right atrial size.

Mitral Valve: The Doppler (color and spectral) study shows trivial mitral regurgitation.

AV and Aortic Root: The Doppler (color and spectral) study shows trivial aortic regurgitation.

Tricuspid Valve: The Doppler (color and spectral) study shows trivial tricuspid regurgitation. As assessed from the tricuspid regurgitant jet, the pulmonary artery systolic pressure is normal.

pulmonic valve: Structurally normal pulmonic valve without stenosis or regurgitation.

Aortic root: Normal aortic root.

V. CONCLUSIONS

Visually estimated left ventricular ejection fraction is 60%.

-Echo showed 60% EF

-Catheterization was done and showed 3V CAD

a) Plan

-start on heparin gtt, stop it 3 hrs before CABG

-start metoprolol 25mg bid po

-CABG tomorrow

-NPO after midnight

-stop aspirin/plavix for CABG

Endocrinology:

#DM 2

-IDDM for 20 yrs

-on insulin lantus 60U BID, 30U TID

plan:

-c/w insulin

#Asthma: well controlled last attack this winter family history+

There are also limited data on the frequency of MI in younger subjects. In the Framingham Heart Study, the incidence of an MI over a 10-year follow-up was 12.9/1000 in men 30 to 34 years old and 5.2/1000 in women 35 to 44 years old [2]. The incidence of MI was eight to nine times greater in men and women aged 55 to 64 years. In other studies, 4 to 10 percent of patients with MI were ≤ 40 or 45 years of age [3-5]. In two series of patients with CHD at ≤ 40 years of age, women comprised 5.6 and 11.4 percent of patients [3,6].

Although CHD is an uncommon entity in young patients, it constitutes an important problem for the patient and the treating physician because of the devastating effect of this disease on the more active lifestyle of young patients. In addition, these patients have different risk factor profiles, clinical presentations, and prognoses than older patients. All of these factors should be taken into consideration when treating young patients with CHD

VI. CORONARY RISK FACTORS

The relative importance of risk factors for the development of CHD according to age was evaluated in a report in which 11,016 men aged 18 to 39 years were followed for 20 years [7]. The relative risks associated with the traditional risk factors were of similar magnitude as in a group of 8955 men aged 40 to 59 years. These included:

- Age — relative risk 1.63 per six year increase
- Serum cholesterol — relative risk 1.92 per 40 mg/dL [1.04 mmol/L] increase
- Systolic blood pressure — relative risk 1.32 per 20 mmHg increase
- Cigarette smoking — relative risk 1.36 per 10 cigarette/day increase

Young patients with MI usually have multiple risk factors for CHD. In some studies, for example, as many as 90 to 97 percent had one or more traditional risk factors for atherosclerosis [8-10]. In a prospective study of over 7000 women of mean age 27 years at baseline followed for an average of 31 years, there were 47 CHD deaths [11]. The CHD mortality rates for those with no risk factors, only one risk factor, or two or more risk factors were 0.7, 2.4, and 5.4 per 1000 person-years, respectively. A comparable relationship was seen for cardiovascular disease mortality and for all-cause mortality. (See "Overview of the risk equivalents and established risk factors for cardiovascular disease".)

a) Smoking

Cigarette smoking is the most common and most modifiable risk factor in young patients. It has been noted in 65 to 92 percent of young patients with MI,

compared to 24 to 56 percent of patients older than 45 years of age [6,9,12-16]. (See "Cardiovascular risk of smoking and benefits of smoking cessation".)

b) *Family history*

Younger patients with CHD more often have a family history of premature CHD: 41 compared to 28 and 12 percent in middle aged or elderly patients, respectively [9]; and 57 versus 43 percent in two series [12]. A higher incidence of a positive family history in young patients (64 percent) was noted in the largest report of 823 patients [6].

In addition, the offspring of patients with premature CHD are more likely to have coronary risk factors than those without such a family history [17]. These include excess body weight and higher levels of serum cholesterol, glucose, and insulin. These offspring are also more likely to have evidence of vascular disease such as endothelial dysfunction and increased carotid artery intima-media thickness [18].

The association between family history and premature CHD can be due to both genetic and environmental factors. This was addressed in a study of 398 families in which 62 vascular biology genes were evaluated [19]. Missense variants of several thrombospondin genes were significantly associated with MI and CHD.

c) *Lipid abnormalities*

Hypercholesterolemia is common in young patients with CHD, but its prevalence is similar to that in older patients. However, when compared to older patients, young patients have lower mean serum high density lipoprotein (HDL) concentrations (35 versus 43 mg/dL [0.9 versus 1.1 mmol/L]) and higher serum triglycerides (239 versus 186 mg/dL [2.7 versus 2.1 mmol/L]) [15]. (See "HDL metabolism and approach to the patient with abnormal HDL-cholesterol levels".)

Hypertriglyceridemia was, in one series, the most common lipid abnormality in young patients with MI [20]. It may be associated with glucose intolerance and a predominance of small atherogenic LDL particles, both of which predispose to atherosclerosis. (See "Approach to the patient with hypertriglyceridemia".)

d) *Diabetes and hypertension*

Two other important coronary risk factors, diabetes mellitus and hypertension, appear to be less common in young patients with CHD than in older patients [6,12]. However, young patients frequently have subtle problems with glucose metabolism. In one study of 108 patients without a history of diabetes mellitus who had an MI before the age of 45, 65 percent had decreased oral glucose tolerance and a hyperinsulinemic response to oral glucose challenge [20]. This finding is consistent with other observations

that impaired glucose tolerance in the absence of overt diabetes is a risk factor for coronary disease. (See "Prevalence of and risk factors for coronary heart disease in diabetes mellitus", section on 'CHD before diabetes'.)

e) *Obesity*

Obesity appears to be an independent risk factor for coronary atherosclerosis, at least in young men. This was illustrated in an autopsy study of approximately 3000 persons between the ages of 15 and 34 who died from noncardiac causes [21]. Increasing body mass index was associated with both fatty streaks and raised atherosclerotic lesions in the right coronary and left anterior descending coronary arteries in young men, but not young women. The effect of obesity on other risk factors (eg, lipid abnormalities, hypertension, glucose intolerance) accounted for only about 15 percent of the relationship between obesity and coronary atherosclerosis.

How this might occur is not known, but other studies have noted an apparently independent effect of obesity as an important coronary risk factor. A report from the Framingham Heart Study suggested that obesity in middle-aged subjects could account for as much as 23 percent of cases of CHD in men and 15 percent in women

f) *Other factors*

A variety of other possible contributing factors have been identified in young patients with MI. These include:

- Oral contraceptive use in young women, primarily when combined with heavy smoking [25]. (See "Risks and side effects associated with estrogen-progestin contraceptives".)
- Frequent cocaine use, which, in the Third National Health and Nutrition Examination Survey of 10,085 adults between the ages of 18 and 45, accounted for 25 percent of nonfatal MIs [26]. (See "Evaluation and management of the cardiovascular complications of cocaine abuse", section on 'Myocardial infarction'.)
- Smoking marijuana may be a rare trigger of MI [27]. (See "Cannabis use disorder: Treatment, prognosis, and long-term medical effects".)
- Factor V Leiden, which is inactivated less efficiently by activated protein C than wild-type factor V, leads to a procoagulant state by increasing thrombin generation. In a report of 107 patients with premature MI but no significant coronary artery stenosis (average age 44), the prevalence of carriers for factor V Leiden was significantly higher in these patients compared to 244 with an MI and significant stenoses and 400 healthy controls (12 versus 4.5 and 5 percent) [28]. At least in young women, the increase in risk with factor V Leiden may be confined to smokers

[29]. (See "Factor V Leiden and activated protein C resistance: Clinical manifestations and diagnosis".)

- Psychosocial factors, such as anger, may be important in the development of premature CHD [30]. (See "Psychosocial factors in coronary and cerebral vascular disease".)
- In women, acute MI may be

g) *Angiographic Findings*

In the majority of patients younger than 45 years of age, angiographic studies were performed because of a history of MI. As expected, major differences were found when compared to older patients.

h) *Coronary disease severity*

Younger patients have a higher incidence of normal coronary arteries, mild luminal irregularities, and single vessel coronary artery disease than do older patients [10,12,13,15,38].

One of the largest reports of angiographic findings in young patients with CHD comes from a substudy of the CASS trial, which compared the results of coronary angiography in 504 young men (≤ 35 years of age) and women (≤ 45 years of age) with a history of an MI to those in over 8300 older patients [12]. The following significant differences were noted:

- Normal coronary arteries were more common in the young patients (18 versus 3 percent). Young women had a higher frequency of angiographically normal coronary arteries than young men, despite a 10 year age difference in the definition of "young."
- Single vessel coronary disease was more common (38 versus 24 percent) and three vessel disease was less common (14 versus 39 percent) in the younger patients.
- Although some series have shown a predilection for involvement of the left anterior descending artery in young patients [13,38], this was not found in the CASS substudy.

In another large series of 823 young patients with CHD, single vessel disease was present in 55 to 60 percent [6].

- Symptomatic coronary heart disease (CHD) is uncommon in young men and women (age less than 40 to 45 years). (See 'Introduction' above.)
- Younger patients with CHD more often have a family history of premature CHD. (See 'Family history' above.)
- Cigarette smoking is the most common and most modifiable risk factor in young patients. (See 'Smoking' above.)
- Diabetes mellitus and hypertension appear to be less common in young patients with CHD than in older patients. (See 'Diabetes and hypertension' above.)

- Other risk factors such as cocaine use, factor V Leiden, and oral contraceptive use are more common in younger individuals with CHD. (See 'Other factors' above.)
- The clinical presentation of CHD in younger patients is different from that in older patients. A higher proportion of young patients do not experience angina, and, in the majority of cases, an acute coronary syndrome that progresses rapidly to MI if left untreated is the first manifestation of CHD. (See 'Clinical presentation' above.)
- Younger patients have a higher incidence of normal coronary arteries, mild luminal irregularities, and single vessel coronary artery disease than do older patients. Rarer causes of CHD such as spontaneous coronary dissection or Kawasaki disease occur more commonly in the young. (See 'Angiographic findings' above.)
- In general, the management of CHD in the young is similar to that in older individuals.