



GLOBAL JOURNAL OF MEDICAL RESEARCH: C
MICROBIOLOGY AND PATHOLOGY
Volume 14 Issue 1 Version 1.0 Year 2014
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Diagnostic Role of the Bethesda System for Reporting Thyroid Lesions: Effective Tool for Managing Thyroid Lesions

By Dr. Vasudha M. Bhagat, Dr. Hemali J. Tailor, Dr. KumarBhargav R. Kaptan,
Dr. Varsha Baladawa, Dr. Gunjan H. Prasad, & Dr. Peeyush K. Saini

Government Medical College, India

Abstract- Introduction: As Fine needle aspiration cytology (FNAC) is the primary investigation for the management of thyroid lesions, its interpretation is very crucial. The Bethesda System for Reporting Thyroid Cytopathology (TBSRTC) for clarity of communication recommends that each case should be reported in 1 of 6 general diagnostic categories facilitating communication among the managing team of doctors and leaves almost no confusion regarding the management of thyroid lesions.

Aim: To study utility of The Bethesda system of reporting thyroid cytopathology.

Materials and Method: During period of 1 year from 1st January to 31st December 2012 aspiration cytology has been carried out in 160 thyroid swellings referred to cytology department of a tertiary care hospital in Surat. Fine needle aspiration cytology was performed using mainly non-aspiration and aspiration techniques. All the cases were reported using TBSRTC. Cases were followed whenever possible.

Keywords: FNAC, non aspiration technique, thyroid lesions, the bethesda.

GJMR-C Classification : NLMC Code: WK 200



DIAGNOSTIC ROLE OF THE BETHESDA SYSTEM FOR REPORTING THYROID LESIONS EFFECTIVE TOOL FOR MANAGING THYROID LESIONS

Strictly as per the compliance and regulations of:



RESEARCH | DIVERSITY | ETHICS

© 2014. Dr. Vasudha M. Bhagat, Dr. Hemali J. Tailor, Dr. KumarBhargav R. Kaptan, Dr. Varsha Baladawa, Dr. Gunjan H. Prasad, & Dr. Peeyush K. Saini. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License (<http://creativecommons.org/licenses/by-nc/3.0/>), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Diagnostic Role of the Bethesda System for Reporting Thyroid Lesions: Effective Tool for Managing Thyroid Lesions

Dr. Vasudha M. Bhagat ^α, Dr. Hemali J. Tailor ^σ, Dr. KumarBhargav R. Kaptan ^ρ, Dr. Varsha Baladawa ^ω
Dr. Gunjan H. Prasad [¥], & Dr. Peeyush K. Saini [§]

Abstract- Introduction: As Fine needle aspiration cytology (FNAC) is the primary investigation for the management of thyroid lesions, its interpretation is very crucial. The Bethesda System for Reporting Thyroid Cytopathology (TBSRTC) for clarity of communication recommends that each case should be reported in 1 of 6 general diagnostic categories facilitating communication among the managing team of doctors and leaves almost no confusion regarding the management of thyroid lesions.

Aim: To study utility of The Bethesda system of reporting thyroid cytopathology.

Materials and Method: During period of 1 year from 1st January to 31st December 2012 aspiration cytology has been carried out in 160 thyroid swellings referred to cytology department of a tertiary care hospital in Surat. Fine needle aspiration cytology was performed using mainly non-aspiration and aspiration techniques. All the cases were reported using TBSRTC. Cases were followed whenever possible.

Observations and Results: The cytological samples were assessed by qualified consultant pathologists and were categorized in category 1 to 6, six tier system according to TBSRTC criteria given by National Cancer Institute USA. Fine needle aspiration cytology analysis revealed 149 (93.12%) non-neoplastic and 11 (6.88%) neoplastic lesions. Major bulks of 140 cases (87.5%) were of category II.

Conclusion: Application of TBSRTC bridges the gap in communication amongst the cytopathologist, endocrinologists, surgeons, radiologists, and other health care providers not only in the confined region but also worldwide and leaves no confusion regarding management of thyroid lesions. Few of the borderline lesions often create the confusion which are eliminated by TBSRTC.

Keywords: FNAC, non aspiration technique, thyroid lesions, the bethesda.

I. INTRODUCTION

Thyroid enlargement is a common occurrence in most regions of the world including India. Being tertiary care hospital we frequently encounter such

cases because southern Gujarat and surrounding mountainous areas are one of the endemic goitre belt in India.

Thyroid lesions are one of the most common lesions subjected to the cytopathology as Fine needle aspiration cytology (FNAC) is the first line investigation apart from other investigations like ultrasonography (USG), thyroid function tests, thyroid scan, and antibody levels are done subsequently to select the patients who require surgery and those that can be managed conservatively. ^[1, 2]

For the primary evaluation of patients FNAC has proven to be a rapid, cost-effective, safe and reliable method of investigation like in lesions of breast, lymph nodes and others. ^[3, 4] However, the success of FNAC is dependent on several important contributing influences including aspirator experience, skilful interpretation rational analysis and its application in management. Data from the Surveillance Epidemiology and End Results (SEER) registry show an increasing prevalence of differentiated thyroid cancer worldwide. ^[5, 6]

The increasing prevalence of thyroid cancer and improvements in the technology and resolution of ultrasound machines have led to an increasing number of cytological diagnostic procedures. ^[7] So being the primary investigation, interpretation and application of FNAC findings is very crucial for further management of thyroid lesions especially for ruling out need of surgery. Uniform communication amongst the cytopathologist, surgeons, endocrinologists, radiologists, and other health care providers will eliminate confusion regarding management. Few borderline thyroid lesions often create confusions regarding treatment. To eliminate such dilemma National Cancer Institute USA in 2007 conference meet was organised in Bethesda with one of the objectives being to standardize the diagnostic terminology for the reporting of thyroid cytopathology results. The recommendations resulting from this conference led to the formation of The Bethesda System for Reporting Thyroid Cytopathology (TBSRTC). This classification scheme has achieved its purpose of standardization of thyroid-reporting cytopathology, as evidenced by several publications. ^[8]

It has been shown that the best way to implement TBSRTC in a pathology laboratory and to

Author ^α: Additional Professor Department of Pathology, Government Medical College, Surat, 395001, Gujarat, India.

Author ^σ: Assistant professor Department of Pathology, Government Medical College, Surat, 395001, Gujarat, India.

Author ^ρ: Senior resident Department of Pathology, Government Medical College, Surat, 395001, Gujarat, India.

Author [§]: Government Medical College, Surat.
e-mail: piyushsaini622@gmail.com

convince the clinicians of its validity and usefulness is to compare the outcome data prior to and after the implementation of TBSRTC. Current study was mainly focussed to study the role of reporting system.

II. MATERIAL AND METHOD

Study includes 160 cases of thyroid swellings patients referred to the cytopathology section of pathology department in a tertiary care hospital in southern Gujarat between January 2012 and December 2012. Patient's details regarding history, clinical examination, thyroid function tests, clinical diagnosis, FNAC and histological data whenever possible were noted. The data were analysed in simple statistical tables.

All the cases of thyroid swelling subjected to FNAC were performed by cytopathologist. Prior to procedure, palpation was carried out to note the mobility of the thyroid during swallowing and the presence of any enlarged cervical lymph node. The patients were made to lie supine with their necks stretched up. A 23-24 gauge needle was used, with non-aspiration technique in most cases and very few cases with aspiration technique by a 10 ml disposable syringe. Two or more passes at different sites were made in each case. No major complications like penetration injury to the trachea, laryngeal nerve, or hematoma were recorded. Slides were prepared from aspirated material. In the case of cystic nodules, the cysts' contents were aspirated, centrifuged, and slides made from the sediment for cytological examination. The slides were stained with May-Grunwald Giemsa (MGG), Papanicolaou [PAP] stain and Haematoxylin and Eosin (H&E) and examined under light microscope. The microscopic diagnosis was interpreted under guidelines laid down by TBSRTC including categories I to VI (table 1) after taking into account of all available clinical, radiology and other data. Whenever possible, further sub typing was given. The cytological diagnosis were correlated with clinical features, thyroid function tests, subjected to histopathological examination whenever possible.

III. RESULTS

Study includes 160 cases with age range between 5 to 70 years. Maximum cases were in 21 – 50 years of age group. Bulk of the cases were females comprising of 136 cases (85%) and 24 cases (15%) were males and female: male ratio was 5.67:1. Long standing history of thyroid swelling was the main presenting symptom. Swelling was mainly diffuse and nodular in few cases. Symptoms like pain in the neck region, dysphagia, hoarseness of voice and cough were rare.

FNAC of 160 patients yielded the following diagnosis as depicted in Table 1.

a) Non neoplastic

Category I: The totals of 9 patients (5.63%) were diagnosed under nondiagnostic or unsatisfactory category.

Category II: It included most of the study cases with 140 cases (87.5%) of total. Age ranges with maximum number of cases were in 20-50 year age group. It consists of 'non-neoplastic' or 'negative for malignancy' cases like colloid goitre with 97 cases (69.29%), Thyroiditis with 36 cases (25.71%) and Adenomatoid goitre 7 cases (5%). All of these benign cases were just followed up and surgery was prevented.

Category III: It includes lesions which were not clearly benign or malignant. Conclusive opinion was not possible. We did not have any case diagnosed in this category.

Category IV: It includes 5 cases (3.12%) of follicular neoplasm (FN) and suspicious of follicular neoplasm (SFN).

Category V: It includes only 1 case (0.63%).

Category VI: It included 5 cases (3.12%) comprising of 2 cases each of papillary thyroid carcinomas and anaplastic carcinoma of thyroid with 1 case of medullary carcinoma of thyroid.

IV. DISCUSSION

As in management of thyroid lesions, FNAC is the gold standard and primary investigation of choice along with other investigations like (USG) ultrasonography examination, thyroid function tests, thyroid scan, and antibody levels are done subsequently to find out patients who require surgery and those that can be managed conservatively.^[1, 2]

Being a tertiary care hospital we have many patients of thyroid disorders from the South Gujarat region including Bharuch, Vapi, Songadh, Vyara and other goitre belts. Majority of these lesions are usually benign and require no aggressive treatment. So interpretation in each case is very crucial for further management. Also we want to establish uniform communication between the pathologist, radiologist, endocrinologist, surgeons and treating physicians. So that there would be no confusion regarding further management.

TBSRTC is a vital guideline which can bridge the communication gap and useful to maintain uniformity not only in the confined region but also worldwide. We followed the TBSRTC guidelines and every case was classified according to six tier reporting guideline from category I to VI.

Study includes total 160 cases of thyroid lesions which comprised of 140 cases (87.5%) of total. Published data suggest FNA has an overall accuracy rate around 75% in the detection of thyroid malignancy.^[9]

The age group which was studied ranged from 5 years to 70 years and maximum no. of cases were in the age group 20-50 years means bulk of thyroid diseases were frequently encountered in young and middle aged group also the majority of cases were the females in reproductive age groups. In present study a female preponderance was noted. Similar female preponderance was noted by Unnikrishnan et al.^[10]

Neoplastic lesions were 11 cases (6.88%). The benign cyst consistent with thyroglossal cyst were 4 cases (2.06%) and others were 4 cases (2.06%). The bulk of the goitre cases were in the age group of 20-50 years and thyroiditis cases were in 11-40 years and mainly in the reproductive age group of the females.

Category I: This category includes cases in which sufficient material was not available like insufficient follicular cells (Satisfactory for evaluation: six groups of well visualised follicular cells with at least ten cells per group), cyst fluid only, obscuring blood, only macrophages, preparation artefact. In such cases repeat FNA was carried out under ultrasound guidance.

Category II: Majority of lesions were benign mainly of colloid goitre 97 cases (69.29%) out of 140. In comparison to various studies benign category includes 60-70%^[8] reason for that is we have goitre belt here. Similar findings were observed Unnikrishnan et al.^[10] The chances of thyroiditis after reproductive age appeared minimal from this study.

Category III: It is reserved for specimens that contain cells (follicular, lymphoid, or other) with architectural and/or nuclear atypia that is not sufficient to be classified as suspicious for a follicular neoplasm, suspicious for malignancy, or malignant. Management in such lesions is repeat FNAC after an appropriate interval. To be noted that this category is of last resort & should not be used indiscriminately.^[8]

Category IV: The goal of this category is to identify all potential follicular carcinomas and refer them for a diagnostic lobectomy. Although these cytomorphologic features do not permit distinction from a follicular adenoma (FA), they are reportable as Follicular Neoplasm (FN) or suspicious of Follicular Neoplasm (SFN), leading to a definitive diagnostic procedure, usually lobectomy.^[11, 12, 13] The term SFN is preferred by some laboratories over FN for this category because a significant proportion of cases (up to 35%) prove not to be neoplasms but rather hyperplastic proliferations of follicular cells, most commonly those of multinodular goiter.^[14-17] About 15% to 30% of cases called FN/SFN prove to be malignant.^[11, 14, 16] The majority of FN/SFN cases turn out to be FAs or adenomatoid nodules of multinodular goiter, both of which are more common and outnumbers the Follicular carcinoma.

Category V: FNAC can diagnose many of the thyroid cancers with fair accuracy, especially papillary thyroid

carcinoma (PTC) which can be diagnosed with certainty by FNA. But the nuclear and architectural changes of some PTCs are subtle and focal. This is particularly true of the follicular variant of PTC, which can be difficult to distinguish from a benign follicular nodule.^[18] Other PTCs may be incompletely sampled and yield only a small number of abnormal cells.^[19] If only 1 or 2 characteristic features of PTC are present or if they are only focal and not widespread throughout the follicular cell population, or if the sample is sparsely cellular, a malignant diagnosis cannot be made with certainty. Such cases occur with some regularity, and they are best classified as suspicious for malignancy," qualified as "suspicious for papillary carcinoma." Such cases suspicious for papillary carcinoma are resected by lobectomy or thyroidectomy. Most (60%-75%) prove to be papillary carcinomas, and the rest are usually FAs.^[11, 14, 16] The same general principle applies to other thyroid malignancies like medullary carcinoma and lymphoma, but these are less frequent than PTC. Such cases were considered after correlating the other findings like serum calcitonin and calcium levels and other relevant data in medullary carcinoma.

Category VI: The malignant category is used whenever the cytomorphologic features are indicative of malignancy. After confirming the malignancy the sub classification was done after summarizing all the results. Approximately 3% to 7% of thyroid FNAs have conclusive features of malignancy, and most are papillary carcinomas.^[11, 14] Malignant lesions are usually treated by thyroidectomy, with some exceptions (e.g., metastatic tumors, non-Hodgkin lymphomas, and undifferentiated carcinomas). According to studies the positive predictive value of a malignant FNA interpretation is 97% to 99%.

We here practiced mainly the non-aspiration technique in almost all cases for studying cytology and found that it is better than aspiration technique. Aspiration technique is associated with low cellularity and more blood as compared to non-aspiration method. We recommend the non-aspiration method for FNAC of thyroid lesions. Similar suggestions by different studies like Maurya et al.^[20] also recommend the non-aspiration technique better for thyroid lesion evaluation by FNAC. The study found that it is difficult to differentiate follicular/Hurthle cell adenoma from carcinoma on cytological assessment because cytology cannot evaluate the criteria of vascular or capsular invasion or of intrathyroid spread. But the papillary carcinoma, Anaplastic carcinoma and medullary carcinoma can be diagnosed by characteristic cytological features.

V. CONCLUSION

TBSRTC is a vital guide for accurate management of thyroid lesions. Classifying the lesions in six categories and following the guidelines given by

The Bethesda USA meetings solves all problems regarding the management of thyroid lesions and leaves no confusion. It plays a big role in establishing the uniform communications between the managing medical personnel. Marked cellularity of the smear is the problem inherent in thyroid FNAC. Increased cellularity of the smear and loss of cohesion may be present in hyperplastic/adenomatous goiter and follicular neoplasm which causes difficulty in differentiating them. This can be solved by using The Bethesda System of Reporting thyroid lesions. We experienced that The Bethesda Reporting System is best for management of thyroid lesions as it gives uniform reporting system.

REFERENCES RÉFÉRENCES REFERENCIAS

- Caruso D, Mazzaferri EL. Fine needle aspiration biopsy in the management of thyroid nodules. *Endocrinologist* 1. 1991: 194-202.
- Giuffrida D, Gharib H. Controversies in the management of cold, hot and occult thyroid nodules. *Am J Med* 1995; 99:642-50.
- Silverman JF, West RE, Larkin EW, Park HM, Finley JL, Swanson MS et al. The role of FNAC in the rapid diagnosis and management of thyroid neoplasm. *Cancer*. 1986; 57: 1164-1170.
- Kini U, Buch A, Bantwal G. Role of FNA in the medical management of minimally enlarged thyroid. *DiagnCytopathol* 2006; 34:196-200.
- Ries LAG, Melbert D, Krapcho M, et al: SEER Cancer Statistics Review, 1975–2005. Bethesda, National Cancer Institute, 2007. [http:// seer.cancer.gov/csr/1975_2005/](http://seer.cancer.gov/csr/1975_2005/)
- Chen AY, Jemal A, Ward EM: Increasing incidence of differentiated thyroid cancer in the United States, 1988–2005. *Cancer* 2009; 115: 3801–3807.
- Ross DS: Predicting thyroid malignancy (editorial). *J ClinEndocrinolMetabol* 2006; 91:4253–4255.
- Ali SZ, Cibas ES: *The Bethesda System for Reporting Thyroid Cytopathology. Definitions, criteria and explanatory notes.* New York, Springer, 2010.
- Robinson IA, Cozens NJ. Does a joint ultrasound guided cytology clinic optimize the cytological evaluation of head and neck masses? *ClinRadiol* 1999; 54:312-6.
- Unnikrishnan et al: Endocrine Society of India management guidelines for patients with thyroid nodules: A position statement, *Indian J EndolMetab.* 2011 Jan-Mar; 15(1): 2–8.
- Gharib H, Goellner JR, Johnson DA. Fine-needle aspiration cytology of the thyroid: a 12-year experience with 11,000 biopsies. *Clin Lab Med.* 1993; 13:699-709.
- Layfield L, Cochand-Priollet B, LiVolsi V, et al. Post thyroid FNA testing and treatment options: a synopsis of the National Cancer Institute Thyroid Fine Needle Aspiration State of the Science Conference. *Diagn Cytopathol.*2008; 36:442-448.
- Mazzaferri EL. NCCN thyroid carcinoma practice guidelines. *Oncology.* 1999; 13:391-442.
- Yang J, Schnadig V, Logrono R, et al. Fine-needle aspiration of thyroid nodules: a study of 4703 patients with histologic and clinical correlations. *Cancer.* 2007; 111:306-315.
- Deveci MS, Deveci G, LiVolsi VA, et al. Fine-needle aspiration of follicular lesions of the thyroid: diagnosis and follow-up. *Cytojournal.* 2006; 3:9.
- Baloch ZW, Fleisher S, LiVolsi VA, et al. Diagnosis of “follicular neoplasm”: a gray zone in thyroid fine-needle aspiration cytology. *DiagnCytopathol.* 2002; 26:41-44.
- Kelman AS, Rathan A, Leibowitz J, et al. Thyroid cytology and the risk of malignancy in thyroid nodules: importance of nuclear atypia in indeterminate specimens. *Thyroid.* 2001; 11:271-277.
- Chung D, Ghossein RA, Lin O. Macrofollicular variant of papillary carcinoma: a potential thyroid FNA pitfall. *DiagnCytopathol.* 2007; 35:560-564.
- Renshaw AA. Focal features of papillary carcinoma of the thyroid in fine-needle aspiration material are strongly associated with papillary carcinoma at resection. *Am J Clin Pathol.* 2002; 118:208-210.
- Mauya et.al: Comparison of aspiration vs non-aspiration techniques in fine-needle cytology of thyroid lesions: *J Cytol.* April 2010, 27(2): 51–54.

Table 1: Distribution of thyroid lesions according to age and sex after applying TBSRTC

Age in years	Male	Female	I	II	III	IV	V	VI	Total
1-10	1	1	0	2	0	0	0	0	2
11-20	2	5	0	6	0	0	0	1	7
21-30	4	43	2	44	0	1	0	0	47
31-40	11	41	4	45	0	0	1	2	52
41-50	4	27	1	25	0	3	0	2	31
51-60	1	8	0	8	0	1	0	0	9
61-70	1	11	2	10	0	0	0	0	12
Total	24 (15%)	136 (85%)	9 (5.63%)	140 (87.5%)	0 (0%)	5 (3.12%)	1 (0.63%)	5 (3.12%)	160

I-Non diagnostic or unsatisfactory, II- Benign, III- Atypia of undetermined significance or Follicular lesions of undetermined significance, IV- Follicular neoplasm or

suspicious of follicular neoplasm, V- Suspicious of malignancy, VI- Malignant lesion.

Table 2 : Diagnostic categories, implied risk of malignancy & recommended clinical management according TBSRTC. (Modified by Ali and Cibas)^[8]

Diagnostic category	Cytological diagnosis	Risk of malignancy	Clinical management
I	Non-diagnostic or Unsatisfactory	1-4 %	Repeat FNA with ultrasound guidance
II	Benign	0-3 %	Clinical follow-up
III	AUS/FLUS*	5-15 %	Repeat FNA
IV	FNS/SFN †	15-30 %	Surgical lobectomy
V	Suspicious of malignancy	60-75 %	Near total thyroidectomy or Surgical lobectomy
VI	Malignant	97-99 %	Near total thyroidectomy

*AUS/FLUS – Atypia of undetermined significance / Follicular lesion of undetermined significance.

† FNS / SFN – Follicular neoplasm / Suspicious of follicular neoplasm.




Figure 1 : Abundant watery thin colloid in Colloid Goitre




Figure 2 : Hurthle cell changes in Hashimoto thyroiditis




Figure 3 : Numerous macrophages only: Nondiagnostic category




Figure 4 : Medullary carcinoma of thyroid with amyloid like material




Figure 5 : Follicular neoplasm with numerous microfollicles




Figure 6 : Anaplastic carcinoma of thyroid

