



GLOBAL JOURNAL OF MEDICAL RESEARCH: C
MICROBIOLOGY AND PATHOLOGY
Volume 14 Issue 6 Version 1.0 Year 2014
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Patterns of Thyroid Lesions: A Histomorphological Study

By VL Ramesh, S Sunitha, R Rupnarayan & C Narayana Murthy

Basaveshwara Medical College, India

Abstract- Background: Thyroid lesions are fairly common in and around Kolar town. This study was undertaken to study the various histomorphological types of neoplastic and non-neoplastic lesions of the thyroid and to correlate these with respect to age and sex.

Methods: All thyroid specimens received at the pathology Department of Sri Devaraj Urs Medical College, Kolar during the period January 2000 to December 2004 were processed. A detailed histomorphological study was done. The histomorphological type was correlated with the age, sex and clinical presentation.

Results: Total 120 cases of thyroid were studied. Most common age group affected was between 3rd and 5th decade. Females were predominantly affected. The non-neoplastic lesions reported in this study were thyroglossal duct cyst 1 case (0.83%), De Quervain thyroiditis 1 case (0.83%), Hashimoto thyroiditis 11 cases (9.16%), colloid goiter 7 cases (5.83%), multinodular goiter 35 cases (29.16%), diffuse toxic goiter 2 cases (1.66%). Among neoplastic lesions follicular adenoma 43 cases (35.83%), atypical follicular adenoma one case (0.83%), papillary carcinoma classic variant 11 cases (9.16%), follicular variant of papillary carcinoma 7 cases (5.83%) and one case (0.83%) of medullary carcinoma.

Keywords: *goiter, thyroid lesions.*

GJMR-C Classification : *NLMC Code: WK 200*



Strictly as per the compliance and regulations of:



Patterns of Thyroid Lesions: A Histomorphological Study

VL Ramesh^α, S Sunitha^σ, R Rupnarayan^ρ & C Narayana Murthy^ω

Abstract- Background: Thyroid lesions are fairly common in and around Kolar town. This study was undertaken to study the various histomorphological types of neoplastic and non-neoplastic lesions of the thyroid and to correlate these with respect to age and sex.

Methods: All thyroid specimens received at the pathology Department of Sri Devaraj Urs Medical College, Kolar during the period January 2000 to December 2004 were processed. A detailed histomorphological study was done. The histomorphological type was correlated with the age, sex and clinical presentation.

Results: Total 120 cases of thyroid were studied. Most common age group affected was between 3rd and 5th decade. Females were predominantly affected. The non-neoplastic lesions reported in this study were thyroglossal duct cyst 1 case (0.83%), De Quervain thyroiditis 1 case (0.83%), Hashimoto thyroiditis 11 cases (9.16%), colloid goiter 7 cases (5.83%), multinodular goiter 35 cases (29.16%), diffuse toxic goiter 2 cases (1.66%). Among neoplastic lesions follicular adenoma 43 cases (35.83%), atypical follicular adenoma one case (0.83%), papillary carcinoma classic variant 11 cases (9.16%), follicular variant of papillary carcinoma 7 cases (5.83%) and one case (0.83%) of medullary carcinoma.

Conclusion: Total 120 thyroid lesions were studied in the present study out of this 57 cases were non-neoplastic and neoplastic were 63 cases. Sub-acute thyroiditis reported was 0.83%. Studies conducted by others range from 0.15% to 4.25%. Hashimoto thyroiditis reported was 9.16%. Other studies range from 4.25% to 5.68%. Colloid goiter reported was 5.83%. Reports of others range from 49.18% to 36%. Multinodular goiter reported was 29.16%. Other studies reported range from 3.19% to 18%. Diffuse toxic goiter reported was 1.66%. Other study reported incidence of 2.12%. Total neoplastic lesions reported was 52.5%. Benign lesions reported were 36.66%. Of these follicular adenoma constituted 35.83%. A typical adenoma reported was 0.83%. Total malignant lesions reported were 15.63%. Total thyroid malignancies reported by other studies range from 14% to 31.91%. Papillary carcinoma classic variant found was 9.16%, follicular variant of papillary carcinoma reported was 5.83%. Papillary carcinoma reported by other studies range from 7.44% to 61.1%. Medullary carcinoma constituted 5.16%. Other study reported as 6.5% of medullary carcinomas. In conclusion, most common symptoms was neck swelling. Majority patients were between 3rd and 6th decade with female preponderance. Follicular adenoma was the most common pathological lesion. Commonest malignancy was the papillary carcinoma.

Keywords: goiter, thyroid lesions.

Author α σ ρ: Department of Pathology, Basaveshwara Medical College and Hospital, Chitradurga, Karnataka, India.
e-mail: roknet.trade@gmail.com

I. INTRODUCTION

Thyroid gland is unique among the endocrine glands in having a wide spectrum of diseases ranging from functional enlargements immunologically mediated enlargements to the neoplastic lesions. These enlargements may be diffuse or nodular at times causing obvious physiological changes. In contrast patient having a papillary carcinoma thyroid with lymph node secondaries may remain asymptomatic till a very late stage. Occasionally a patient may present with obvious metastatic disease with an undetectable primary (occult or hidden malignancy of thyroid).

Thyroid gland lesions appear to be common in and around the city of Kolar. So the classification of various histomorphological types of tumor is important to categorize the lesion into non-neoplastic and neoplastic lesion of thyroid. The WHO published its second edition on the histological classification of thyroid tumors in 1988¹. Based on WHO we can classify our neoplastic lesions. It will be of great value for clinicians for further therapy and prognosis.

The present study is intended to study the various histomorphological changes of non – neoplastic and neoplastic lesions of the thyroid, as there are no studies on the patterns of thyroid lesions in and around Kolar, which has high number of patients with thyroid enlargements.

II. MATERIALS AND METHODS

The material for the present study comprised of specimens received at Department of Pathology, Sri Devaraj Urs Medical College, Tamaka, Kolar, between January 2000 and December 2004 from patients admitted to R.L. Jalappa Hospital and S.N.R. Hospital, Kolar. All cases registered in our department files for thyroidectomy and diagnosed between January 2000 and December 2004 for a period of five years were reviewed. The period of retrospective study was from Jan 2000 to Dec 2003 and prospective study from Jan 2003 to Dec 2004.

The specimen were fixed in 10% formalin for 24-48 hour. Large specimens were cut serially (at 1cm thickness) before fixing. After fixation, representative areas were selected for paraffin embedding. In case of encapsulated lesions, adequate representation from tumour capsule – thyroid interface was given. Section were cut at 4-5 microns thick and stained with

hematoxylin and eosin and studied. This was done for all cases received between January 2003 and December 2004.

Special stains like methyl violet, vanGieson, masson trichrome and congo red were performed for necessary cases.

Stained histopathology slides were studied in detail. All details of the case consisting of clinical history, external examination, gross features, microscopic features and final diagnosis were filled in a proforma. Details from all proforma were tabulated in a master chart.

Retrospective study for three years from January 2000 to December 2002 (48 cases). Prospective study for two years from January 2003 to December 2004 (75cases).

A total number of 8,638 specimens were received during this period. Of these 123 cases were clinically thyroid neck swellings. Among these 3 cases were excluded, which were histopathologically diagnosed as granulation tissue (Sl.No.28 and 42) and normal lymph node (Sl.No.118). Remaining 120 cases were thyroid lesions and included in this study.

III. RESULTS

The present study is undertaken for a period of five years between January 2000 and December 2004.

Table 1 : Age and Sex Distribution

Sl.No	Age	No.of.Cases	Male	Female
1	<10	1	-	1
2	10-19	4	-	4
3	20-29	33	4	29
4	30-39	40	2	38
5	40-49	21	5	16
6	50-59	11	1	10
7	60-69	9	-	9
8	70-79	1	-	1
	Total	120	12(10%)	108(90%)

Table 2 : Symptoms with which the patient presented

Sl.No	Symptoms	No.of.Cases
1	Neck Swelling	120 (100%)
2	Dysphagia	24 (20%)
3	Dyspnoea	15 (12.5%)

Table 3 : External examination of neck swelling had following features

Sl.No	Signs	No.of.Cases
1	Diffuse. a. Sub acute thyroiditis b. Hashimoto thyroiditis c. Colloid goiter d. Diffuse toxic goiter e. Papillary carcinoma	34 (28.3%)
2	Solitary nodule. a.Follicular adenoma b. Atypical adenoma c. Thyroglossal duct cyst d. Papillary carcinoma e. Medullary carcinoma	51 (42.5%)
3	Multiple nodules. a. Multinodular goiter	35 (29.2%)

Table 4 : Morphologic types of thyroid lesions

Sl.No	Morphologic type	No.of.Cases	%
1	Non - Neoplastic lesions	57	47.5
2	Neoplastic lesions	63	52.5

Table 5 : Histomorphologic types and their incidence in different sex and age groups

SL.No	Age in years	<10		10-19		20-29		30-39		40-49		50-59		60-69		70-79		TOTAL
	Types	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	
1	Thyroglossal duct cyst	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1
2	Sub- acute thyroiditis	-	-	-	-	-	-	-	1	-	-	-	-	-	-	-	-	1
3	Hashimoto thyroiditis	-	-	-	-	-	1	-	3	1	2	-	3	-	1	-	-	11
4	Colloid goiter	-	-	-	-	1	2	-	2	-	1	-	-	-	1	-	-	7
5	Multinodular goiter	-	-	-	2	2	6	-	11	1	8	-	3	-	2	-	-	35
6	Diffuse toxic goiter	-	-	-	-	-	1	-	1	-	-	-	-	-	-	-	-	2
7	Follicular adenoma	-	-	-	2	3	12	-	11	2	6	-	3	-	3	-	1	43
8	Atypical adenoma	-	-	-	-	-	1	-	-	-	-	-	-	-	-	-	-	1
9	Papillary Ca.Classic	-	-	-	-	-	2	1	3	1	-	1	2	-	1	-	-	11
10	Papillary.Ca.Follicular	-	-	-	1	-	1	1	4	-	-	-	-	-	-	-	-	7
11	Medullary Carcinoma	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	1
	TOTAL	-	1	-	5	6	26	2	36	5	17	1	11	-	9	-	1	120

Table 6 : Histomorphologic types of non-neoplastic lesions of thyroid

Sl.No	Types	No.of.Cases	%
1	Thyroglossal duct cyst	1	0.83
2	Sub- acute thyroiditis	1	0.83
3	Hashimoto thyroiditis	11	9.16
4	Colloid goiter	7	5.83
5	Multinodular goiter	35	29.16
6	Diffuse toxic goiter	2	1.66
	Total	57	47.5

Table 7 : Histomorphologic types of neoplastic lesions

Sl. No	Types	No of cases	%
1	Follicular adenoma	43	36
2	Atypical adenoma	1	1
3	Papillary .Ca. Classic	11	9
4	Papillary .Ca. Follicular	7	6
5	Medullary Carcinoma.	1	1
	Total	63	(52.5%)

IV. DISCUSSION

Total 120 thyroid lesions were studied in the present study. Of this 57 cases were non-neoplastic and 63 cases were neoplastic consisting of 47.5% and 52.5% respectively. A study conducted by Sankaran⁹ reviewed 127 cases and found the percentage of non-neoplastic lesions as 85.8% and neoplastic as 14.2%. Non-neoplastic lesions, in this study there was one case of thyroglossal cyst (0.83%) out of 120 cases. This was a 5 years old female child. One case of sub-acute thyroiditis was reported (0.83%) in a 38 years female patient out of 120 cases. A study conducted by Arora and Gupta^{6,10} reviewed 94 cases and found the percentage of sub-acute thyroiditis was 4.25% (4 cases). Another study conducted by Meachim and Young⁸ reviewed 1285 cases and found the percentage of sub-acute thyroiditis was 0.15% (2 cases). Hashimoto thyroiditis accounted for 11 cases (9.16%) out of 120 cases. A study conducted by Arora and Gupta^{2,6,10} found Hashimoto thyroiditis were 4.25% (4 cases) out of 94 cases studied. Another study conducted by Meachim and Young⁸ reviewed 1285 cases and found the percentage of Hashimoto thyroiditis was 5.68% (73 cases). Total all types of the thyroiditis reported were 12 cases (10%) out of 120 cases. Total all types of thyroiditis reported in the study conducted by Arora and Gupta^{6,10} was 9.57% (9 cases) out of 94 cases. In another study conducted by Meachim and Young^{6,8} total all types of thyroiditis was 5.99% (77 case) out of 1285 cases studied. Colloid goiter formed 5.83% (7 cases). Maximum cases were in the 3rd to 5th decade of life and one male case was reported. There was a wide range in the incidence of the colloid goiter reported by several authors. In a study conducted by Sankaran^{6,9} the incidence of colloid goiter was 36%. The average age being 33 years with female preponderance. In another study conducted by Arora and Gupta¹⁰ the incidence of colloid goiter was 15.95%. In the study conducted by Meachim and Young⁸ the incidence of colloid goiter was 49.18%. Multinodular goiter was the most common non-neoplastic lesion in this study. There were 35 cases (29.16%) with peak age incidence seen between 3rd and 5th decade of life and was more common in females. In a study conducted by Sankaran^{6,9} the incidence of

multinodular goiter was 18% and average age incidence was 35 years. In the study conducted by Arora and Gupta¹⁰ the incidence of multinodular goiter was 3.19%. Diffuse toxic goiter accounted to 1.66% (2 cases). Both were female patients. The study by Arora and Gupta^{5,10} reported an incidence of 2.12%. Compared to the overall incidence of goiter (all types) in this study (36.65%). Kalpatrick et al^{6,11} reported the overall incidence as 39.4 %, predominantly in the 20-49 years age group.

Neoplastic lesions, benign and malignant tumors together formed 63 cases (52.5%) out of total 120 cases studied.

Benign lesions found were in 36.66% (44 cases). Of this follicular adenoma was reported in 35.83%(43 cases). Follicular adenoma was the most common lesion in this study and it was the most common neoplastic lesion. Maximum incidence was seen between 3rd and 5th decade of life with female preponderance. Five male patients were reported. In a study conducted by Arora and Gupta^{1,3,7,10} represents 36.17% of follicular adenoma out of 94 cases studied. In another study conducted by Thomas¹² follicular adenoma represented 21.3% out of 121 cases studied.

The comparison between different histological sub-types of follicular adenoma in this study with incidence reported by various authors is shown below:

Types	Present study (43 cases)	Arora and Gupta ^{7,10} (94 cases)	Thomas ^{6,12} (34 cases)
Micro follicular (foetal)	- 03	05	02
Macrofollicular (colloid)	- 25	27	56
Normofollicular (simple)	- 15	Nil	29
Hurthle cell adenoma	- Nil	02	02.

Atypical adenoma was found in one case (0.83%). This was female patient aged 27 years.

Malignant tumors (19 cases) constituted 15.63% of the total 120 cases studied.

In contrast, Sankaran⁹ reported an incidence of 14%. Arora and Gutpa^{7,10} reported an incidence of

31.91% and Thomas^{3,5,12} reported an incidence of 19%. **Papillary carcinoma classic variant** constituted 9.16%(11 cases). Most cases were aged 40 years and below. Two youngest patients were 22 years old females. The oldest patient was a 65 years female with lymph node metastasis. There were only three male patients.

Comparative analysis of histological types of the thyroid carcinoma

Sl. No	Types	Arora & Gupta ¹⁰ (94cases)	Thomas ¹² (23cases)	Woolner et al ¹³ (885 cases)	Burn & Taylor ¹⁴ (152 cases)	Present Study (19 case)
1	Papillary .Ca	23.33%	34.8%	61.1%	28.5%	94.73%
2	Follicular .Ca	63.33%	60.8%	17.7%	28.5%	-
3	Follicular + Papillary .Ca	-	4.4%	-	-	-
4	Medullary .Ca	-	-	6.5%	-	5.26
5	Anaplastic .Ca	13.33%	-	14.7%	43	-

V. CONCLUSION

Most of the patients presented with a symptoms of neck swelling. Majority of the patients were between 3^d and 6th decade. Females were predominantly affected.

The commonest lesion was follicular adenoma followed by multinodular goiter. Most common malignant lesion was papillary carcinoma.

The present study was undertaken to review the recent literature in recognising the histomorphologic criteria for the thyroid lesions and to correlate the histomorphological type of thyroid lesion with age and sex of patient in and around Kolar town. The drawback of this study was that the present data being hospital generated cannot be regarded as representative of the incidence of thyroid lesion in the general population.

REFERENCE RÉFÉRENCES REFERENCIAS

- Hedinger C, Williams ED, Sobin LH. The WHO classification of thyroid tumors: A commentary on the second edition. *Cancer* 1989; 63 : 908 – 911.
- Alrich EM, Blank RH, Allen MS. Carcinoma of the thyroid. *Ann Surg* 1955;153: 762 – 767.
- Carcangiu ML, DeLellis RA. Thyroid gland, In; Damjanov I, Lindoer J, editors. *Anderson Pathology Vol 2*, 10th edn. St. Louis: Mosby, 1996; p. 1943 – 1979.
- Keele CA, Neil E, Joels N. Thyroid, In; Samson Wright Applied Physiology, 13th edn. Delhi: Oxford University Press; 1985; p. 537 – 546.
- Virginia A, Livolsi R. Surgical Pathology of Thyroid, In; Major problems in Pathology Vol 22. Philadelphia: WB Saunders Co; 1990; p. 150 – 159.
- Cotran RS, Kumar V, Collins T. Thyroid, In : Robbins Pathologic Basis of Disease 6th edn. Philadelphia: WB Saunders Co, 1999; 1130 – 1147.
- Rosai J. Thyroid gland, In: Ackerman's Surgical pathology Vol 1 9th edn. St.Louis: Mosby, 2004; p. 515 – 594.
- Meachim G, Young MH. De Quervain's subacute granulomatous thyroiditis: Histologic identification and incidence. *J Clin Pathol* 1963; 16: 189 – 199.
- Sankaran V. Swelling of the thyroid. *J Ind Med Assoc* 1960; 34: 484 – 488.
- Arora HL, Gupta DP. Geographic pathology of thyroid diseases in Rajasthan. *J Ind Med Assoc* 1967; 48: 424-428.
- Kilpatrick R, Milne JS, Rushbrooke M, Wilson ESB, Wilson GM. A Survey of thyroid enlargement in two general practices in Great Britain. *Brit Med J* 1963; 29-34.
- Thomas PA. Thyroid adenoma. *J Ind Med Assoc* 1966; 46:189-193.
- Woolner LB, Behrs OH, Black BM, McKonahey WM, Keating FR. Classification and prognosis of thyroid carcinoma. *Am J Surg* 1961; 102:354-386.

14. Burn JI, Taylor SF. Natural history of thyroid carcinoma-A study of 152 treated patients. Brit Med J 1962; 1218-1223.

