



GLOBAL JOURNAL OF MEDICAL RESEARCH: F  
DISEASES

Volume 15 Issue 1 Version 1.0 Year 2015

Type: Double Blind Peer Reviewed International Research Journal

Publisher: Global Journals Inc. (USA)

Online ISSN: 2249-4618 & Print ISSN: 0975-5888

## Astigmatism among other Refractive Errors in Children of Southern Sri Lanka

By Prof. Saman Wimalasundera

*University of Ruhuna, Sri Lanka*

**Abstract-** The main goal of this study was to find the problem of astigmatism among children with refractive errors who were in the age group of 3 – 14 years. A two stage screening process was envisaged and 5649 children were screened to find visual abnormalities and other defects. Of them, 1233 (21.8%) were selected for second stage screening. The total ophthalmological manifestations found were 7.8% (78/1000). The analysis of different morbidities included refractive errors 6.2% and other eye diseases 1.6%. Among refractive errors 2.3% were unilateral and 3.9% were bilateral involvement. The prevalence of simple myopia was 1.65% and of hypermetropia was 0.95%. The astigmatism found was 1.5% unilaterally out of 2.3 of total unilateral refractive errors. Bilateral astigmatism was 2.1% out of 3.9% of total bilateral refractive errors. Findings indicate the existence of more astigmatic errors among Sri Lankan children either unilaterally or bilaterally.

**Keywords:** *refractive errors, astigmatism, simple myopia, hypermetropia, screening.*

**GJMR-F Classification :** *NLMC Code: WW 310*



*Strictly as per the compliance and regulations of:*



# Astigmatism among other Refractive Errors in Children of Southern Sri Lanka

Prof. Saman Wimalasundera

**Abstract-** The main goal of this study was to find the problem of astigmatism among children with refractive errors who are were in the age group of 3 – 14 years. A two stage screening process was envisaged and 5649 children were screened to find visual abnormalities and other defects. Of them, 1233 (21.8%) were selected for second stage screening. The total ophthalmological manifestations found were 7.8% (78/1000). The analysis of different morbidities included refractive errors 6.2% and other eye diseases 1.6%. Among refractive errors 2.3% were unilateral and 3.9% were bilateral involvement. The prevalence of simple myopia was 1.65% and of hypermetropia was 0.95%. The astigmatism found was 1.5% unilaterally out of 2.3 of total unilateral refractive errors. Bilateral astigmatism was 2.1% out of 3.9% of total bilateral refractive errors. Findings indicate the existence of more astigmatic errors among Sri Lankan children either unilaterally or bilaterally.

**Keywords:** refractive errors, astigmatism, simple myopia, hypermetropia, screening.

## I. INTRODUCTION

One important major disease condition that leads to visual failure and related symptoms such as headache, and asthenopia include refractive errors of light rays entering the eyes.

The refraction of light means, when the rays of light travels through air, and enter a denser transparent medium the speed of light is reduced and the light rays proceed at a different angle (bending of rays) the light is hence called refracted. The ray of light, if incident perpendicularly to the denser medium, it does not bend but only reduces the speed. The refraction power of the light is derived from the index of refraction of various parts of the eye through which light passes in to the eye.

When the light from an object reaches the eye it has to pass the cornea, aqueous humor, crystalline lens and vitreous humor to get at the retina where the stimulus of light ray is converted to neuro-electrical impulses. (Andcea 1987)

The curvature of the cornea and the lens play an important role in the refraction of the light. The cornea contributes approximately two thirds of the refractive power and the lens contributes about one third of the refractive power of the eye. The cornea has the index of refraction of 1.376 and lens has the average of 1.41. The total power of the eye does not equal to the sum of the power of the cornea and the lens because they are separated at aqueous and vitreous interfaces

and it is around + 58 diopters. The curvature of the human lens could be altered using the circular ciliary muscle and its refractive state could be actively changed according to the need.

When the refraction of the eye does not take place accurately to focus the rays on to the macula area of the retina, the condition is called the refractive error (AMETROPIA). In an eye with a refractive error the parallel light rays that come in to the eye will therefore focus in front or behind the plane of the retina when the eye's optical system is at rest. This process will result in the blurring of the image. In a schematic eye a 0.3mm shift in the focal place needs a correction of one diopter.

The optical system described above assumed to have spherical structures. That means all meridians of curved structures have equal spheroidal surfaces. Many optical systems in the eye are however are not so and resembles a toric surface. In a toric surface the curvature varies in different meridians. In such instances the light rays passing through steep meridians will be more deflected than rays which pass through flatter meridians. This is a complicated process and will result in defocusing the part of the object from which the light rays pass in to the eye through anomalous meridian. This condition is called Astigmatism.

Corneal toricity is the cause of most of the Astigmatism. Astigmatism could be simple myopic or hypermetropic depending on where the ametropic rays get their focal point. If the part of the rays is focused in front of the retina, it is called myopic astigmatism. When part of the rays is focused behind, it is called hypermetropic stigmatism. When only one meridian is ametropic (either myopic or hypermetropic) it is called simple myopic or hypermetropic astigmatism. In the mixed type of astigmatism one principal meridian is myopic type and the other one is hypermetropic type. When both meridians are ametropic, it is called compound astigmatism. If both meridians are hypermetropically deviated differently it is called compound hypermetropic astigmatism and if the same entity happens myopically the condition is called compound myopic astigmatism. In mixed type of astigmatism one meridian becomes hypermetropic whilst the other is myopic.

Astigmatism is again classified as regular or irregular. In the regular type the principal meridian are ninety degrees apart. (Perpendicular to each other). In irregular astigmatism the principal meridians are not

**Author:** DO, PhD Department of Community Medicine, Faculty of Medicine, Galle. e-mail: samanwimalasundera@gmail.com



perpendicular. Most astigmatism are regular type. (Gary Heitingod) (Jackson and Finlay 1985)

Refractive errors among children could be easily detected and treated. Delay in seeking treatment lead to significant ocular morbidity including amblyopia, low vision and poor quality of life (Resnikoff 2004) Having recognized the importance of the prevention of low vision, the 66th world health assembly initiated a global action plan for 2014 – 2019 to promote universal eye health. This plan requests member states to take measures to reduce avoidable visual impairment by 25% by the year 2019 (WHO 2013) (Deborah 1985)

Astigmatism can be corrected with special type of lenses called toric lenses. In a toric lens one meridian is more curved than all other meridians. The meridian of least curvature and greatest curvature are always at right angles in a toric lens. Toric lenses can be plus lenses (to correct hypermetropic astigmatism) or minus lenses (to correct myopic astigmatism) Toric lenses are called spherocylinders, in which the power of the lens act only perpendicular to its axis.

Clinically people with astigmatism experience headache, fatigue of eyes after work, eye strain and blurred vision at distances. These symptoms may not necessarily due to astigmatism only but exclusion of astigmatism is important in people with such symptoms.

## II. OBJECTIVE OF THE STUDY

1. To find the prevalence of astigmatism among school children and its different types.

### a) Study setting

One of the health unit areas of Galle (Bop-Poddala) district was selected for this study. All children of 3 – 14 years were taken as study subjects.

### b) The study design


The study design included three stages. In stage one the primary visual screening at schools, pre schools and in the community was done. The positive cases difficult cases, refusals and absentees were re screened at stage two screening in the field. The selected cases from second screening were referred to a third stage complete ophthalmological examination for diagnostic and treatment purposes at a well equipped eye clinic.

Primary and secondary screening of children was done by trained, medically qualified doctors with the help of the chief investigator in the field. The third stage examination was done by the ophthalmic qualified principal investigator. The health department field staff helped in other activities like finding subjects from the community, schools and pre schools.

## III. RESULTS

### a) Coverage of subjects

#### Screening of children




\* - School records


\*\* - PHM records

b) Stages of screening and referrals

Stage 1



Stage 2 screening



3<sup>rd</sup> stage examination




Table 1 : Break down of confirmed eye diseases after examination

Disease category	Visual defect		Total	Prevalence N=5649
	+	-		
Refractive errors	351	-	351	<b>6.2%</b>
Other eye diseases	39	49	88	<b>1.6%</b>
Total eye diseases	390	49	439	<b>7.8%</b>



**Table 2 :** Distribution of refractive errors – The involvement of the eye

Distribution	Eye		Prevalence
Unilateral	Right	40	0.7%
	Left	88	1.6%
Bilateral		223	3.9%
Total		351 eyes	6.2%

**Table 3 :** Distribution of refractive errors  
Different types – unilateral involvements

Types of refractive errors	involvements				Total		Prevalence N=5649
	Right		Left		No	%	
	No	%	No	%			
Simple myopia	8	20.5	23	25.8	31	24.2	0.5
Simple Hypermetropia	9	23.5	7	7.9	16	12.5	0.3
Myopic Astigmatism	16	41.0	51	57.3	67	52.3	1.2
Hypermetropic Astigmatism	2	5.1	1	1.1	3	2.3	0.05
Compound Myopic Astigmatism	-		2	2.2	2	1.6	0.04
Compound Hypermetropic Astigmatism	2	5.1	1	1.1	3	2.3	0.05
Mixed Astigmatism	2	5.1	4	4.5	6	4.7	0.11
<b>Total</b>	39	100	89		128	100	
<b>Prevalence (N=5649)</b>	<b>0.7%</b>		<b>1.6%</b>				<b>2.3%</b>

*Prevalence of unilateral refractive errors = 23/1000*

*Prevalence of total unilateral Astigmatism = 15/1000*

**Table 4 :** Distribution of refractive errors – different types  
Bilateral involvements

Type	Right		Left		Total		Prevalence %
	N0	%	N0	%	N0	%	
Simple myopia	64	28.7	66	29.6	130	29.1	1.15
Simple Hypermetropia	41	18.4	34	15.2	75	16.8	0.65
Myopic Astigmatism	83	37.2	91	40.8	174	39.0	1.55
Hypermetropic Astigmatism	9	4.0	10	4.5	19	4.3	0.15
Compound Myopic Astigmatism	17	7.6	14	6.3	31	7.0	0.25
Compound Hypermetropic Astigmatism	-	-	1	0.4	1	0.2	0.01
Mixed Astigmatism	9	4.0	7	3.1	16	3.6	0.15
<b>Total</b>	223	100	223	100	<b>446 eyes</b>		
<b>Prevalence (N=5649)</b>							<b>3.9%</b>

*Prevalence of bilateral refractive errors = 39/1000*

*Prevalence of bilateral Astigmatism = 21/1000*

Table 5 : Average Astigmatism

	Unilateral	Bilateral			Average of Astigmatism unilateral and bilateral
		Right	Left	Average	
Myopic astigmatism	52.3%	37.2%	40.8%	39.0%	45.7%
Hypermetropic astigmatism	2.3%	4.0%	4.5%	4.25%	3.3%
Compound myopic astigmatism	1.6%	7.6%	6.3%	6.95%	4.3%
Compound Hypermetropic astigmatism	2.3%	-	0.4%	0.20%	1.3%
Mixed astigmatism	4.6%	4%	3.1%	3.55%	4.1%
<b>Total astigmatism out of all refractive errors.</b>	<b>63.1%</b>			<b>54.0%</b>	<b>59.0%</b>

#### IV. DISCUSSION

The main aim of this study was to highlight the astigmatism in Sri Lankan children including other types of refractive errors. Uncorrected refractive errors can hamper the performance at schools and lead to various symptoms such as headache, pain in the eyes and aesthenopic symptoms.

A study among children aged 5 – 17 years from Asia, Afro America, Hispanics and Whites revealed that Asians had the highest age and sex adjusted prevalence of myopia (18.5%) and astigmatism (33.6%) (Kleinstejn, Lee, and Zadnik 2003)

A study conducted among 7 – 15 years old children (N = 44.4) in a rural population of India showed that the cause of visual impairment is mainly refractive errors and it was 61%. (Dandona et al 2002)

Another study done in Hyderabad India in children aged 3 – 18 years revealed astigmatism of 10.3% (Kalikirani, Naduvilath, Bantal and Dandora 1997).

Seimon (2005) analysed the prescriptions issued by over 150 optometrists that includes children <15 years over a period of one year and found that 5612 children had sought correction for all kinds of refractive errors.

In this study the total sample of children between 3-14 years was 5649, representing the children of same age group living in southern Sri Lanka. Of them 1233 (21.8%) were selected for further second stage screening using the Snellen's chart and direct examination of the anterior segment. Those selected cases included the first stage screen failures and the difficult to screen cases. For the second stage 98% of referred cases attended and of them 512 children were qualified for the third stage examination. After the examination, 439 cases were confirmed having an ophthalmological diagnosis, giving a prevalence of 7.8% (78/1000).

The analysis of different disease conditions gave the prevalence of Refractive errors of 6.2%(62/1000) and other eye conditions of 1.6% (16/1000).Table 1.

The analysis of the distribution of Refractive errors showed 2.3% (23/1000)of unilateral involvement and 3.9%(39/1000) of bilateral refractive errors in children.-Tables2, 3, & 4.

The prevalence of total simple myopia (bilateral and unilateral) was 1.65% (16.5 / 1000). Simple hypermetropia 0.95 (...1.0% or 10 / 1000)

The astigmatism in children is the main concern of this paper and the the analysis showed a prevalence of unilateral Astigmatism of 1.5% or 15/1000 compared to total unilateral refractive errors of 23/ 1000 in children. The prevalence of bilateral astigmatism was 21/1000 compared to total unilateral refractive errors of 39/1000.In the calculation In the calculation of total Astigmatism, the average of bilateral similar types were considered-Table 5.

The average Astigmatism calculated was 59.0%of all Refractive errors and the most prevalent form of Astigmatism in school children was found to be Myopic Astigmatism, which amounts to 45.7% of all refractive errors. As a prevalence the Myopic Astigmatism alone was 2.8% (28/ 1000 )either unilaterally or bilaterally in school children in southern Sri Lanka making it the most common form of refractive error.

This study highlights that the refractive errors constitute a considerable amount of visual problems among children. Among all refractive errors in Sri Lankan children Astigmatism constitute a bulk. Although Sri Lanka has a well established primary health care system, the burden of refractive errors implies that further consideration in to the issue of screening is needed.

#### REFERENCES RÉFÉRENCES REFERENCIAS

1. Andcea C.T(December 1987) The refractive errors in children. The paediatric clinics of north Americavol. 34, no.6 1425-1438.
2. Dandona R Dandona L. Srinivas M, Sahare P, Narsaiah S, Mun SR, Pokhrael GP and Ellwein LB (2002) Refractive errors in children in a rural

population in India, Investigative ophthalmology and visual Sciences 43 (3) 615 – 622.

3. Deborah, Pavan – Langston (1985) Manual of ocular diagnoses and therapy Asian edition 344-348,
4. Gary Heiting or [www.allaboutvision.com](http://www.allaboutvision.com)
5. Jackson CRS and Finlay RD (1985) the eye in general practice (8<sup>th</sup> ed) Edinburgh, London cherchil Livingstone.
6. Kalkivayi V, Naduvisath TJ Bansal AK, Dandora L (1997) visual impairment in school children in southern India Indian J. ophthalmology 45, 129 – 34.
7. Kleinstien RN, Lee RJ and Zadnik K, (2003) refractive error and ethnicity in children. Arch. Ophthalmology 121, 1148 – 4
8. Resnikoff S, Pascolini D, Mariotti SP, Pokharel GP (2008) Global magnitude of visual impairment caused by uncorrected refractive errors in 2004. Bull world health org 86(1) 63 – 70
9. Seimon R(2005) the distribution of refractive errors in Sri Lankan children. The journal of the college of ophthalmologists of Sri Lanka. 11(1) 24 – 26
10. World health organization (2013). Draft action plan for the prevention of avoidable blindness and visual impairment 2014 – 2019 retrieved from <http://apps.who.int/gb/ebwha/pdffiles/EB132/B132.9.enpdf>