



GLOBAL JOURNAL OF MEDICAL RESEARCH: E  
GYNECOLOGY AND OBSTETRICS  
Volume 15 Issue 1 Version 1.0 Year 2015  
Type: Double Blind Peer Reviewed International Research Journal  
Publisher: Global Journals Inc. (USA)  
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

## Effect Normal Pregnancy and Duration on Liver Enzymes Tests

By Dunia M. R. M.Sc

*University of Kufa, Iraq*

**Abstract-** The current study was designed to investigate the changes of liver enzymes during normal pregnancy. To achieve the intended aim, 185 pregnant women of aged 20 – 37 years (60 women in first trimester, 65 women in second trimester and 60 women in third trimester of pregnancy), also the study contain 70 women (control individuals) in age near to age of pregnant women. The levels of Alanine amino transferase (ALT), Aspartate amino transferase (AST) and Alkaline Phosphatase (ALP) were determined by enzymatic methods. The results indicated a significant ( $P < 0.05$ ) increase of ALT and significant ( $P < 0.01$ ) increase of AST activities in pregnant women in third trimester when compared with those of the control group, while ALP indicated higher significantly ( $P < 0.0005$ ) in third and second trimester when compared with control group.

**Keywords:** *alanine amino transferase (ALT), aspartate transferase (AST) and alkaline phosphatase (ALP).*

**GJMR-E Classification :** *NLMC Code: WJ 190*



*Strictly as per the compliance and regulations of:*



# Effect Normal Pregnancy and Duration on Liver Enzymes Tests

Dunia M. R. M.Sc

**Abstract-** The current study was designed to investigate the changes of liver enzymes during normal pregnancy. To achieve the intended aim, 185 pregnant women of aged 20 – 37 years (60 women in first trimester, 65 women in second trimester and 60 women in third trimester of pregnancy), also the study contain 70 women (control individuals) in age near to age of pregnant women. The levels of Alanine amino transferase (ALT), Aspartate amino transferase (AST) and Alkaline Phosphatase (ALP) were determined by enzymatic methods. The results indicated a significant ( $P < 0.05$ ) increase of ALT and significant ( $P < 0.01$ ) increase of AST activities in pregnant women in third trimester when compared with those of the control group, while ALP indicated higher significantly ( $P < 0.0005$ ) in third and second trimester when compared with control group. The liner regression analysis demonstrated significant ( $r = 0.85$ ,  $P < 0.0005$ ) positive correlation for ALT levels when compared with AST and significant positive correlation for ALT with ALP ( $r = 0.89$ ,  $P < 0.0005$ ) and AST levels with ALP levels ( $r = 0.9$ ,  $P < 0.0005$ ). I am found from my study, the elevated liver enzymes during pregnancy as part of physiological changes.

**Keywords :** alanine amino transferase (ALT), aspartate transferase (AST) and Ikaline phosphatase (ALP).

## I. INTRODUCTION

The liver in the body is the most important organ after the heart. Performing many important functions including metabolism, detoxification and formation of important compounds including blood clotting factors and albumin (16). The pregnant women experiences physiological changes to support fetal growth and development (1,2,3). The levels of estrogens (estradiol) and progesterone increase progressively during pregnancy (4,5). These sex hormones have effects on hepatic metabolic, synthesis, and excretory functions (6,7,8). The biliary excretion of bromosulophthalein decreases during late pregnancy and the clearance of some compounds that are secreted into bile may therefore be impaired (9,10). The phenomenon of hemodilution secondary to the increase in plasma volume decreases the serum protein concentrations. Consequently, certain changes in values of liver function tests occur during normal pregnancy (11,12,13). Pregnancy does not change liver size but in the third trimester the enlarging uterus displaces the liver superiorly and posteriorly, therefore a palpable liver

*Author: Dept. of Bio chemistry, college of medicine, university of Kufa, IRAQ. e-mail: Duniam.mohammedali@uokufa.edu.iq*

disease (14,15). Liver cell injury or necrosis is measured by detergent Glutamate Oxaloacetate Transaminase (AST) and Glutamate Pyruvate Transaminase (ALT) levels (17). While liver synthetic function is quantified by determining albumin level and prothrombin time. Biliary obstruction are elevated by measuring alkaline phosphatase (18). The most commonly used indicators of liver damage (hepatocellular) are the alanine aminotransferase (ALT) and aspartate aminotransferase (AST), formerly referred to as SGPT and SGOT (19). These are enzymes normally found in liver cells that leak out of these cells and make their way to the blood when liver cells are injured. The ALT is felt to be a more specific indicator of liver inflammation as AST is also found in other organs such as the heart and skeletal muscle, the level of the ALT and AST may be used as a general measure of the degree of liver inflammation or damage (19,20). Measurement of serum alanine aminotransferase (ALT) and aspartate aminotransferase (AST) activities levels is the most useful tests for the routine diagnosis of liver diseases (18,19). While serum Alkaline phosphatase (ALP) activity level increase in late pregnancy, mainly during the third trimester.

## II. MATERIALS AND METHODS

Four groups of individuals were included in this study. Group 1 contained 60 pregnant women in first trimester of pregnancy (1 – 3 months). Group 2 consisted of 65 pregnant women in second trimester of pregnancy (4 – 6 months). Group 3 comprised 60 pregnant women of pregnancy (7 – 9 months) and Group 4 contained 70 non pregnant women as control in this study.

Disposable syringes and needles were used for blood collection. Venous blood samples, about 5ml were collected from pregnant and non pregnant women (control group). The blood collected in a polyethylene tubes without anticoagulant, allowed to clot at room temperature for 15 min, blood samples were centrifuged at 3000Xg for 15 min, sera were removed and stored at -17 C until analysis. Labrotary data were obtained by using available kits; serum ALT, serum AST (Randox Kit) and serum ALP (Kind and King). The results were expressed as mean  $\pm$ SD students t test was used for comparison of different groups with controls.

### III. RESULTS

a) Serum ALT, AST and ALP in normal pregnancy during the three times of pregnancy and control group: The characteristics of the study groups are presented in table1 which consists of data of both pregnant women and control group not receiving oral contraception. The results were analyzed using students- test. There was significant ( $P<0.05$ ) increase in ALT activity level during the third trimester ( $9.5\pm3.3$ ) when compared with those of the control group( $7.0\pm2.5$ ), while there was no significant difference in ALT during the

second( $7.8\pm2.8$ ) and first( $7.1\pm2.8$ ) trimester, also the serum AST activity level found to increase significantly ( $P<0.01$ ) during the third trimester ( $38.9\pm4.5$ ) when compared with those of control group( $14.5\pm2.5$ ) and no significant difference in AST in second( $23.7\pm6.1$ ) and first( $18.9\pm3.3$ ) trimester. On the other hand, serum ALP activity level show higher significant ( $P<0.0005$ ) during the third( $379.0\pm70.2$ ) and second( $173.1\pm46.8$ ) trimester when compared with those of the control group( $75.2\pm11.1$ ) and no significant difference in ALP activity level in first( $79.2\pm25.2$ ) trimester.

*Table 1 :* Serum ALT, AST, and ALP in normal pregnancy during the three times of pregnancy and control group.

Parameter	Subjects	NO	Mean $\pm$ SD	P Value
ALT (U/L)	Control	70	7.0 $\pm$ 2.5	
	1 <sup>st</sup> . trimester	60	7.1 $\pm$ 2.8	N.S
	2 <sup>nd</sup> .	65	7.8 $\pm$ 2.8	N.S
	3 <sup>rd</sup> .	60	9.5 $\pm$ 3.3	<0.05
AST (U/L)	Control	70	14.5 $\pm$ 2.5	
	1 <sup>st</sup> .	60	18.9 $\pm$ 3.3	N.S
	2 <sup>nd</sup> .	65	23.7 $\pm$ 6.1	N.S
	3 <sup>rd</sup> .	60	38.9 $\pm$ 4.5	<0.01
ALP(U/L)	Control	70	75.2 $\pm$ 11.1	N.S
	1 <sup>st</sup> .	60	79.2 $\pm$ 25.2	<0.0005
	2 <sup>nd</sup> .	65	173.1 $\pm$ 46.8	<0.0005
	3 <sup>rd</sup> .	60	379.0 $\pm$ 70.2	<0.0005

b) Correlation factors of ALT, AST and ALP levels in normal pregnant women: The linear regression analysis stated significant ( $r=0.85, P<0.0005$ ) positive correlation for ALT with AST activities and

significant positive correlation for ALT activity with ALP activity ( $r=0.89, P<0.0005$ ) and AST activity with ALP activity ( $r=0.9, P<0.0005$ ) in pregnant women (Table2).

*Table 2 :* Correlation Factors of serum ALT, AST and ALP activities in normal pregnant women

Parameter	ALT		AST		ALP	
	r	P	r	P	r	P
ALT			0.85	<0.0005	0.89	<0.0005
AST	0.85	<0.0005			0.9	<0.0005
ALP	0.89	<0.0005	0.9	<0.0005		

c) Influence the duration of pregnancy time on liver enzymes activities. To demonstrate the influence of duration of pregnancy time on ALT, AST and ALP values in pregnant women. As shown in Table3 no significant in ALT and AST activities in second trimester when compared with those of the first trimester, while a significant ( $P<0.01$ ) less elevation of ALP activity in the same comperation. On the

other hand there were significant ( $P<0.0001$ ) increases in ALT, AST and Alp activities levels in third trimester when compared with those of first trimester, the table show also a significant ( $P<0.001$ ) increase in activities levels of AST and ALP during the third trimester when compared with those of second trimester and less elevation in significant ( $P<0.01$ ) for ALT activity.

*Table 3 :* Influence the duration of pregnancy on liver enzymes activities

Parameter	1 <sup>st</sup> Vs 2 <sup>nd</sup> Trimester	1 <sup>st</sup> Vs 3 <sup>rd</sup> Trimester	2 <sup>nd</sup> Vs 3 <sup>rd</sup> Trimester
ALT	N.S	0.001	0.01
AST	N.S	0.001	0.001
ALP	0.01	0.001	0.001

### IV. DISCUSSION

In the present study ALT, AST and ALP activities were measured in 185 healthy pregnant women and 70

control group not receiving oral contraception. None of the women included had evidence of liver disease. When liver cells are damaged or destroyed, the enzymes in the cell leak out into the blood, where they

check the blood for two main liver enzymes ALT and AST(22,23).

In the present investigation that serum ALT activity was significantly higher during the third trimester than in controls ( $P < 0.05$ ). The present results were in agreement with previous works (24, 25), while Bacq et al (12) found that serum ALT activity was significantly higher during the second trimester than in controls but was no different during the third trimester. The current results illustrated that serum AST activity was significantly higher during the third trimester than in controls ( $p < 0.01$ ), two other studies found the same results (14,26), while Bacq et al (12) have stated that serum AST activity was during all three trimesters not significantly higher than in control group. Other study (27) found a significant increase in AST levels between first and third trimester of pregnancy. An increase in ALT and AST levels was found during labor, which might be caused by contractions of uterine muscle (28, 29).

The results indicate that serum ALP activity was significantly higher during the third and second trimesters as compared to control group ( $P < 0.0005$ ). This is primarily due to placental isoenzyme production (30,31). During the third trimester, there was also increase in the production of the bone isoenzyme. The results of this study, showed serum ALT, AST and ALP increased in normal pregnancy as compared to non pregnant women.

## V. CONCLUSIONS

1. The results indicated a significant increase of ALT in pregnant women in third trimester when compared with those of the control group.
2. The levels of AST activity in cease significantly in third trimester when compared with those of control group.
3. The ALP activity indicated higher significantly in third and second trimester when compared with control group.
4. Liver enzymes activities elevated during normal pregnancy.

## VI. ACKNOWLEDGMENTS

First of all, thanks to good for giving me the power and the insistence to complete this work. I want to thank the staff of the AL- Sadder Teaching Hospital in Najaf Governorate for their help during the work. Special thanks are due to Mr. Layth AL faham for this help. I would also like to express my gratitude to my family.

## REFERENCE RÉFÉRENCES REFERENCIAS

1. Black burn ST., Loper DL.: Maternal, fetal and neonatal physiology, A clinical perspective. Philadelphia: WB Saunders, 1992.
2. Campbell E., Dickinson C., Slater J. et al: Clinical Physiology. 5th ed. Black Well. Scientific publication, 1984 ; P 651.
3. Magueda I., Armada E., Diaz J., et al: Practice Guidelines of the Spanith Society of Cardiology for management of cardiac disease in pregnancy, 2000.
4. Taylor D.: Biological Science 2. Oxford university press, New yourk, 1995; P 665.
5. Van thiel D H., Gavaler JS: Pregnancy- associated Sex steroids and their effects on the liver Semin liver Dis, 1987; 7(1): 1- 7.
6. Vigil D. and Gratia P.: Acute fatty liver in pregnancy. Current concepts. Rev. Med. Panama, 2004; 22: 1621.
7. Bacq Y: liver and pregnancy. Patho. Biol, 1999; 47: 958- 965.
8. Marpeau L., Verspyck E., Descarques G.: Management of cholestasis in pregnancy. Press. Med, 1999; 28:2132- 2134.
9. Tindall VR., Beazley JM.: An assessment of changes in liver function during normal pregnancy- using a modified bromsulphthaline test. J Obslet Gynaecol, 1965; 75(5): 717- 737.
10. Ylostalo P. (1970): Liver function in hepatosis of pregnancy and preeclampsia with special refrance to modified bromsulphthaline tests. Acta obstet Gynecol Scand, 1970; 49(supple 4): 1-53.
11. Everson GT.: Liver problems in pregnancy; Distinguishing normal from abnormal hepatic changes. Med. Scope women Health, 1998; 3(2): 3.
12. Bacq Y., Zarka O., et al: Liver function test in normal pregnancy: A prospective study of 103 pregnant women and 103 matched controls. Hepatology, 1996; 23: 1030- 1034.
13. Alonso AG.: Effect of pregnancy on preexisting liver disease physiological changes during pregnancy. Annals of Hepatology, 2006; 5(3): 184- 186.
14. Knopp R., Well P., et al: Clinical chemistry Alterations in pregnancy and contraceptive use. Obstetries and Gynecology, 1998; 66: 682- 690.
15. Hilsden R. and Shaffer E.: Liver disease in pregnancy. Chpter 14, 2005.
16. Hunt CM., Sharara ALA: Liver disease in pregnancy. American Academy of family physician, 1999; Vol 59/No.4.
17. Chopra S., Griffin PH.: Laboratory tests and diagnostic procedures in evaluation of liver disease. A M J med: 1985; 79: 221- 230.
18. Guyton A. and Hall G.: Text book of medical Physiology. 9th ed. W.B. Saunders. Company, 1996; P855.
19. Yonegama K. and Ikeda J: Changes in maternal bone mineral density during pregnancy and relation ship between the density and fofus growth a prospective study. Nippon. Kosu. Eisei, Zasshi, 2000; 47:661- 669.
20. Morgan M., Hainsuonth I. and Kinghom J.: Prospective study on liver dysfunction in pregnancy in south west wales, 2005.

21. 21-Knopp RH. , Bergelin RO.: Clinical chemistry alterations in pregnancy and oral contraceptive use. *Obstet Gynecol*, 1985; 66(5):682- 690.
22. Riely CA: Hepatic disease in pregnancy. *Am j Med* 1994; 96(1):18-22.
23. Samuels p., Cohen Aw.: Pregnancies complicates by liver disease and liver dysfunction. *Obstet Gynecol Clin North Am* 1992; 19:745-763.
24. Cerutti R, Ferraris S, etal: Bahaviour of serum enzymes in pregnancy. *Clin Exp Obstet Gynecol* 1976; 3:22-24.
25. Salgo L, Pal A: Variation in some enzymes in amniotic fluid and maternal serum during pregnancy. *Enzyme* 1989; 41:101-107.
26. Elliott JR, Kell RT: Normal Clinical values for pregnant women at term. *Clin Chem* 1971; 17:156-157.
27. Guntupalli RS, Steingrub J: Hepatic disease and pregnancy: An over view of diagnosis and management. *Crit Care Med*: 2005;33(10. Suppl.):332-339.
28. Meade BW, Rosalki SB: Serum enzyme activity in normal pregnancy and newborn. *J Obstet Gynaecol* 1993; 70 : 693-700.
29. Loganathan G, Rachel G, etal: Liver function tests in normal pregnancy: a study from Southern India, *Indian J of gastroenterology* 2005; 24(6):268-269.
30. Siogren MH: Hepatic emergencies in pregnancy. *Med Clin North Am* 1993; 77:1115-1127.
31. Smoleniec JS, James DK: Gastro intestinal crises during pregnancy. *Dig Dis* 1993; 11:313-324.