

# GLOBAL JOURNAL

OF MEDICAL RESEARCH: G

## Veterinary Science & Veterinary Medicine

Sero-Prevalence of Toxoplasma

Prevalence of Bovine Cysticercosis

Highlights

Prevalence of Fasciola Infection

Major Causes of Liver Condemnation

Discovering Thoughts, Inventing Future

VOLUME 16    ISSUE 3    VERSION 1.0



GLOBAL JOURNAL OF MEDICAL RESEARCH: G  
VETERINARY SCIENCE AND VETERINARY MEDICINE

---



GLOBAL JOURNAL OF MEDICAL RESEARCH: G  
VETERINARY SCIENCE AND VETERINARY MEDICINE

---

VOLUME 16 ISSUE 3 (VER. 1.0)

OPEN ASSOCIATION OF RESEARCH SOCIETY

© Global Journal of Medical Research. 2016.

All rights reserved.

This is a special issue published in version 1.0 of "Global Journal of Medical Research." By Global Journals Inc.

All articles are open access articles distributed under "Global Journal of Medical Research"

Reading License, which permits restricted use. Entire contents are copyright by of "Global Journal of Medical Research" unless otherwise noted on specific articles.

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopy, recording, or any information storage and retrieval system, without written permission.

The opinions and statements made in this book are those of the authors concerned. Ultraculture has not verified and neither confirms nor denies any of the foregoing and no warranty or fitness is implied.

Engage with the contents herein at your own risk.

The use of this journal, and the terms and conditions for our providing information, is governed by our Disclaimer, Terms and Conditions and Privacy Policy given on our website <http://globaljournals.us/terms-and-condition/menu-id-1463/>

By referring / using / reading / any type of association / referencing this journal, this signifies and you acknowledge that you have read them and that you accept and will be bound by the terms thereof.

All information, journals, this journal, activities undertaken, materials, services and our website, terms and conditions, privacy policy, and this journal is subject to change anytime without any prior notice.

Incorporation No.: 0423089  
License No.: 42125/022010/1186  
Registration No.: 430374  
Import-Export Code: 1109007027  
Employer Identification Number (EIN):  
USA Tax ID: 98-0673427

## Global Journals Inc.

(A Delaware USA Incorporation with "Good Standing"; Reg. Number: 0423089)  
Sponsors: Open Association of Research Society  
Open Scientific Standards

### *Publisher's Headquarters office*

Global Journals® Headquarters  
945th Concord Streets,  
Framingham Massachusetts Pin: 01701,  
United States of America  
USA Toll Free: +001-888-839-7392  
USA Toll Free Fax: +001-888-839-7392

### *Offset Typesetting*

Global Journals Incorporated  
2nd, Lansdowne, Lansdowne Rd., Croydon-Surrey,  
Pin: CR9 2ER, United Kingdom

### *Packaging & Continental Dispatching*

Global Journals  
E-3130 Sudama Nagar, Near Gopur Square,  
Indore, M.P., Pin: 452009, India

### *Find a correspondence nodal officer near you*

To find nodal officer of your country, please  
email us at [local@globaljournals.org](mailto:local@globaljournals.org)

### *eContacts*

Press Inquiries: [press@globaljournals.org](mailto:press@globaljournals.org)  
Investor Inquiries: [investors@globaljournals.org](mailto:investors@globaljournals.org)  
Technical Support: [technology@globaljournals.org](mailto:technology@globaljournals.org)  
Media & Releases: [media@globaljournals.org](mailto:media@globaljournals.org)

### *Pricing (Including by Air Parcel Charges):*

#### *For Authors:*

22 USD (B/W) & 50 USD (Color)  
Yearly Subscription (Personal & Institutional):  
200 USD (B/W) & 250 USD (Color)

# GLOBAL JOURNALS CONSTITUTIONAL EDITORIAL BOARD

~INTEGRATED~

## *Dr. Charles A. Rarick*

Ph.D.  
Professor of International Business  
College of Business  
Purdue University Northwest  
Hammond, Indiana USA

## *Dr. Osman Balci, Professor*

Department of Computer Science  
Virginia Tech, Virginia University  
Ph.D. and M.S. Syracuse University, Syracuse, New York  
M.S. and B.S. Bogazici University, Istanbul, Turkey  
Web: [manta.cs.vt.edu/balci](http://manta.cs.vt.edu/balci)

## *Dr. A. Heidari*

Ph.D, D.Sc, Faculty of Chemistry  
California South University (CSU),  
United Stated

## *Dr. Miklas Scholz*

B.Eng. (equiv), PgC, MSc, Ph.D, CWEM, C.Env., CSci,  
C.Eng.  
Nigeria Health, Wellness and Fitness  
University of Lund

## *Dr. Maria Gullo*

Ph.D, Food Science and Technology  
University of Catania  
Department of Agricultural and Food Sciences  
University of Modena and Reggio Emilia, Italy

## *Dr. Qiang Wu*

Ph.D University of Technology, Sydney  
Department of Mathematics,  
Physics and Electrical Engineering  
Northumbria University

## *Dr. Bingyun Li*

Ph.D Fellow, IAES  
Guest Researcher, NIOSH, CDC, Morgantown, WV  
Institute of Nano and Biotechnologies  
West Virginia University, US

## *Dr. Audeh Ahmad Ahmad*

Amman Arab University For Higher Education  
Ph.D, Accounting-Ais  
Faculty of Business Administration  
Alalbyt University, Jordan, Amman

## *Dr. Lucian Baia*

Ph.D Julius-Maximilians University Würzburg, Germany  
Associate professor  
Department of Condensed Matter Physics and  
Advanced Technologies, Babes-Bolyai University,  
Romania

## *Dr. Sahraoui Chaieb*

PhD Physics and Chemical Physics  
M.S. Theoretical Physics  
B.S. Physics, École Normale Supérieure, Paris  
Associate Professor, Bioscience  
King Abdullah University of Science and Technology

## *Dr. Houfa Shen*

Ph.D Manufacturing Engineering,  
Mechanical Engineering, Structural Engineering  
Department of Mechanical Engineering  
Tsinghua University, China

## *Dr. Arshak Poghossian*

Ph.D Solid-State Physics  
Leningrad Electrotechnic Institute, Russia  
Institute of Nano and Biotechnologies  
Aachen University of Applied Sciences, Germany

*Dr. A. Stegou-Sagia*

Ph.D Mechanical Engineering, Environmental  
Engineering School of Mechanical Engineering  
National Technical University of Athens

*Giuseppe A Provenzano*

Irrigation and Water Management, Soil Science,  
Water Science Hydraulic Engineering  
Dept. of Agricultural and Forest Sciences  
Universita di Palermo, Italy

*Dr. Ciprian LĂPUȘAN*

Ph. D in Mechanical Engineering  
Technical University of Cluj-Napoca  
Cluj-Napoca (Romania)

*Dr. Haijian Shi*

Ph.D Civil Engineering Structural Engineering  
Oakland, CA, United States

*Dr. Yogita Bajpai*

Ph.D Senior Aerospace/Mechanical/  
Aeronautical Engineering professional  
M.Sc. Mechanical Engineering  
M.Sc. Aeronautical Engineering  
B.Sc. Vehicle Engineering  
Orange County, California, USA

*Dr. Abdurrahman Arslanyilmaz*

Computer Science & Information Systems Department  
Youngstown State University  
Ph.D., Texas A&M University  
University of Missouri, Columbia  
Gazi University, Turkey  
Web:[cis.yzu.edu/~aarslanyilmaz/professional\\_web](http://cis.yzu.edu/~aarslanyilmaz/professional_web)

*Dr. Chao Wang*

Ph.D. in Computational Mechanics  
Rosharon, TX, USA

*Dr. Adel Al Jumaily*

Ph.D Electrical Engineering (AI)  
Faculty of Engineering and IT  
University of Technology, Sydney

*Kitipong Jaojaruek*

B. Eng, M. Eng D. Eng (Energy Technology, Asian  
Institute of Technology).  
Kasetsart University Kamphaeng Saen (KPS) Campus  
Energy Research Laboratory of Mechanical Engineering

*Dr. Mauro Lenzi*

Ph.D, Biological Science, Pisa University, Italy  
Lagoon Ecology and Aquaculture Laboratory  
Orbetello Pesca Lagunare Company

*Dr. Omid Gohardani*

M.Sc. (Computer Science), FICCT, U.S.A.  
Email: [yogita@computerresearch.org](mailto:yogita@computerresearch.org)

*Dr. Yap Yee Jiun*

B.Sc.(Manchester), Ph.D.(Brunel), M.Inst.P.(UK)  
Institute of Mathematical Sciences,  
University of Malaya,  
Kuala Lumpur, Malaysia

*Dr. Thomas Wischgoll*

Computer Science and Engineering,  
Wright State University, Dayton, Ohio  
B.S., M.S., Ph.D.  
(University of Kaiserslautern)  
Web:[avida.cs.wright.edu/personal/wischgol/index\\_eng.html](http://avida.cs.wright.edu/personal/wischgol/index_eng.html)

*Dr. Baziotis Ioannis*

Ph.D. in Petrology-Geochemistry-Mineralogy  
Lipson, Athens, Greece

*Dr. Xiaohong He*

Professor of International Business  
University of Quinnipiac  
BS, Jilin Institute of Technology; MA, MS, Ph.D,  
(University of Texas-Dallas)  
Web: [quinnipiac.edu/x1606.xml](http://quinnipiac.edu/x1606.xml)

*Dr. Burcin Becerik-Gerber*

University of Southern Californi  
Ph.D in Civil Engineering  
DDes from Harvard University  
M.S. from University of California, Berkeley  
M.S. from Istanbul Technical University  
Web: [i-lab.usc.edu](http://i-lab.usc.edu)

*Dr. Söhnke M. Bartram*

Department of Accounting and Finance  
Lancaster University Management School  
Ph.D. (WHU Koblenz)  
MBA/BBA (University of Saarbrücken)  
Web: [lancs.ac.uk/staff/bartras1/](http://lancs.ac.uk/staff/bartras1/)

*Dr. Söhnke M. Bartram*

Ph.D, (IT) in Faculty of Engg. & Tech.  
Professor & Head,  
Dept. of ISE at NMAM Institute of Technology

*Dr. Balasubramani R*

Department of Accounting and Finance  
Lancaster University Management School  
Ph.D. (WHU Koblenz)  
MBA/BBA (University of Saarbrücken)  
Web: [lancs.ac.uk/staff/bartras1/](http://lancs.ac.uk/staff/bartras1/)

*M. Meguellati*

Department of Electronics,  
University of Batna, Batna 05000, Algeria

*Dr. T. David A. Forbes*

Associate Professor and Range Nutritionist  
Ph.D Edinburg University - Animal Nutrition  
M.S. Aberdeen University - Animal Nutrition  
B.A. University of Dublin- Zoology.  
Web: [essm.tamu.edu/people-info/faculty/forbes-david](http://essm.tamu.edu/people-info/faculty/forbes-david)

*Dr. Bassey Benjamin Esu*

B.Sc. Marketing; MBA Marketing; Ph.D Marketing  
Lecturer, Department of Marketing, University of Calabar  
Tourism Consultant, Cross River State Tourism  
Development Department  
Co-rdinator , Sustainable Tourism Initiative, Calabar,  
Nigeria

*Dr. Maciej Gućma*

Asistant Professor,  
Maritime University of Szczecin Szczecin, Poland  
Ph.D. Eng. Master Mariner  
Web: [www.mendeley.com/profiles/maciej-gucma/](http://www.mendeley.com/profiles/maciej-gucma/)

*Dr. Shun-Chung Lee*

Department of Resources Engineering,  
National Cheng Kung University, Taiwan

*Dr. Fotini Labropulu*

Mathematics - Luther College, University of Regina  
Ph.D, M.Sc. in Mathematics  
B.A. (Honours) in Mathematics, University of Windsor  
Web: [luthercollege.edu/Default.aspx](http://luthercollege.edu/Default.aspx)

*Dr. Vesna Stanković Pejnović*

Ph. D. Philosohy , Zagreb, Croatia  
Rusveltova, Skopje, Macedonia

*Dr. Miguel Angel Ariño*

Professor of Decision Sciences  
IESE Business School  
Barcelona, Spain (Universidad de Navarra)  
CEIBS (China Europe International Business School).  
Beijing, Shanghai and Shenzhen  
Ph.D. in Mathematics, University of Barcelona  
BA in Mathematics (Licenciatura)  
University of Barcelona  
Web: [web.iese.edu/MAArino/overview.axd](http://web.iese.edu/MAArino/overview.axd)

*Dr. Philip G. Moscoso*

Technology and Operations Management  
IESE Business School, University of Navarra  
Ph.D in Industrial Engineering and Management,  
ETH Zurich , M.Sc. in Chemical Engineering,  
ETH Zurich Link: Philip G. Moscoso personal webpage

*Dr. Mihaly Mezei*

Associate Professor  
Department of Structural and Chemical Biology  
Mount Sinai School of Medical Center  
Ph.D., Etsv Lornd University, Postdoctoral Training,  
New York University, MSSM home:  
<https://www.mountsinai.org/Find%20A%20Faculty/profile.do?id=0000072500001497192632>  
Lab home - software,  
publications: <https://inka.mssm.edu/~mezei>  
Department: <https://atlas.physbio.mssm.edu>

*Dr. Vivek Dubey (HON.)*

MS (Industrial Engineering),  
MS (Mechanical Engineering)  
University of Wisconsin  
FICCT  
Editor-in-Chief, USA

*Dr. Carlos García Pont*

Associate Professor of Marketing  
IESE Business School, University of Navarra  
Doctor of Philosophy (Management),  
Massachusetts Institute of Technology (MIT)  
Master in Business Administration, IESE,  
University of Navarra  
Degree in Industrial Engineering,  
Universitat Politècnica de Catalunya  
Web: [iese.edu/aplicaciones/faculty/facultyDetail.asp](http://iese.edu/aplicaciones/faculty/facultyDetail.asp)

*Dr. Sanjay Dixit, M.D.*

Director, EP Laboratories, Philadelphia VA Medical Center  
Cardiovascular Medicine - Cardiac Arrhythmia  
University of Penn School of Medicine  
Web: [pennmedicine.org/wagform/MainPage.aspx?](http://pennmedicine.org/wagform/MainPage.aspx?)

*Dr. Pina C. Sanelli*

Associate Professor of Radiology  
Associate Professor of Public Health  
Weill Cornell Medical College  
Associate Attending Radiologist  
NewYork-Presbyterian Hospital  
MRI, MRA, CT, and CTA  
Neuroradiology and Diagnostic Radiology  
M.D., State University of New York at Buffalo,  
School of Medicine and Biomedical Sciences  
Web: [weillcornell.org/pinasanelli/](http://weillcornell.org/pinasanelli/)

*Er. Suyog Dixit*

(M.Tech), BE (HONS. in CSE), FICCT  
SAP Certified Consultant  
CEO at IOSRD, GAOR OSS  
Technical Dean, Global Journals Inc.(US)  
Website: [www.suyogdixit.com](http://www.suyogdixit.com)  
Email: [suyog@suyogdixit.com](mailto:suyog@suyogdixit.com)



*Er. Pritesh Rajvaidya*

Computer Science Department  
California State University  
BE (Computer Science), FICCT  
Technical Dean, USA  
Email: [pritesh@computerresearch.org](mailto:pritesh@computerresearch.org),  
[deanusa@globaljournals.org](mailto:deanusa@globaljournals.org)

*Dr. Apostolos Ch. Zarros*

DM, Degree (Ptychio) holder in Medicine,  
National and Kapodistrian University of Athens  
MRes, Master of Research in Molecular Functions in  
Disease,  
University of Glasgow  
FRNS, Fellow, Royal Numismatic Society  
Member, European Society for Neurochemistry  
Member, Royal Institute of Philosophy  
Scotland, United Kingdom

*Dr. Han-Xiang Deng*

MD., Ph.D  
Associate Professor and Research Department  
Division of Neuromuscular Medicine  
Davee Department of Neurology and Clinical  
Neurosciences  
Northwestern University Feinberg School of Medicine  
Web: [neurology.northwestern.edu/faculty/deng.html](http://neurology.northwestern.edu/faculty/deng.html)

*Dr. Roberto Sanchez*

Associate Professor  
Department of Structural and Chemical Biology  
Mount Sinai School of Medicine  
Ph.D., The Rockefeller University  
Web: [mountsinai.org/](http://mountsinai.org/)

*Jixin Zhong*

Department of Medicine,  
Affiliated Hospital of Guangdong Medical College,  
Zhanjiang, China Davis Heart and Lung Research Institute,  
The Ohio State University, Columbus, OH 43210, USA

*Dr. Wen-Yih Sun*

Professor of Earth and Atmospheric Sciences  
Purdue University, Director  
National Center for Typhoon and Flooding Research,  
Taiwan  
University Chair Professor  
Department of Atmospheric Sciences,  
National Central University, Chung-Li, Taiwan  
University Chair Professor  
Institute of Environmental Engineering,  
National Chiao Tung University, Hsin-chu, Taiwan.  
Ph.D., MS The University of Chicago, Geophysical Sciences  
BS National Taiwan University, Atmospheric Sciences  
Web: [event.nchc.org.tw/2009](http://event.nchc.org.tw/2009)

*Dr. Michael R. Rudnick*

M.D., FACP  
Associate Professor of Medicine  
Chief, Renal Electrolyte and Hypertension Division (PMC)  
Penn Medicine, University of Pennsylvania  
Presbyterian Medical Center, Philadelphia  
Nephrology and Internal Medicine  
Certified by the American Board of Internal Medicine  
Web: [uhs.upenn.edu/](http://uhs.upenn.edu/)

*Dr. Aziz M. Barbar, Ph.D.*

IEEE Senior Member  
Chairperson, Department of Computer Science  
AUST - American University of Science & Technology  
Alfred Naccash Avenue - Ashrafieh

*Dr. Minghua He*

Department of Civil Engineering  
Tsinghua University  
Beijing, 100084, China

*Anis Bey*

Dept. of Comput. Sci.,  
Badji Mokhtar-Annaba Univ.,  
Annaba, Algeria

*Chutisant Kerdvibulvech*

Dept. of Inf.& Commun. Technol.,  
Rangsit University, Pathum Thani, Thailand  
Chulalongkorn University, Thailand  
Keio University, Tokyo, Japan

*Dr. Wael Abdullah*

Elhelece Lecturer of Chemistry,  
Faculty of science, Gazan Univeristy,  
KSA. Ph. D. in Inorganic Chemistry,  
Faculty of Science, Tanta University, Egypt

*Yaping Ren*

School of Statistics and Mathematics  
Yunnan University of Finance and Economics  
Kunming 650221, China

*Ye Tian*

The Pennsylvania State University  
121 Electrical Engineering East  
University Park, PA 16802, USA

*Dr. Diego González-Aguilera*

Ph.D. Dep. Cartographic and Land Engineering,  
University of Salamanca, Ávila, Spain

*Dr. Hai-Linh Tran*

PhD in Biological Engineering  
Department of Biological Engineering  
College of Engineering Inha University, Incheon, Korea

*Dr. Tao Yang*

Ph.D, Ohio State University  
M.S. Kansas State University  
B.E. Zhejiang University

*Dr. Feng Feng*

Boston University  
Microbiology, 72 East Concord Street R702  
Duke University  
United States of America

*Shengbing Deng*

Departamento de Ingeniería Matemática,  
Universidad de Chile.  
Facultad de Ciencias Físicas y Matemáticas.  
Blanco Encalada 2120, piso 4.  
Casilla 170-3. Correo 3. - Santiago, Chile

*Claudio Cuevas*

Department of Mathematics  
Universidade Federal de Pernambuco  
Recife PE Brazil

*Dr. Alis Puteh*

Ph.D. (Edu.Policy) UUM  
Sintok, Kedah, Malaysia  
M.Ed (Curr. & Inst.), University of Houston, USA

*Dr. R.K. Dixit(HON.)*

M.Sc., Ph.D., FICCT Chief Author, India  
Email: [authorind@globaljournals.org](mailto:authorind@globaljournals.org)

*Dr. Dodi Irawanto*

PhD, M.Com, B.Econ Hons.  
Department of Management,  
Faculty of Economics and Business, Brawijaya University  
Malang, Indonesia

*Ivona Vrdoljak Raguz*

University of Dubrovnik, Head,  
Department of Economics and Business Economics,  
Croatia

*Dr. Prof Adrian Armstrong*

BSc Geography, LSE, 1970  
PhD Geography (Geomorphology)  
Kings College London 1980  
Ordained Priest, Church of England 1988  
Taunton, Somerset, United Kingdom

*Thierry FEUILLET*

Géolittomer – LETG UMR 6554 CNRS  
(Université de Nantes)  
Institut de Géographie et d'Aménagement  
Régional de l'Université de Nantes.  
Chemin de la Censive du Tertre – BP, Rodez

*Dr. Yongbing Jiao*

Ph.D. of Marketing  
School of Economics & Management  
Ningbo University of Technology  
Zhejiang Province, P. R. China

*Cosimo Magazzino*

Roma Tre University  
Rome, 00145, Italy

*Dr. Shaoping Xiao*

BS, MS, Ph.D Mechanical Engineering,  
Northwestern University  
The University of Iowa  
Department of Mechanical and Industrial Engineering  
Center for Computer-Aided Design

*Dr. Alex W. Dawotola*

Hydraulic Engineering Section,  
Delft University of Technology,  
Stevinweg, Delft, Netherlands

*Dr. Luisa dall'Acqua*

PhD in Sociology (Decisional Risk sector),  
Master MU2, College Teacher in Philosophy (Italy),  
Edu-Research Group, Zürich/Lugano

*Xianghong Qi*

University of Tennessee  
Oak Ridge National Laboratory  
Center for Molecular Biophysics  
Oak Ridge National Laboratory  
Knoxville, TN 37922, United States

*Gerard G. Dumancas*

Postdoctoral Research Fellow,  
Arthritis and Clinical Immunology Research Program,  
Oklahoma Medical Research Foundation  
Oklahoma City, OK  
United States

*Vladimir Burtman*

Research Scientist  
The University of Utah, Geophysics  
Frederick Albert Sutton Building, 115 S 1460 E Room 383  
Salt Lake City, UT 84112, USA

*Jalal Kafashan*

Mechanical Engineering, Division of Mechatronics  
KU Leuven, BELGIUM

*Zhibin Lin*

Center for Infrastructure Engineering Studies  
Missouri University of Science and Technology  
ERL, 500 W. 16th St. Rolla,  
Missouri 65409, USA

*Dr. Lzzet Yavuz*

MSc, PhD, D Ped Dent.  
Associate Professor,  
Pediatric Dentistry Faculty of Dentistry,  
University of Dicle, Diyarbakir, Turkey

*Prof. Dr. Eman M. Gouda*

Biochemistry Department,  
Faculty of Veterinary Medicine, Cairo University,  
Giza, Egypt

*Della Ata*

BS in Biological Sciences  
MA in Regional Economics  
Hospital Pharmacy  
Pharmacy Technician Educator

*Dr. Muhammad Hassan Raza, PhD*

Engineering Mathematics  
Internetworking Engineering, Dalhousie University,  
Canada

*Dr. Asunción López-Varela*

BA, MA (Hons), Ph.D (Hons)  
Facultad de Filología.  
Universidad Complutense Madrid  
29040 Madrid, Spain

*Dr. Bondage Devanand Dhondiram*

Ph.D  
No. 8, Alley 2, Lane 9, Hongdao station,  
Xizhi district, New Taipei city 221, Taiwan (ROC)

*Dr. Latifa Oubedda*

National School of Applied Sciences,  
University Ibn Zohr, Agadir, Morocco  
Lotissement Elkhier N°66  
Bettana Salé Maroc

*Dr. Belen Riverio, PhD*

School of Industrial Enigneering  
University of Vigo  
Spain

## CONTENTS OF THE ISSUE

---

- i. Copyright Notice
- ii. Editorial Board Members
- iii. Chief Author and Dean
- iv. Contents of the Issue
  1. Prevalence of *Fasciola* Infection in Slaughtered Animals in Kashmir. **1-10**
  2. Sero-Prevalence of *Toxoplasma Gondii* Infection and Associated Risk Factors in Animals Presented to Sholla and Akaki-Kality Veterinary Clinics, Addis Ababa. **11-17**
  3. Prevalence of Bovine Cysticercosis and Status of Human Taeniasis in and Around Asella Town, Tiyyoworeda, South East Ethiopia. **19-26**
  4. Epidemiological Study of Small Ruminant Diseases in Selected Districts of Kaffa and Bench- Maji Zone, Southern Nations Nationalities and Peoples Regional State (SNNPRs), Ethiopia. **27-37**
  5. Major Causes of Liver Condemnation and Associated Direct Financial Losses in Bovine Saughtered at Assela Municipal Abattoir Arsi, South Eastern Ethiopia. **39-44**
- v. Fellows
- vi. Auxiliary Memberships
- vii. Process of Submission of Research Paper
- viii. Preferred Author Guidelines
- ix. Index



GLOBAL JOURNAL OF MEDICAL RESEARCH: G  
VETERINARY SCIENCE AND VETERINARY MEDICINE  
Volume 16 Issue 3 Version 1.0 Year 2016  
Type: Double Blind Peer Reviewed International Research Journal  
Publisher: Global Journals Inc. (USA)  
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

## Prevalence of *Fasciola* Infection in Slaughtered Animals in Kashmir

By Nazima Gul, Hidayatullah Tak, Khalid M. Fazilli, Iram Abdullah & Tanveer A. Sofi

*University of Kashmir*

**Abstract-** Fasciolosis is denoted as a significant veterinary health problem. During current study, a total of 714 cattle slaughtered at different abattoirs of Srinagar city (J&K) were examined for the presence of *Fasciola* sps in the liver from January 2014 to January 2016. There was moderate prevalence of 26.84% in the studied area. Predominance of *Fasciola gigantica* (20.86%) was seen as compared to *Fasciola hepatica* (3.361%) infection with mixed infection of 2.66%. Epidemiological determinants like age, gender, breed and body condition showed statistically significant ( $p < 0.05$ ) effect on bovine Fasciolosis. Seasonal data showed highest prevalence in autumn (39.87%) followed by winter (28.84%) with lowest prevalence in spring (16.40%).

**Keywords:** epidemiology, fasciola, cattle, abattoir and srinagar.

**GJMR-G Classification:** NLMC Code: WC 900



*Strictly as per the compliance and regulations of:*



© 2016. Nazima Gul, Hidayatullah Tak, Khalid M. Fazilli, Iram Abdullah & Tanveer A. Sofi. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License <http://creativecommons.org/licenses/by-nc/3.0/>), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

# Prevalence of *Fasciola* Infection in Slaughtered Animals in Kashmir

Nazima Gul <sup>α</sup>, Hidayatullah Tak <sup>σ</sup>, Khalid M. Fazilli <sup>ρ</sup>, Iram Abdullah <sup>ω</sup> & Tanveer A. Sofi <sup>¥</sup>

**Abstract-** Fasciolosis is denoted as a significant veterinary health problem. During current study, a total of 714 cattle slaughtered at different abattoirs of Srinagar city (J&K) were examined for the presence of *Fasciola* sps in the liver from January 2014 to January 2016. There was moderate prevalence of 26.84% in the studied area. Predominance of *Fasciola gigantica* (20.86%) was seen as compared to *Fasciola hepatica* (3.361%) infection with mixed infection of 2.66%. Epidemiological determinants like age, gender, breed and body condition showed statistically significant ( $p < 0.05$ ) effect on bovine Fasciolosis. Seasonal data showed highest prevalence in autumn (39.87%) followed by winter (28.84%) with lowest prevalence in spring (16.40%).

**Keywords:** epidemiology, fasciola, cattle, abattoir and srinagar.

## I. INTRODUCTION

Ruminant productivity around the world is majorly affected by trematode parasitism (Vercruysse and Claerebout 2001). Among them, Fasciolosis gains public concern not only due to its prevalence and

economic significance to animal stock in all continents (Schweizer *et al.*, 2005, Mungube *et al.*, 2006) but also to its zoonotic aspect. Bovine Fasciolosis is an impediment in profitable bovine farming and for butchers and consumers too. Parasite of genus *Fasciola* i.e *Fasciola hepatica* and *Fasciola gigantica* is the causative agent of Fasciolosis which occur in a wide range of definitive hosts. Over the last decade there has been a substantial increase in the number of fasciolosis cases recorded. It is spurred on by both environmental changes (warmer, wetter climate) and man-made modifications such as an increase in animal movements and intensification of livestock farming (Mas-Coma *et al.*, 2005).

According to Annual Reports of Department of Animal Husbandry, Dairying and Fisheries, species-wise incidence of Bovine Fasciolosis in India is tabulated as under:

Year	Outbreaks	Attacks	Deaths
2008-2009	85	391402	2
2009-2010	84	375237	6
2010-2011	105	345108	27
2011-2012	130	316363	74
2012-2013	195	509195	31
2013-2014	137	802698	11
2014-2015	129	3606	4

While comparing the apparent prevalence of liver fluke infection, detected by liver, faeces and bile examination it has been reported that examination of liver or bile samples was more sensitive than faecal examination (Braun *et al.*, 1995 and Kumar *et al.*, 2002). Thus the abattoir study was carried out to determine the prevalence.

## II. MATERIAL AND METHODS

A two-year prospective systematic sampling study was undertaken from January 2014 to January 2016 to determine the relative occurrence of *Fasciola* infection in the livers of cattle presented to six abattoirs across the Kashmir. Samples were taken from the three studied localities i.e., Hazratbal, Parimpoora, and

Gouskimber of Srinagar district but sampling effort was more important in Parimpoora locality, where four slaughterhouses were closely located.

The sample size was calculated using the formula given by Thrustfield, M. (2005).

$$n = \frac{1.96^2 \cdot P_{\text{exp}} (1 - P_{\text{exp}})}{d^2}$$

Where n = required sample size

$P_{\text{exp}}$  = expected prevalence = 50%

d = desired absolute precision = 5%

Hence, d = 0.05 and p = 0.5 (50%).

The expected prevalence in the study area was 55 % (Akhoun and Peer, 2014). Thus the minimum desired annual sample size was calculated to 381. However, due to drastic floods only 316 cattle were examined in Year 2014 as collection areas were inaccessible and sample size was extended to 396 in Year 2015.

Author <sup>α</sup> <sup>σ</sup> <sup>ω</sup> <sup>¥</sup>: Department of Zoology, University of Kashmir, Jammu and Kashmir -190006, India. e-mail: nazimagul1@gmail.com

Author <sup>ρ</sup>: Department of Biotechnology, University of Kashmir, Jammu and Kashmir -190006, India.

### III. STUDY OF EPIDEMIOLOGICAL PARAMETES

#### a) *Antimortem analysis*

- Age, Gender and breed of animal

The age of each animal was confirmed by looking at the physical appearance of body and examining the dental pad and incisor teeth (Cockrill, 1974). The data was collected according to predesigned proforma: Young (1Yr-3Yrs), adult (3-6Yrs) and aged (Above 6 years). During survey the gender and breed of animals was also recorded.

- Assessment of Body condition

Body scoring of the cattle was made based on the method described by Nicholson and Butterworth (1986). Each scoring were given number from 1(L-, very lean) to 9 (F+, very fat) and these scores finally included under three body condition scores, good, medium and poor.

- Season

On the basis of temperature and precipitation, four seasons in a year recognized in Kashmir valley are: winter (December to February); spring (March to May); summer (June to August); autumn (September to November) (Dar *et al.*, 2002).

#### b) *Postmortem examination*

- Types of infection

Infection based on causative agent were classified as *Fasciola hepatica*, *Fasciola gigantica*, mixed *Fasciola* species (*Fasciola hepatica*, *Fasciola gigantica*) infection.

#### c) *Postmortem fluke recovery*

Worms were recovered from infected livers by squeezing them manually to macerate the parenchyma and the flukes were carefully removed and placed in petridish containing 0.15M Dubecco's PBS buffer (pH 7.3) for initial washing. The flukes were stored in

collection vials containing PBS and were transported to the laboratory of Department of Zoology, University of Kashmir, Srinagar. Fasciolids were identified primarily on differences in body shape and size of the adults, with the smaller *F. hepatica* exhibiting wide and defined shoulders compared to the slender *F. gigantica* having less defined shoulders and shorter cephalic cones (Soulsby, 1986). For permanent slide preparation flukes were rapidly killed in 70% ethyl alcohol to avoid shrinkage. The flukes were then transferred to vials containing 6-10% formalin for preservation. Flukes were stained with Borax Carmine, dehydrated in ascending grades of ethanol, cleared in Xylene and mounted in Balsam Canada and viewed under monocular light microscope.

#### d) *Data Analysis*

Data was recorded, entered and managed into MS Excel work sheet and analyzed using Minitab Version 13. Prevalence was calculated as percentage of infected among the examined samples. Chi square test was employed to examine the effect of above mentioned epidemiological determinants on the level of parasitism in host. In all statistical analysis, confidence level was held at 95% and P-value is <0.05 (at 5% level of significance) was considered as significant.

### IV. RESULTS

Fasciolosis in an area is influenced by a multifactorial system which comprises both definitive and intermediate hosts, parasite and environmental effects. Numerous factors (both intrinsic and extrinsic) form an association posing a potential epidemiological threat and it is important that the existence and localization of such an association should be recognized beforehand so that the situation can be brought under control. Thus in this portion of result, these factors have be assessed and potential reason behind the association have been well documented




Pg 1: *Fasciola hepatica*



Pg 2: *Fasciola gigantica*





**Pg 1:** Infected Liver Samples. (A) Bovine liver showing oozing parasite. (B, C) Parasites lying free and soaked in foul smelling fluid (D) Yellowish tissue

**Overall Prevalence (Table 1)**

The overall prevalence of Fasciolosis for the period of two years (2014-2015) was found to be 26.84% in the current study areas. In 2015, the percentage prevalence was higher (27.02%) than in 2014 (25.31%). There was an increase of 1.71% in prevalence rate from 2014 to 2015. But difference in prevalence rate was not statistically significant ( $p > 0.05$ )

as there was sampling error in year 2014 because of scarcity of data collection for a period of 2 months (September and October) due to Floods that affected the whole valley.

The result of current study indicated that Fasciolosis in cattle is spread relatively with moderate prevalence rate of 26.84% in the study area as compared to high prevalence of 51.42%, 42.06% and

43.63% in Ladakh and Srinagar province of Jammu and Kashmir (Kuchai *et al.* 2011; Akhoun and Peer, 2014 respectively). The reported difference may be attributed to different factors like mode of infestation, agroclimatic variations, technique used for data collection and different management conditions under which cattle are reared. However, the result of the present study is in close proximity to the prevalence rates of 29.38% and 25.40% reported earlier by Sheikh *et al.* 2007 and Fatima *et al.* 2012 in neighboring areas of Kashmir using the same abattoir survey. The prevalence rate was also within ranges of findings of other authors like 25.46% by Khan *et al.* 2009 from Pakistan; 27.26% and 25.2% by Kabir *et al.* 2010 and Afroze *et al.* 2013 from different provinces of Bangladesh; 25.9% by Mungube *et al.* 2012 from Kenya; 26.55% by Nega *et al.* 2012 from Ethiopia; 23.96% reported by Asressa *et al.* 2012 from Andassa Livestock Research Center in North-West of Ethiopia.

The results revealed that the lowest prevalence of Fasciolosis for Year 2014 was in the month of May (14.2%) and highest being in the month of August (35.8%). However in Year 2015, the prevalence rate was highest in the month of September (44.66%) followed by October (39.66%) and lowest in May (9.3%). Moreover, the infection was reported throughout the year due to resistance of metacercariae for desiccation, especially during the dry season and continued presence of the shallow water, enough vegetation and humidity for continued exposure of the animals to encysted metacercariae and no restriction on cattle grazing habits and movement between the infected and treated localities which was also suggested by El Bahy, 1998.

These results are in agreement with Pfukenyi *et al.* 2006 and Faria *et al.* 2005 who reported high intensity in August-September in Zambian cattle and in dairy cattle herd in Minas Gerais, Brazil respectively. Similarly, Qureshi *et al.* 2012 recorded lowest prevalence in the month of May in Buffaloes of Northwestern Punjab, Pakistan which supports the findings of the current study. In both years, the lowest infection in May-June can be related to progression of hot dry weather, as the temperature was high and humidity was low in these months.

Table 1: Yearwise Prevalence of Fasciolosis

YEAR	EX.	INF.	PREV	$\chi^2$ (P-Value)
2014	316	80	25.31%	0.183 0.669
2015	396	107	27.02%	
Total	714	192	26.84%	

Month-wise prevalence (Fig. 1)




Figure 1: Monthwise prevalence of Fasciolosis (2014-2015)

Season wise Prevalence (Table and Fig 2)

On seasonal basis, the current study showed maximum spread of disease in Autumn Season i.e. 33.33% and 40% in Year 2014 and 2015 respectively. The minimum infection was recorded in spring season showing prevalence of 20% and 12.9% in consecutive studied years. There was no statistically significant difference between seasons in year 2014 which has already been stated could be attributed to skipping the data of two months due to natural disaster Kashmir valley faced. However statistically significant difference was observed between seasons in year 2015.

This difference could be due to a variety of weather condition in each year. The highest prevalence in autumn was also reported by Chaudhri *et al.* 1993; Maqbool *et al.* 1994 and Ghirmire and Karki 1996; Abrous *et al.* 1999; Maqbool *et al.* 2002; Pfukenyi *et al.* 2005 and Haridy *et al.* 2006 who emphasized that the possible reason for the same could be availability of favourable temperature and moisture for the rapid propagation of the parasitic trematode life cycle in this very season.

Table 2: Season wise prevalence of Fasciolosis (2014-2015)

Year	2014			2015		
Season	Ex.	Inf.	Prev.	Ex.	Inf.	Prev.
Spring	115	23	20%	82	10	12.9%
Summer	99	26	26.26%	102	20	19.6%
Autumn	12	4	33.33%	146	59	40%
Winter	90	27	30%	66	18	27.27%
$\chi^2$ (p-Value)	3.218(0.486)			25.26(0.000)		




Figure 2: Cumulative Season wise prevalence

Distribution on the basis of infection type (Table 3)

Of the total 192 affected livers by fasciolosis, 149 (77.60%), 24 (12.5%) and 19 (9.89%) respectively showed *Fasciola gigantica*, *Fasciola hepatica* and mixed infection (*Fasciola hepatica* and *Fasciola gigantica*).

The finding of this study was in consistence with the earlier investigation by Ashrafi *et al.* 2004 from Gilan province; Mir *et al.* 2008 from Kashmir and Khan *et al.* 2009 from Punjab Province (Pakistan) and by Phiri *et al.*

2005; Abunna *et al.* 2009; Fufa *et al.* 2009; Mwabonimana, *et al.* 2009. The predominance of *Fasciola gigantica* could be due to the availability of appropriate environmental conditions and topography (lowland and middle altitude zone) which are favorable habitat to its intermediate host *L. natalensis* (Urquhart *et al.* 1996). However, inverse distribution was reported by Melugeta *et al.* 2011; Belay *et al.* 2012; Chakiso, *et al.* 2014 and Alemu and Abebe 2015. Mixed infection of *F.*

*hepatica* and *F. gigantica* occurs presumably as a result of the movement of stock between high and low ground or through overlapping of the territories of the snail

vector at altitudinal range of 1200-1800 M.a.s.l. (Kendel 1954 and Graber, 1975).

Table 3: Prevalence based on type of infection

Infection Type	Infected	Prev. Among Infected Ones (N=192)	Overall Prevalence (N=714)
<i>F. gigantica</i>	149	77.60%	20.86%
<i>F. hepatica</i>	24	12.5%	3.361%
Mixed	19	9.89%	2.66%
$\chi^2$		254.29(p=0.000)	186.22(p=0.000)

#### Age-wise distribution (Table 4)

Out of 714 cattles, 166 heads were of age group <1-3Years, 396 of age between 3-6 years and 152 having age >6 Years. Among these 3 age categories, prevalence of Fasciolain livers was highest in >3-6 years age group (30.30%) followed by age group >6 years (28.28%) and least infection in bovines of age 1-3Years(17.46%).The results in current study were in consistency with Keyu *et al.* 2005;Rehman *et al.*2015.The sound explanation behind the lower prevalence in age group >6 yrs compared to younger age group(3-6yrs) could be due to self-cure phenomenon (Assanji, 1988) or high acquired immunity

which increase with age (Dwinger *et al.* 1982). It has been also reported that *Fasciola* infected hosts may recover from parasitic infection with increasing age and hence become resistant (Yilma and Mesfin, 2000; Shiferaw *et al.*2011; Mufti, 2011.Mulcahy *et al.* 1999 suggested that resistance is not wholly immunological based rather resistance to reinfection may be due to hepatic fibrosis resulting from primary infection. Least infection in age group <1-3 years is possibly due to less chances of acquiring infection due to short exposure time as compared to older animals which is in agreement with (Anderson *et al.* 1999)and Teklu *et al.* 2015.

Table 4: Agewise prevalence of fasciolosis

Age	Ex.	Inf.	Prevalence	$\chi^2$ p-Value
1Yr-3Yrs	166	29	17.46%	9.991 0.007
3Yrs-6Yrs	396	120	30.30%	
>6Yrs	152	43	28.28%	

#### Genderwise prevalence (Table 5)

Out of 531 males and 183 females slaughtered during the survey period, males won by retaining lesser infection of 19.96% and were par to females who showed higher prevalence of 46.99%. The difference was highly significant and thus revealed sex as determinant influencing the prevalence of Fasciolosis rate. Our findings are in agreement with results of Daniel 1995; Molina *et al.* 2005; Bhutto *et al.* 2012 and Teklu *et al.* 2015 who reported higher prevalence of this parasite in female than male.

In the current studied abattoirs, the number of slaughtered male cattles (531) was far higher than the females (183). The number of positive females was higher in proportion than males even if the number of female cattle that come to abattoir were fewer in number. These results were in consistent to Kara *et al.* 2009. High infection rate in females can be multifactorial like high stress during parturition period (Spithill *et al.* 1999), weak and malnourished making them more susceptible to infection (Blood and Radostits, 2000) or due to the feeding conditions i.e females are generally being let loose to graze freely in pastures. The other

possible reason for the same could be that the most of people traditionally feed their lactating cows with grasses during dry season which are grown around rivers and marshy areas for the sake of getting high milk yield as suggested by Gracy *et al.* 1999 and Tilahun *et al.* 2014. However, some authors revealed that male cattle are more prone to Fasciolosis than female counterparts like Khan and Maqbool 2012. But, Rahmeto 1992 and Dagne 1994; Keyyu *et al.* 2005; Phiri *et al.* 2005; Khan *et al.* 2009; Kabir *et al.* 2010; Kanyar *et al.* 2010; Assefa *et al.* 2015 reported no significant difference between the gender of animal and infection rate which could be associated with similar management given to both group of animals or probably due to common grazing pastures on which both are fed together, which expose them to the same risk of infection.

Table 5: Genderwise prevalence of Fasciolosis

	Examined	Infected	Prevalence
Males	531	106	19.96%
Females	183	86	46.99%
$\chi^2$ (p-value)	49.221(0.000)		

#### Association of body condition with infection (Table 6)

Among all examined animals (n = 714), 30.53% (n = 218) were marked as poor (body score 1-3), 35.05% (n =250) as Medium (4-6) and 34.44% (n = 246) as Good (7-9) body conditions. 42.66% of infection ( $n_i=93$ ) was recorded in animals with poor body condition, 22.40% of infection ( $n_i=56$ ) in animals with medium body score and 17.47% of infection ( $n_i=43$ ) in animals having good body condition. Thus, an inverse association was found between the body condition and infectious rate of Fasciolosis which was statistically significant ( $p<0.05$ ). These findings are in accordance to Mihreteab *et al.* 2010; Tilahun *et al.* 2014 and Teklu *et al.* 2015.

The current findings that lean animals are associated with higher *Fasciola* infection compared to animals with medium and normal body condition could be attributed to emaciation due to lack of nutrients, loss of blood and tissue fluid and damage to liver parenchyma caused by presence of flukes. Similar findings were reported by Beckele *et al.* 2010; Tesfay *et al.* 2012 and Assefa *et al.* 2015. Radostits *et al.* 1994 has mentioned that Chronic fasciolosis is the commonest disease in cattle and significantly characterized with weight loss.

Table 6: Effect of body condition on prevalence of Fasciolosis

Body Condition	Ex.	Inf.	Prevalence	$\chi^2$ p-Value
Poor	218	93	42.66%	41.223 0.000
Medium	250	56	22.40%	
Good	246	43	17.47%	

#### Breedwise prevalence of Fasciolosis (Table7)

Out of the total 71 cattle examined, 213 were reared locally and 501 were imported from other states to the valley for slaughter purpose. The prevalence of fasciolosis was 40.80% and 20.90% for local and nonlocal breed cattle, respectively. There was statistically significant ( $\chi^2 = 29.06$ ,  $P = 0.000$ )

association of fasciolosis with breeds. Our results are in agreement with study conducted by Teklu *et al.* 2015.

This difference in prevalence based on breed might be due to the management of the animals as most of the local animals were reared in the extensive system of management which makes them easily susceptible to the parasites

Table 7: Breed wise Prevalence of Fasciolosis

Breed	Ex.	Inf.	Prevalence	$\chi^2$ p-Value
Locals	213	87	40.80%	29.06 0.000
Non-locals	501	105	20.90%	

## V. CONCLUSION

Moderate intensity of fasciolosis was recorded in the current study. In line with above findings, it is recommended that strategic application of flukicides should be done and further epidemiological studies on biology and ecology of intermediate host should be carried out so to develop substantiable planning for considerable success in control of Fasciolosis. There is need to carry out economic analysis so as to give appropriate economic losses directly by liver condemnation.

## VI. ACKNOWLEDGEMENT

I extend my heartily thanks to Dr Javed, Assitant Professor, Dept of Statistic, University of Kashmir for proficiency and support regarding Biostatistics computation and analysis. I would like to thank butchers of abattoirs visited during current study for their interest to participate in research work by providing liver samples of cattle which form the basic tool for this piece of research work.

## BIBLIOGRAPHY

1. Abrous, M., Rondelaud, D.; Dreyfuss, G.; Cabaret, J. (1999): *Paramphistomum daubneyi* and *F. hepatica*: Influence of temperature changes on the shedding of cercariae from dually infected *Lymnaea truncatula*. *Parasitol. Res.* **85**: 765-769.
2. Abunna, F.; Asfaw, L.; Megers, B and A. Regassa, A. (2009). Bovine Fasciolosis: Coprological, Abattoir Survey and its Economic Impact due to Liver Condemnation at Soddo Municipal Abattoir, Southern Ethiopia. *Tropical Animal Health and Production*, **42**: 289-292.
3. Affroze, S.; Begum, N.; Islam, M.S.; Rony; S.A.; Islam, M.A. and Mondal, M.M.H. (2013). Risk Factors and Gross Pathology of Bovine Liver Fluke Infection at Netrokona District, Bangladesh. *J. Anim. Sci. Adv.* **3**(2): 83-89.
4. Akhoun, Z. A. and Peer, F.U. (2014). Prevalence of fascioliasis in district Srinagar of Kashmir valley. *Vet. Sci. Res. J.* **5**(1&2): 54-57.
5. Alemu, A. and Abebe, B. (2015). Fasciolosis: Prevalence, Evaluation of Floatation and Simple Sedimentation Diagnostic Techniques and Monetary Loss due to Liver Condemnation in Cattle Slaughtered at Wolaita Soddo Municipal Abattoir, Southern Ethiopia. *Food Science and Quality Management*. **43**.
6. Anderson, N., Luong, T.T., Vo, N.G., Bui, K.L., Smooker, P.M., Spithill, T.W. (1999). The sensitivity and specificity of two methods for detecting *Fasciola* infection in cattle. *Vet. Parasitol.* **83**: 15-24.
7. Ashrafi, K.; J Massoud 1, K Holakouei 2, M Mahmoodi 2, MA Joafshani 3, MA Valero 4, MV Fuentes 4, M Khoubbane 4, P Artigas 4, MD Bargues 4, S Mas-Coma (2004). Evidence Suggesting that *Fasciola gigantica* Might be the Most Prevalent Causal Agent of Fascioliasis in Northern Iran. *Iranian J Publ Health*, **33**(4) 31-37.
8. Asressa Yeneneh, Hassen Kebede, Tewodros Fentahun, Mersha Chanie, 2012. Prevalence of cattle fluke's infection at Andassa Livestock Research Center in north-west of Ethiopia. *Veterinary Research Forum*, **3** (2): 85-89.
9. Assanji, M. F. (1988). Helminth infection in livestock. *J. Helminthol.*, **62**: 243-249.
10. Assefa, A.; Assefa, Z.; Beyene, D. and Desissa, F. (2015). Prevalence of bovine Fasciolosis in and around Inchini town, West Showa Zone, Ada'a Bega Woreda, Central Ethiopia. *J. Vet. Med. Anim. Health.* **7**(6): 241-248.
11. Bekele, M., Haftom, T & Yehenew, G. (2010). Bovine fasciolosis: Prevalence and its economic loss due to liver condemnation at Adwa municipal abattoir, North Ethiopia. *Ethiopian Journal of Agricultural Science and Technology*. **1**: 39-47.
12. Belay, E.; Molla, W. and Amare, A. (2012). Prevalence and Economic Losses of Bovine Fasciolosis in Dessie Municipal Abattoir, South Wollo Zone, Ethiopia. *European Journal of Biological Sciences.* **4** (2): 53-59.
13. Bhutto, B.; Arijo, A.; Phullan, M.S. and Rind, A. (2012). Prevalence of fascioliasis in buffaloes under different agro-climatic areas of Sindh Province of Pakistan. *International Journal of Agricultural Biology.* **14**: 241-245.
14. Braun, U., Wolfensberger, R., Hertzberg, H., 1995. Diagnosis of liver flukes in cows- a coparison of the findings in the liver, in the faeces and in the bile. *Schweizer Archiv for Tierheilkunde* **137** (5), 187-92.
15. Chakiso, B.; Menkir, S. and Desta, M.(2014). On Farm Study of Bovine Fasciolosis in Lemo District and its economic loss due to liver condemnation at Hossana Municipal abattoir, Southern Ethiopia. *Int.J. Curr.Microbiol.App.Sci* **3**(4): 1122-1132.
16. Chaudhri, S. S.; Gupta, R. P.; Kumar, S.; Singh, J. and Sangwan, A. K. (1993): Epidemiology and control of *Fasciola gigantica* infection of cattle and buffaloes in Eastern Haryana, India. *Indian. J. Anin. Sci.* **63**: 600-605.
17. Dagne, M. (1994). Survey on Prevalence and Economic Significance of Bovine Fasciolosis in Debre Berhan region. DVM Thesis, Faculty of Veterinary Medicine, Addis Ababa University, Debre Zeit, Ethiopia.
18. Daniel, F. (1995). Economic importance of organs condemnation due to fasciolosis and hydatidosis in cattle and sheep slaughtered at Dire Dawa Abattoir, DVM thesis, Faculty of Veterinary Medicine, Addis Ababa University Debre zeit, Ethiopia.
19. Dar, G. H.; Bhagat, R. C. and Khan, M. A. (2002). Biodiversity of the Kashmir Himalaya. (Valley Book House, Srinagar - 190006, J&K), pp. 19.
20. Dwinger, R.H.; Leriche, P.D. and Kuhne, G.I. (1982). Fasciolosis in beef cattle in North West Argentina. *Trop. Anim. Health Prod.* **14**: 167-171.
21. El-Bahy, N. M. (1998). Strategic Control of Fascioliasis in Egypt. Review article. Continual Scientific Committee of Pathology, Microbiology and Parasitology, Egypt.
22. Faria, R. N.; Cury, M.C. and Lima, W.S. (2005). Prevalence and dynamics of natural infection with *Fasciola hepatica* (Linnaeus, 1758) in Brazilian cattles. *Revue Méd. Vét.* **156**(2): 85-86.
23. Fatima, M.; Chishti, M. Z.; Ahmad, F. and Lone, B. A.(2012). Epidemiological Study of Fasciolosis in Cattle of Kashmir Valley. *Advances in Biological Research* **6** (3): 106-109.
24. Fufa, A.; Loma, A.; Bekele, M. and Alemayehu, R. (2009). Bovine fasciolosis: Carpoological, abattoir survey and its economic impact due to liver condemnation at Soddo municipal abattoir,

- Southern Ethiopia. *Tropical Animal Health Production*. **42**: 289-292.
25. Ghirmire, N. P. and Karki, N. P. S. (1996): Prevalence of fascioliasis and efficacy of various anthelmintics in buffaloes of Rural Kathmandu vetcon. *N. U. A.* pp. 43.
  26. Graber, M.M. (1975). Helminths and helminthiasis of domestic and wild animals of Ethiopia. *Bulletin Animal Health Production*. **23**: 57- 86.
  27. Gracy, J.; David, S.; Collins and Robert Huey (1999) Meat Hygiene, 10 edition, Harcourt Brace pp: 681-686.
  28. Kabir, M. H. B.; Eliyas, M.; Hashem, M. A.; Mohiuddin and Miaz, O. F. (2010). Prevalence of zoonotic parasitic diseases of domestic animals in different abattoir of Comilla and Brahman Baria region in Bangladesh. *Univ. J. Zool. Rajshahu Univ.* **28**: 21-25 <http://journals.sfu.ca/bd/index.php/UJZRU>
  29. Kendall, S.B. (1954). Fascioliasis in Pakistan. *Ann. Trop. Med. Parasitol.*, **48**(3): 307-313.
  30. Keyyu J D, Monrad J, Kyvssgard N C and Kassuku A A (2005). Epidemiology of *Fasciola gigantica* and paramphistomes in cattle on traditional, smallscale dairy and largescale dairy farms in the southern highlands of Tanzania. *Tropical Animal Health and Production* **37**: 303314.
  31. Khan, M. K.; Sajid, M. S.; Khan, M. N.; Iqbal, Z. and Iqbal, M. U. (2009) Bovine fasciolosis: Prevalence, effects of treatment on productivity and cost benefit analysis in five districts of Punjab, Pakistan. *Research in Veterinary Science* **87**: 70–75.
  32. Kuchai, J.A.; Chishti, M.Z.; Zaki, M.M.; Dar, S.A.; Muzaffar Rasool; Ahmad, J. and Hidayatullah Tak (2011). Some Epidemiological Aspects of Fascioliasis among Cattle of Ladakh. *Global Veterinaria* **7** (4): 342-346.
  33. Kumar, V.; Banerjee, P. S.; Hira Ram; Rajat Garg and Yadov, C. L. (2002). Faecal examination vis-à-vis bile examination for the diagnosis of bubaline fasciolosis. *Journal of Veterinary Parasitology* **16**: 185-186.
  34. Maqbool, A.; Arshad, M. J.; Mahmood, F. and Hussain, A. (1994): Epidemiology and chemotherapy of fascioliasis in buffaloes. *Assiut. Vet. Med. J.* **30**, 115-123.
  35. Maqbool, A.; Hayat, C. S.; Akhtar, T. and Hashmi, H.A. (2002). Epidemiology of fasciolosis in buffaloes under different managemental conditions. *Vet. arhiv* **72**, 221-228,
  36. Mas-Coma, S.; Bargues, M.P. and Valero, M.A. (2005). Fascioliasis and other plant-borne trematode zoonoses. *International Journal for Parasitology*, **35**: 1255– 1278.
  37. Mihreteab, B., Haftom, T. and Yehenew, G. 2010. Bovine Fasciolosis: Prevalence and its Economic loss due to liver condemnation at Adwa Municipal Abattoir, North Ethiopia. School of Veterinary Medicine, College of Agriculture and Veterinary Medicine, Jimma. *Ethiopian Journal of Applied sciences and Technology*, **1**(1): 39-47.
  38. Mir, R.A.; Chishti, M.Z.; Zarger, M.A.; Tak, H. and Dar, F.A. (2008). Seasonal prevalence of trematode parasites of sheep (*Ovis aries* L.) in Kashmir Valley, India. *Nigerian Journal of Parasitology* **29**(2): 80-83.
  39. Molina, E. C.; Gonzaga, E.A. and Lumbae, L.A. (2005). Prevalence of infection with *fasciola* gigantic and its relationship to carcass and liver weights and fluke and egg counts in slaughter cattle and buffaloes in Southern Mindanao, Philippines. *Tropical Animal Health Production*. **37**: 215-221.
  40. Mufti, S., 2011. An epidemiological study of bovine fasciolosis in potohar region. Faculty of Sciences, Arid Agriculture University, Rawalpindi, Pakistan.
  41. Mulcahy, G., P. Joyce and J.P. Dalton. Immunology of *Fasciola hepatica* infection In: Dalton, JP (Ed), *Fasciolosis* CAB International, pp 341-376 (1999).
  42. Mulugeta, S.; Begna, F. and Tsegaye, E. (2011). Prevalence of Bovine Fasciolosis and its Economic Significance in and Around Assela, Ethiopia. *Global Journal of Medical Research*. **11**(3).
  43. Mungube, E.O.; Sila, D.M.; Kariuki, C.W.; Bauni, S.M.; Tenhagen, B.A.; Wamae, L.; Nginyi, J. and Omondi, G.A. (2012). A cross-sectional survey on fasciolosis in selected settlements of Taveta Division, Coast Province, Kenya. *Livestock Research for Rural Development* **24** (4): 1-8.
  44. Mwabonimana, M.F., A.A. Kassuku, H.A. Ngowi, L.S.B. Mellau, H.E. Nonga and E.D. Karimuribo, (2009). Prevalence and Economic Significance of Bovine Fasciolosis in Slaughtered Cattle at Arusha Abattoir Tanzania. *Tanzania Veterinary J.*, **26**: 68-74.
  45. Nega, M.; Bogale, B.; Chanie, M.; Melaku, A. and Fentahun, T. (2012). Comparison of Coprological and Postmortem Examinations Techniques for the Determination of Prevalence and Economic Significance of Bovine Fasciolosis. *Journal of Advanced Veterinary Research*. **2**: 18-23.
  46. Ogunrinade, A. and G. O. Adegoke, 1982. Bovine fascioliasis in Nigeria. Intercurrent parasitic and bacterial infection. *Tropical Animal Health Production*, **14**: 121-125.
  47. Pfukenyi, D. M., J. Monrad, S. Mukaratirwa (2005a): Epidemiology and control of trematode infections in cattle in Zimbabwe: a review. *J. S. Afr. Vet. Assoc* **76**, 9-17.
  48. Phiri, A.M., I.K. Phiri, C.S. Sikasunge and J. Monrad, (2005). Prevalence of Fasciolosis in Zambian Cattle Observed at Selected Abattoirs with Emphasis on Age, Sex and Origin. *J. Veterinary Medicine B*, **52**: 414-416.
  49. Qureshi AW, Tanveer A, Maqbool A & Niaz S (2012). Seasonal and monthly prevalence pattern of fascioliasis in buffaloes and its relation to some climatic factors in north eastern areas of Punjab,

- Pakistan. Iranian Journal of Veterinary Research, 13(2): 134-137.
50. Radostitis, O.M., Blood, D.C., Gay, C.C. (1994). A text book of the disease of cattle, sheep, goats, pigs and horse: Veterinary medicine. 8th ed. London, UK: Bailliere, Tindall. 105-106.
  51. Rahmeto A (1992). Fasciolosis Clinical occurrences, Coprological, Abattoir and Snail survey in and around Wolisso. DVM thesis, Faculty of Veterinary Medicine, Addis Ababa University, Debre Zeit, Ethiopia.
  52. Rehman, T. U.; Khan, M. N.; Sajid, M. S. and Javed, M.T. (2013). Slaughter house based epidemiology and estimation of economic losses of bovine Fascioliasis in Tehsil Sargodha. Pakistan Journal of Science 65 (4):
  53. Schweizer, G.; Braun, U.; Deplazes, P. and Torgerson, P.R. (2005). Estimating the financial losses due to bovine fasciolosis in Switzerland. Veterinary Record .157:188-193.
  54. Sheikh, G.N.; Qadri, G.J. and Gunjan Das (2007). Seasonal prevalence of bovine Fascioliasis in Kashmir valley. Indian Journal of Veterinary Medicine 27(2): 119-120.
  55. Shiferaw Mulugeta, Feyisa Begna, Ephrem Tsegaye, 2011. Prevalence of Bovine Fasciolosis and its economic significane in and around Assela, Ethiopia. Global Journals Inc. 11: 1-2.
  56. Soulsby E.J.L (1986) Helminths, Arthropods and Protozoa of Domesticated Animals, 7th. Ed edn. Bailliere Tindall, London, pp 212–342.
  57. Spithill, T. W., P. M. Smooker and D. B. Copeman, 1999. Fasciola gigantica: epidemiology, control, immunology and molecular biology. Wallingford, Oxon, UK: CABI. Pp. 465–525.
  58. Teklu, H.; Abebe, N and Kumar, N. (2015). Abattoir prevalence of bovine fasciolosis in the manucipal abattoir of Wukro, Northern Ethiopia. Journal of International Academic Research for Multidisciplinary 2(12): 430-438.
  59. Tesfay, H.; Dejene, T. and Kebede, E. (2012). Prevalence of bovine fasciolosis and its associated risk factors in Mekelle municipal abattoir. *Journal of the Drylands*, 10-17.
  60. Thrustfield, M. (2005). Veterinary Epidemiology. 3<sup>rd</sup> Edition, Black Well Science Ltd. Cambridge, USA pp: 225-228.
  61. Tilahun, Z.; A. Nemomsa, D. Haimanot and K. Girma (2014) Study on Prevalence of Bovine Fasciolosis at Nekemte Veterinary Clinic, East Wollega Zone, Oromia, Ethiopia. European Journal of Biological Sciences 6 (2): 40-45.
  62. Urquhart, G. M.; Armour, J.; Duncan, J.L.; Dunn, A.M. and Jennings, F. W. (1996). Veterinary Parasitology. 2<sup>nd</sup> edition. Blackwell Science Ltd., Osney Mead, London, UK.
  63. Vercruysse, J. and Claerebout, E. (2001). Treatment vs non treatment of helminth infection in cattle defining the threshold. Veterinary Parasitology. 98: 195-214.
  64. Yilma Jabore and Mesfin Ali, 2000. Dry season bovine fasciolosis in northwestern part of Ethiopia. Revue Medical Veterinary, 151(6): 493-500.





GLOBAL JOURNAL OF MEDICAL RESEARCH: G  
VETERINARY SCIENCE AND VETERINARY MEDICINE  
Volume 16 Issue 3 Version 1.0 Year 2016  
Type: Double Blind Peer Reviewed International Research Journal  
Publisher: Global Journals Inc. (USA)  
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

## Sero-Prevalence of *Toxoplasma Gondii* Infection and Associated Risk Factors in Animals Presented to Sholla and Akaki-Kality Veterinary Clinics, Addis Ababa

By Abebaw Getachew, Alebachew Tilahun, Alemu Aylate & Wale Tesfaye  
*Wolaita Sodo University*

**Abstract-** A cross-sectional study on *Toxoplasma Gondii* in livestock was carried from October 2011 to March 2012 in Addis Ababa, Ethiopia to determine sero-prevalence and associated risk factors. A total 347 serum samples were collected from the jugular veins of each animal and heart of swine, presented veterinary clinics and abattoirs, respectively. The overall prevalence in the six animal species out of 347 animals sampled was 126 (36.1%) which were detected as sero-positive for toxoplasmosis. Prevalence based on animal species, out of 347 different animal species sampled were 71 (36.6%), 15 (37.5%), 11 (25%), 19 (47.5%), 7 (35%) and 3 (33.3%) in ovine, caprine, bovine, swine, equine and camel, respectively. In attempt look for the association between risk factors and sero-prevalence, a questionnaire survey was conducted and the result obtained showed economic loss due to abortion (30%), stillbirth (12%), neonatal mortality (18%), dystocia (17%), retained fetal membrane (8%) and endometritis (6%). Seventy two (72) of the respondents had cats in their household.

**Keywords:** *toxoplasmosis, veterinary clinics, seroprevalence, risk factors.*

**GJMR-G Classification:** NLMC Code: QX 140



*Strictly as per the compliance and regulations of:*



© 2016. Abebaw Getachew, Alebachew Tilahun, Alemu Aylate & Wale Tesfaye. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License <http://creativecommons.org/licenses/by-nc/3.0/>), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

# Sero-Prevalence of *Toxoplasma Gondii* Infection and Associated Risk Factors in Animals Presented to Sholla and Akaki-Kality Veterinary Clinics, Addis Ababa

Abebaw Getachew <sup>α</sup>, Alebachew Tilahun <sup>σ</sup>, Alemu Aylate <sup>ρ</sup> & Wale Tesfaye <sup>ω</sup>

**Abstract-** A cross-sectional study on *Toxoplasma Gondii* in livestock was carried from October 2011 to March 2012 in Addis Ababa, Ethiopia to determine sero-prevalence and associated risk factors. A total 347 serum samples were collected from the jugular veins of each animal and heart of swine, presented veterinary clinics and abattoirs, respectively. The overall prevalence in the six animal species out of 347 animals sampled was 126 (36.1%) which were detected as sero-positive for toxoplasmosis. Prevalence based on animal species, out of 347 different animal species sampled were 71 (36.6%), 15 (37.5%), 11 (25%), 19 (47.5%), 7 (35%) and 3 (33.3%) in ovine, caprine, bovine, swine, equine and camel, respectively. In attempt look for the association between risk factors and sero-prevalence, a questionnaire survey was conducted and the result obtained showed economic loss due to abortion (30%), stillbirth (12%), neonatal mortality (18%), dystocia (17%), retained fetal membrane (8%) and endometritis (6%). Seventy two (72) of the respondents had cats in their household. The sero-prevalence on the basis of sex was significantly higher ( $p < 0.05$ ) in males than females. No statistically significant association was noted among other factors and sero-positivity. Results of present study suggest widespread environmental contamination with *T. gondii* oocysts and that livestock could be a potentially important source of *T. gondii* infection if their infected meat is consumed undercooked.

**Keywords:** toxoplasmosis, veterinary clinics, sero-prevalence, risk factors.

## I. INTRODUCTION

*Toxoplasma gondii* infections are prevalent in humans and animals worldwide (Dubey and Beattie, 1988). Felids are the key animal species in the life cycle of this parasite because they are the hosts that can excrete the environmentally-resistant stage, the oocyst. Humans become infected postnatal by ingesting tissue cysts from undercooked meat, consuming food or drink contaminated with oocysts, or by accidentally ingesting oocysts from the environment. However, only a small percentage of exposed adult humans or other animals develop clinical signs of disease. It is unknown

**Author α:** University of Gondar, College of Veterinary Medicine and Animal Science, Biomedical Science Unit, Gondar, Ethiopia, P.Box-196. e-mail: abebaw80@gmail.com

**Author σ ρ ω:** Wolaita Sodo University, School of Veterinary Medicine, Ethiopia. e-mails: alebmen19@gmail.com, wsusvm2014@gmail.com, twalelgn@yahoo.com

whether the severity of toxoplasmosis in immune-competent hosts is due to the parasite strain, host variability or other factors. Recently, attention has been focused on genetic variability among *T. gondii* isolates from apparently healthy and sick hosts. It has been 100 years since the discovery and naming of *T. gondii*. The parasite was first found in laboratory animals (Dubey, 2007). Its medical importance remained unknown until 1939 when *T. gondii* was identified conclusively in tissues of a congenitally-infected infant in New York City, USA (Wolf *et al.*, 1939), and its veterinary importance became known when it was found to cause abortion storms in sheep in 1957 in Australia (Hartley and Marshall, 1957).

Although infection does not clinically affect cattle, transmission of infection to humans from tissue cysts when eating raw or undercooked beef should not be discounted. Toxoplasmosis may be important in Ethiopia where raw or partially cooked meat is regarded as a delicacy (Bekele and Kasali, 1989). In Ethiopia, there are documented reports on serological survey of Caprine toxoplasmosis by Teshale and his colleagues in Central and Southern Ethiopia (Teshale *et al.*, 2006). The serological survey results on toxoplasmosis by Negash and his associates further confirm the presence of *T. gondii* infection in sheep and goat population in Ethiopia (Negash *et al.*, 2004). The results of a questionnaire survey in Debre Birhan and the surrounding area revealed that abortion was the major cause of lamb loss during 12 months studied period (Getachew and Tilaye, 2002). In addition, the Sero-prevalence, assessment of its zoonotic importance and identification of factors associated with Sero-prevalence was documented in Nazareth town, Ethiopia (Negash *et al.*, 2008).

Toxoplasmosis is recognized as disease of great economic importance since it causes heavy losses through abortion, stillbirth, neonatal mortality, encephalitis and pneumonia particularly in sheep and goats (Radostits *et al.*, 2007 and Singh and Msolla, 1994). If animals are important in the epidemiology of human toxoplasmosis it is well to have information concerning serological study in those hosts (Morris *et al.*, 2007). In the present paper, we summarize

information on serological prevalence of *T. gondii* infection in different animals species presented Veterinary Clinics and economic impact of the disease in the study area.

## II. MATERIALS AND METHODS

### a) Study Area

The study was conducted in Addis Ababa which lies at an altitude of 2000-3000 meters above sea level. The mean annual rainfall is 1800 mm with a bimodal pattern. There are dry and rainy seasons in the area. The long rainy season extends from June to September, contributes about 84% of the total annual rainfall while the dry season lasts from October to February. The short rainy season lasts from March to May. The study was conducted from November 2011-March 2012 in Sholla and Akaki-Kality Veterinary Clinics, Addis Ababa. The mean annual minimum and maximum temperature are 14°C and 21°C respectively with an average rate of 17°C the mean relative humidity is 61.3% (CSA, 2009).

### b) Study Population

The study included all animals which came to Sholla and Akaki-Kaliti Veterinary Clinics and consisted of bovine, ovine, caprine, swine, camel and equine species regardless of their age, breed and disease case. Blood sample for swine and camel was taken from Addis Ababa Abattoir Enterprise.

### c) Study Design

A cross-sectional study was conducted from October 2011 to March 2012 to determine the prevalence toxoplasmosis among animals coming to Sholla and Akaki-Kaliti Veterinary Clinics for various health problems. After reviewing daily patient flow (case) to the clinic, the expected patient population in the study period was taken as a sampling frame. A simple random sampling method was used to collect blood sample from different animal species.

### d) Sample Size Determination

The required sample size for the study animals was determined by the formula given by (Thrusfield, 1995) assuming 95% of confidence interval and at 5% desired precision. This was estimated with the assumed toxoplasmosis prevalence of 34.5% based on previous study by (Getachew and Tilaye, 2002) by taking the average prevalence of 34% (sheep) and 35% (goat) in Debre Birhan and the surrounding areas. Accordingly the desired sample size was 347.

### e) Study Methodology

#### i. Serum collection and serological examination

Approximately 5ml of blood was taken from jugular vein of each study animal but for swine it was taken directly from heart in abattoir and the serum was separated and stored at deep freezer until tested. *Toxoplasma gondii* antibody was determined by the

Slide Agglutination Test using a commercial kit (HUMA TEX TOXO, Human Gesellsfür Biochemica und Diagnostoica mbH Max-Plank-ring21.65205 Wiesbaden. Germany). This method is quick, simple and requires smaller quantity of reagents. Agglutination reactions are more sensitive than immuno-precipitation tests. The tests are simple and have an added advantage of easily readability (Chauhan and Agarawal, 2006).

Comparable assessment of slide agglutination test shows that as sensitive as, and a more specific than latex agglutination test. The predictive value of a negative slide agglutination test is less than the latex agglutination test but produced results within minutes, although, quantitative results is not comparable to other assays. Slide agglutination presents a rapid alternative to the latex agglutination test as a screening assay toxoplasmosis, although patients at risk of life threatening infection require detailed serological examination using additional methods (Dunford and Johnson, 1991). In this test a clean dry glass with 6 cells was taken and a drop of antigen suspension was placed over the middle of area of each 6 cell and one drop of positive control serum (goat) in one of cell while the negative control serum was placed on the other cell and one drop of test serum was placed on the rest 4 cells. Then mixed with separate disposable sticks and spreader the fluid over the entire area of the particular cell the slide was tilted back and forth of 4 minutes so that rotates slowly inside the cell. Finally it was observed for clumping (agglutination) by naked eye and magnifying lenses in comparison with the two controls (positive and negative). The negative result was identified as negative control result which did not form agglutination (homogenous appearance) but distinct agglutination was indicator for positive toxo-Ab of at least 4 IU/ml similar with positive control.

#### ii. Questionnaire Survey

A pre-tested structural questionnaire was prepared to animal owners with respect to the case they brought which included both open ended and closed ended questions. The questions was concerned with hygiene, environment, management, nutrition, reproductive disorder history and nervous signs, ownership of cats, purpose of cat keeping as well as the mechanism of cat feces disposal. In addition, the habit of exposure of raw meat and milk were important questions that gave useful information for epidemiology of human toxoplasmosis.

### f) Data Analysis

All data obtained from the study were entered into Ms Excel 2007 data sheet and analyzed using STATA 11, statistical software programme. The Seroprevalence was calculated later on by dividing the sera were found positive to slide agglutination test to the total sample size multiplied by 100. The risk factors associated with toxoplasmosis were determined using

percent values and using Pearson's Chi-square ( $\chi^2$ ). A statistically significant association between variables was said to exist if the calculated level of significance is less than 5% ( $p < 0.05$ ) at 95% confidence level. The strength of associations between the exposure to the risk factors and sero-positivity is measured using odds ratio (Wasserthiel-smaller, 1995).

### III. RESULTS

#### a) Sero-prevalence

The overall Sero-prevalence rate of the test result were found in 126 of the 347(36.31%) animals (table 1) (6 different species of animals) examined for slid agglutination test (CI=31.22, 41.40, 95% level of confidence). Sero-prevalence by origin, age, species, management system, hygiene, reproductive abnormality, cat ownership as well as associated clinical findings is not significant while a statistically significant difference( $p < 0.05$ ). Sero-prevalence among males than females being observed. Higher prevalence was observed in males and females (table 5).

#### b) Risk Factors Associated with Sero-positivity

Factors closely related to the natural history of toxoplasmosis are presence of cats, origin, history of abortion or neonatal mortality or births of weak lambs and reproductive abnormalities, management practices, hygiene and clinical finding. These factors and its association with sero-positivity ( $p > 0.05$ ) is explained in (table 2,3,4,6,7,8,9, 10). Breed was not included in the analysis since most farmers in the area had local breeds.

#### c) Result of the Questionnaire Survey

A questionnaire survey was conducted on 100 livestock owners revealed that during the previous months lamb loss amounted to 60% (30% abortion, 12% stillbirth and 18% neonatal mortality). Birth of weak lambs amounted to 20% while reproductive abnormalities were 31% (17% dystocia, 8% retained fetal membrane and 6% endometritis). Seventy two respondents confirmed the presence of cats in their house hold kept for clearing rodents. Only 10% of interviewed individuals reported disposal of cat feces by burying in the ground. Thirty percent of them reported the disposal of cat feces on the backyard or grazing land, which increased the risk of exposure to toxoplasma oocyst and cats had close contact with most family members. The survey also showed that 83% of the interviewed people had a history of consumption of raw or under cooked meat. Sixty five percent of them had animals with poor hygienic management in their grazing area and drinking water. Fifty four percent of these owners informed that their livestock had a contact with dead animal carcass, which is improperly disposed. Thus further had contract the infection through ingestion of oocysts from these areas is a probable source of toxoplasmosis. In addition livestock owners revealed that there were wild cats coming to the grazing area of their livestock which act as a definitive host.

Table 1: Overall relative prevalence of toxoplasmosis in different species

Species	No. of animals examined	Positive	Prevalence (%)
Ovine	194	71	55.91
Caprine	40	15	11.53
Bovine	44	11	12.68
Swine	40	19	11.53
Equine	20	7	5.76
Camel	9	3	2.59
Total	347	126	100.00

$$\chi^2 = 1.2429 \text{ P} = 0.537$$

Table 2: Association between prevalence and origin of animals

Origin	No of examined	No of positive
Sholla vet clinic	74(21.33%)	28(22.22%)
Akaki Kaliti	215(61.96%)	75(59.22%)
A.A Abattoir Enterprise	58(16.71%)	23(18.25%)
Total	347(100.00%)	126(100.00%)

$$\chi^2 = 0.5445 \text{ P} = 0.762$$

**Table 3:** Association between Sero-prevalence and reproductive loss

Reproductive loss	No of examined	No. positive (%)
Present	269(77.52)	95(75.40)
Absent	78(22.48)	31(24.00)
Total	347(100.00)	126(100.00)

$\chi^2=0.5126$  p = 0.474

**Table 4:** Association between prevalence and cat ownership

Cat ownership	No of examined	No. positive (%)
Present	113(32.56)	35(27.78)
Absent	234(67.44)	91(72.22)
Total	347(100.00)	126(100.00)

$\chi^2=2.0645$  p=0.151

**Table 5:** Association between Sero-prevalence and sex

Sex	No of examined	No. positive (%)
Male	170(48.99)	76(60.32)
Female	177(51.01)	50(39.68)
Total	347(100)	126(100.00)

$\chi^2=10.1556$  p =0.001

**Table 6:** Prevalence on the basis of age

Age	No of examined	No. positive (%)
Old	106(30.55)	34 (26.98)
Adult	149(42.94)	56(44.44)
Young	92(26.51)	36(28.57)
Total	347(100.00)	126(100.00)

$\chi^2= 1.2429$  p = 0.537

**Table 7:** Prevalence on the basis of sex and species

Species	Sex	No of examined	No. Positive	Prevalence (%)
Ovine	Male	76	36	47.37
	Female	118	35	29.66
Caprine	Male	29	9	31.03
	Female	11	6	54.55
Bovine	Male	26	8	30.77
	Female	18	3	16.67
Swine	Male	31	18	58.06
	Female	9	1	11.11
Equine	Male	7	4	57.14
	Female	13	3	23.08
Camel	Male	1	1	100.00
	Female	8	2	25.00

**Table 8 :** Association between prevalence and management system

Management	No of examined (%)	Positive (%)
Extensive	228(65.71)	147(66.72)
Semi intensive	119(34.29)	74(33.48)
Total	347(100.00)	221(100.00)

$\chi^2=0.1771$  p=0.674

Table 9: Association between prevalence and hygienic condition

Hygiene	No of examined (%)	Positive
Poor	151(43.52)	55(43.65)
Good	196(56.46)	71(56.35)
Total	347(100.00)	221 (100.00)

$\chi^2=0.00125$   $p=0.969$

Table 10: Association between prevalence and clinical finding

Clinical Finding	No of examined (%)	Positive (%)
Respiratory	116(33.43)	38(30.16)
GI	69(19.88)	20(15.87)
Nervous	2(0.58)	2(1.59)
Skin and Mucosal	38(10.95)	16(12.70)
Metabolic	1(0.29)	1(0.79)
Poisoning (Toxicosis)	1(0.29)	0(0.00)
Normal	103(29.68)	44(34.92)
Traumatic Wound	3(0.86)	2(1.59)
Reproductive	14(4.03)	3(2.38)

#### IV. DISCUSSION

The overall Sero-prevalence of 126(36.1%) in the 347 study animals of different six species. Of which ovine (56.35%), caprine (11.9%), bovine (8.73%), swine (15.08%), equine (5.56%) and camel (2.38%) lies midway between the three previous studies in Ethiopia. The overall prevalence of 36.6% in sheep in this study lies between the three previous studies in Ethiopia. Bekele and Kassali, Getachew and Tilaye, and Tamiru reported Sero-prevalence of 22.9%, 33% and 54.7% in sheep, respectively. This is in agreement with studies in other African countries with prevalence rates ranging from 11.5% to 34% (Deconinck *et al.*, 1996). In goats the overall Sero-prevalence in this work was higher than the results of the two previous studies in Ethiopia with prevalence of 11.9% and 26.7% but less than those reported from other African countries with prevalence rates ranging from 31.9% to 63% (Tamiru, 2000). On the other hand, the finding agrees with recent study reports of Getachew and Tilaye of 35% in goats (Getachew and Tilaye, 2002). In bovine the overall Sero-prevalence (25%) out of 44 cattle examined. Thought number of animals studied were not proportional relative to previous studies, it is higher prevalent than the reports of Bekele and Kasali (2002) who reported a prevalence of 6.6%. In swine although those animals serum was taken from Addis Ababa abattoir and its number of sample size was not proportional relative to other species constitute the largest Sero-prevalence which is 47.5%. This result agrees with published reports in other parts of the world, the sero-positive prevalence in swine is 22% with a range of 0-97 % (Radostits *et al.*, 2007). The overall Sero-prevalence in equine species (horse and donkey) was 35%. The overall Sero-prevalence in camel (*Camelus dromedarius*) species was 33.3% which is high. This result revealed that a higher Sero-

prevalence when compared with previous studies in three ecologically different areas of Sudan (22.2%) (Khali *et al.*, 2007).

In the present study, no statistically significant difference in Sero-prevalence was noted among different origin, species, age groups, management system, hygienic condition, reproductive abnormalities, cat ownership and the associated clinical finding. This seems contradictory with the established facts, however, it is difficult to made firm conclusion as number of study animals is low in proportion with these factors. The odd ratio of the four factors (management system, reproductive abnormality, hygienic condition and cat ownership) is explained in figure 2. The prevalence would have been significantly higher in warm and moist areas than in cold or hot dry areas, increased with age, in extensive (small holder) management system than intensive type, prevalent in cat ownership and a major cause of abortion and neonatal mortality. The disagreement of the assessment of risk factors associated with sero-positivity to *T. gondii* in addition to the above reasons could be due to lack of the specificity of the serological test used. The slight variation in the results of Sero-prevalence observed can be attributed to variation in ecological conditions as most animals came from different areas to Addis Ababa. Variation may also be due to the diagnostic technique utilized (Assadi-Rad *et al.*, 1995). Even though the test used (slide agglutination test) was not done in Ethiopia, the results showed a higher Sero-prevalence than the previous studies it is evidenced that it is more sensitive and the result is valued in all species.

The results of the questionnaire survey indicated that economic loss due to abortion, stillbirth, neonatal mortality and related reproductive abnormality are important in the study area. It is also important that the inclusion of the questionnaire survey on ownership

of cats and purpose of cat keeping as well as the mechanism of cat feces disposal and the habit of raw meat and milk consumption. As oocysts are essential in the life cycle of *T. gondii* and in which both domestic cats and other felids may shed oocyst. These can contract the infection through contamination of grazing area as well as close contact with human (domestic cats) leading its public health hazard in addition to the owners habit of raw or under cooked meat as tissue cysts are the end stage of the parasite waiting to be eaten by animals and human (Morris *et al.*, 2007). The results of questionnaire survey based on its economic impact agrees with previous studies by Getachew and Tilaye in 2002 (abortion 30%, stillbirth 12%, neonatal mortality 18%, dystocia 17%, retained fetal membrane 8% and endometritis 6%). Seventy two (72) percent of the interviewed individuals had cats in their premises and kept for clearing rodents only together with livestock and almost all fed cats raw or under cooked meat. In general the maintenance of the cycle is achieved among the intermediate hosts, definitive host (cat) and environment (contaminated by infective stage of the parasite).

The Sero-prevalence in this study was significantly high both from public health and economic perspectives. Toxoplasmosis is a disease of economic importance as it is a major cause of abortion, stillbirth and neonatal mortality in sheep and goats (Getachew and Tilaye, 2002). Ovine abortion and neonatal mortality due to *T. gondii* are important problems in New Zealand, Australia, Canada, United States and the United Kingdom; in countries they are second in importance only to Chlamydia (Radiostits *et al.*, 2007). Several studies conducted so far indicated that *T. gondii* infection in humans is widely distributed in most tropical countries (Negash, 2000). The high Sero-prevalence in the study animals and the results of the questionnaire survey ensured that Toxoplasmosis in Addis Ababa suggests a high risk to humans. The recent study in Adama town of Ethiopia is an evidence for its prevalence in that of 65% people examined for anti-*Toxoplasma gondii* antibodies by the MDAT, serologic evidence of Toxoplasmosis was found in 60% (39/65) (Negash *et al.*, 2008).

Statistical analysis revealed that there is significant association between males and females with males having higher Sero-prevalence than females. This is in agreement with findings by Getachew and Tilaye observed in goats (Getachew and Tilaye, 2002). This could be due to the fact male animals are stressed due to transport from different ecological areas and as most of them were kept for feedlot and breeding purpose for long time.

## V. CONCLUSION AND RECOMMENDATIONS

In general, the Sero-prevalence survey conducted in this study showed that toxoplasmosis is a

widespread and well established infection among the six species (ovine, caprine, bovine, equine swine and camel) two veterinary clinics (Sholla and Akaki Kality) and Addis Ababa Abattoir Enterprise. The significance of toxoplasmosis as a disease of zoonotic importance and its economic impact was demonstrated. Therefore, prevention efforts should focus on educating cat owners about the importance of collecting cat feces in litter boxes, spaying cats, reducing the numbers of feral cats, cooking all meats, and promoting rigorous hand hygiene, reducing the numbers of wild rats is also important for control of toxoplasmosis. We think that further studies should be conducted to determine whether any host reservoirs exist amongst domestic and wild animals in this area, in which the disease was previously not found. This study will be the basis for further studies that will deepen our knowledge of the epidemiology of *T. gondii*. More extended studies are required to determine the sero-prevalence rates among populations of wild rats and other wild animals in difference areas, and the implications of *T. gondii* prevalence on both animal and human health.

### Conflict of interest

The authors have no declared any conflict of interest

## VI. ACKNOWLEDGEMENTS

Authors would like to thank Sholla and Akaki-Kaliti Veterinary Clinics and Addis Ababa Abattoir Enterprise for their technical support throughout the study period.

## REFERENCES RÉFÉRENCES REFERENCIAS

1. Assadi Rad, A.M., New, J.C. and Patton, S. 1995. Risk factors associated with transmission of *Toxoplasma gondii* to sows kept in difference management systems in Tennessee. *Vet. Parasitol.* 57: 289-297.
2. Bekele, T. and Kasali, O. B. 1989. Toxoplasmosis in sheep, goats and cattle in Ethiopia. *Veterinary Research Communication.* 13:5, 371-37.
3. Chauhan, R.S. and Agarawal, D.K. 2006. Text Book of Veterinary Clinical and Laboratory Diagnosis. 2<sup>nd</sup> ed. Jaypee Brothers Medical Publisher (P) Ltd, Newdelhi. 27: 226.
4. CSA (Central Statistics Authority), 2009. Agricultural sample survey (2008/9). Statistical bulletin 302. Adiss Ababa. Ethiopia.
5. Deconinck, P., Pangui, L.J., Akakpo, J., Garrouste, A., Ouattara, L., Roger, F. et al., 1996, 'Sero-epidemiology of toxoplasmosis in sheep and goats from Africa', *Revue de Médecine Vétérinaire*; 147(5), 377-378.
6. Dubey JP, Beattie CP. General Biology. In: Dubey JP, Beattie CP. Toxoplasmosis of Animals and Man. Boca Raton: CRC Press; 1988. p. 1-40.

7. Dubey, J.P., 2007. The history and life cycle of *Toxoplasma gondii*. In: Weiss, L.M., Kim, K. (Eds.), *Toxoplasma gondii. The Model Apicomplexan: Perspectives and Methods*. Academic Press, New York, pp. 1–17.
8. Dunford, P. and Johnson, J. 1991. Assessment of a slide agglutination test. *Lab. Sci Apr.* 48(2):137-141. Public Health Laboratory Services George 'S Hospital, London, England, UK.
9. Getachew, T. and Tilaye, D. 2002. Study on Toxoplasmosis in Debre Birhan and the surrounding areas in Ethiopia. Faculty of Veterinary Medicine Debrezeit and Institute of Pathobiology, Addis Ababa University, Addis Ababa, respectively. *Bull. Anim. Hlth. Prod. Afri.* 50, 138-147.
10. Hartley, W.J., Marshall, S.C., 1957. Toxoplasmosis as a cause of ovine perinatal mortality. *N.Z. Vet. J.* 5, 119–124.
11. Kahli, K.M, Gadir, A.A, Rahman, M.M, Yassir, O.M, Ahmed, O.M, Ahmed, A.A and Elrayah, I.E. 2007. Prevalence of *Toxoplasma Gondii* antibodies in camel and their herders in 3 ecologically different areas in Sudan. *Jor. Cam. Prac. Res* 14(1): 11-13.
12. Morris, J. A., Aulisie, C. G. And McCown, J. M. 2007. Serological evidence of toxoplasmosis in animals. *Journal of Infectious diseases*. Oxford University Press.
13. Negash, T., Tilahun, G. and Medhin, G. 2008. Sero-prevalence of *toxoplasma gondii* in Nazareth Town, Ethiopia. *East African Journal of public health*, 5: 3.
14. Negash, T., Tilahun, G., Patton, S., Prevot, F. and Dorchie, P. H. 2004. Serological survey on Toxoplasmosis in sheep and goats in Nazareth, Ethiopia. *Revue Med. Vet.*, 155, 10:486-487.
15. Radostitis, O.M., Gay, C.C. Hinch Cliff, K.W. and Constable, P.D. 2007: A text book of the disease of cattle, horse, sheep, goats and pigs. *Veterinary medicine*, 10<sup>th</sup> ed Sounders. Elsevier, New York, Pp 1517-1522.
16. Singh. And Msolla, P. 1994. In: Proceedings of the Second Tanzanian Veterinary Association Scientific Conference. *J. Tanz. Vet. Assoc.* 2: 210.
17. Tamiru N. (2000). Study on toxoplasmosis. Faculty of Veterinary Medicine. Addis Ababa University. Debre-Zeit, DVM-Thesis.
18. Teshale, S., Dumedere, A., Darde, M. L., Mega. And Dorchie, P. 2006. Serological survey of caprine toxoplasmosis in Ethiopia: prevalence and risk factors. Faculty of Veterinary Medicine, Deberzeit, Ethiopia.
19. Thrusfield, M. (1995). *Veterinary Epidemiology*. 2nd ed. Black well science Ltd. UK.
20. Wassserthiel-Smaller. 1995. *Biostastics and Epidemiology: A Primer for Health Professionals*. 2<sup>nd</sup> ed. Springer, Verlag. New York, U.S.A. 304.
21. Wolf, A., Cowen, D., Paige, B., 1939. Human toxoplasmosis: occurrence in infants as an ncephalomyelitis verification by transmission to animals. *Science* 89, 226–227.



This page is intentionally left blank





GLOBAL JOURNAL OF MEDICAL RESEARCH: G  
VETERINARY SCIENCE AND VETERINARY MEDICINE  
Volume 16 Issue 3 Version 1.0 Year 2016  
Type: Double Blind Peer Reviewed International Research Journal  
Publisher: Global Journals Inc. (USA)  
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

## Prevalence of Bovine Cysticercosis and Status of Human Taeniasis in and Around Asella Town, Tiyoworeda, South East Ethiopia

By Adem Edao, Feyera Gemed Dima & Feyissa Begna Deressa

*Jimma University*

**Abstract- Background:** Bovine cysticercosis and human taeniasis is an important parasitic disease and more common in developing countries including Ethiopia where meat is an important component of human diet and traditionally consumed raw on several occasions.

**Methodology:** Cross-sectional study was conducted from November 2014 to April 2015 at Asella municipal abattoir in Arsi zone of Oromia to determine prevalence of bovine cysticercosis and status of human taeniasis with its associated risk factors around Asella town. Routine meat inspection method and questionnaire survey were performed for this study.

**Result:** From the total of 430 carcass inspected, 5(1.2%) were positive for bovine cysticercosis. The statistical analysis revealed that risk factors like sex, breed, body condition and ages are not associated with the prevalence of cysticercosis. The questionnaire survey showed that 44.3% of total respondents were infested with *Taenia saginata* at least once in their life time and out of these positive respondents 32% and 6.7% become positive only before and after 2000 E.C respectively, while 5.6 % were infested many times.

**Keywords:** bovine, cysticercosis, taenia saginata, prevalence, risk factors, taeniasis.

**GJMR-G Classification:** NLMC Code: WC 838



*Strictly as per the compliance and regulations of:*



© 2016. Adem Edao, Feyera Gemed Dima & Feyissa Begna Deressa. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License (<http://creativecommons.org/licenses/by-nc/3.0/>), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

# Prevalence of Bovine Cysticercosis and Status of Human Taeniasis in and Around Asella Town, Tiyyoworeda, South East Ethiopia

Adem Edao <sup>α</sup>, Feyera Gemedo Dima <sup>ο</sup> & Feyissa Begna Deressa <sup>ρ</sup>

**Abstract- Background:** Bovine cysticercosis and human taeniasis is an important parasitic disease and more common in developing countries including Ethiopia where meat is an important component of human diet and traditionally consumed raw on several occasions.

**Methodology:** Cross-sectional study was conducted from November 2014 to April 2015 at Asella municipal abattoir in Arsi zone of Oromia to determine prevalence of bovine cysticercosis and status of human taeniasis with its associated risk factors around Asella town. Routine meat inspection method and questionnaire survey were performed for this study.

**Result:** From the total of 430 carcass inspected, 5(1.2%) were positive for bovine cysticercosis. The statistical analysis revealed that risk factors like sex, breed, body condition and ages are not associated with the prevalence of cysticercosis. The questionnaire survey showed that 44.3% of total respondents were infested with *Taenia saginata* at least once in their life time and out of these positive respondents 32% and 6.7% become positive only before and after 2000 E.C respectively, while 5.6 % were infested many times. A statistical analysis showed that human taeniasis prevalence is significantly different ( $p < 0.05$ ) among the categories of the considered risk factors like: residence area, age, sex, raw meat consumption, education and occupational status. Religion difference was not a problem for Taeniasis prevalence distribution.

**Conclusion:** In general, the current study revealed that the presence of bovine cysticercosis was small but it's still a public health hazard in the study area which needs increased awareness about the health impact of taeniasis.

**Keywords:** bovine, cysticercosis, taenia saginata, prevalence, risk factors, taeniasis.

## I. INTRODUCTION

Livestock in developing countries play a crucial role in improving food security, generating cash income and are an asset. The total livestock population in Ethiopia according to 2014 estimation was 56.71 million cattle, 29.33 million sheep and 29.11 million goats (CSA, 2015), which places Ethiopia first in Africa and ninth in the world in terms of total stock population. From the total cattle population 98.95% are local breeds and the remaining are hybrid and exotic breeds. Cattle constitute large portion of livestock population and are managed

by small holder farmers under extensive low input traditional system (CSA, 2015). However, its productivity remains marginal due to prevalent diseases, malnutrition and management constraints. Among that *T. saginata*/bovine cysticercosis is the one which remains a major public and animal health problem (EARO, 2000).

Bovine cysticercosis is an infection of cattle caused by the larval stage *T. saginata* which live in human intestinal. This parasite is universally distributed in developing as well as in developed countries (Gracey and Collins, 1992; Cabaret *et al.*, 2002; Dorny *et al.*, 2009). In humans, the disease is called taeniasis which is accompanied with symptoms like nausea, abdominal discomfort, epigastric pain, diarrhea, excessive appetite or loss of appetite, weakness, loss of weight and intestinal blockage. Sometimes, the mobile gravid segments may make their way to unusual sites such as the appendix and biliary tract and may cause serious disorders (WHO, 2013). Live cattle having *C. bovis* shows no symptoms, however, heavy infestation by the larvae may cause myocarditis or heart failure (Gracey and Collins, 1992). Cysticerci can remain alive in cattle anywhere from weeks to years and such infection in cattle is a public health problem as the infected raw or undercooked beef causes taeniasis in human (Garcia, 2003; Garcia *et al.*, 2007). It has economic significance as well as the economic losses accruing from the condemned and downgraded carcasses and due to treatment of carcasses before human consumption is substantial (Yoder *et al.*, 1994; Onyango-Abuje, 1996; Giesecke, 1997).

Bovine cysticercosis and taeniasis are common where hygienic conditions are poor and the inhabitants traditionally eat raw or insufficiently cooked or sun-cured meat (Minozzo *et al.*, 2002). Inadequate health education and low availability of taenicides are the major obstacles for the control of such infections (Pawlowski, 1996). Due to these reasons, taeniasis is more common in developing countries including Ethiopia where meat is an important component of human diet and traditionally consumed raw on several occasions. Lack of awareness about raw meat consumption, existence of highest population density, poor hygiene and sanitary facilities some of the factors that facilitate transmission (Jones *et al.*, 1997).

**Author p:** DVM, MSc in Epidemiology and Zoonotic Microbiology School Veterinary Medicine, College of Agriculture and Veterinary Medicine, Jimma University. e-mail: fey\_abe2009@yahoo.com

The epidemiology of bovine cysticercosis/human taeniasis varies from one area to another so control measures appropriate in one area is not necessarily of value in another. Hence, it is essential to have adequate knowledge of the epidemiology of the disease before contemplating control programmes. In Ethiopia some studies have been conducted on bovine cysticercosis at different times. But the studies performed were limited to few parts of the country and there was no information on prevalence of bovine cysticercosis and human taeniasis in and around Asella district. Therefore, the objectives of this study were:

- To determine the prevalence of bovine cysticercosis at Asella municipal abattoir,
- To estimate status of taeniasis and associated risk factors in and around the study area

## II. MATERIAL AND METHODOLOGY

### a) Description of the Study Area

The present study was conducted at Asella municipal abattoir, and Asella town and its surrounding, Tiyyoworeda. Asella town is a capital of Arsi Zone, Oromia regional state, Ethiopia. It is located about 175 km Southeast of Addis Ababa at 6° 59' to 8° 49' N latitudes and 38° 41' to 40° 44'E longitudes. The altitude of the area ranges from 2500 to 3000 m.a.s.l. Asella and its surrounding is characterized by mid sub-tropical weather, with minimum and maximum temperature ranging from 8.4 to 22.6°C, and the relative humidity ranging from 43 to 60%. The average rainfall is 2000mm. The area has a bimodal rainfall occurring from March to April (short rainy season) and July to October (long rainy season). According to Arsi Planning Economic and Development Office (APEDO, (2007)), the area is densely populated, with livestock population of 85,893 cattle, 57,118 sheep, 10,725 goats, 7841 horses, 15,642 donkeys, 517 mules and 35,489 poultry. The farmers in the area practice mixed crop-livestock farming system.

### b) Study Population

The study animals comprise indigenous cattle brought to Asella Municipal abattoir for slaughter from different districts in and around Assela town and it includes cattle of different age, sex, breed and body condition categories. For the survey data, the target populations were residents of Asella town and surrounding kebeles (Burqa cilalo, Dosh, Harobilallo, Café Misoma and Gora silingo) and includes all age group >18 and both female and males

### c) Study Design

A cross-sectional study was used to study the prevalence of bovine cysticercosis at Asella municipal abattoir by using routine meat inspection technique in municipality slaughter house for the presence of *C. bovis*. Moreover, a cross-sectional study was conducted

by a semi-structured questionnaire survey to assess the status of *T. saginata*/taeniasis and associated risk factors.

## III. SAMPLE COLLECTION METHOD

### a) Active Abattoir Survey

Animal samples were collected by active abattoir survey. The study animals were selected using simple random sampling method and age, breed, sex and body condition of each study animals was recorded on prepared format paper at ante-mortem. Body condition scoring of the cattle was made based on the guideline provide by (Nicholson and Butter, 1986) and all animals included in the study were animals with, medium and good body condition. Age determination was carried out by means of their dentition as described by (De-lahunta and Habel, 1989) and all the inspected animals were at the age of adult and old age group. Prior to sampling, each selected animal were given an identification number by writing a code on its head by using unwashable ink.

Meat inspection was made in accordance with the procedures of Ethiopian Ministry of Agriculture Meat Inspection Regulation (MOA, 1972) for the detection of *T. saginata*'s cysts. Because of owners' discomfort on multiple incisions for the thorough examination of major muscles, only the masseters muscle and internal organs such as tongue, heart, liver, kidney, lung and diaphragm were used as indicators of the presence of cysts in the carcass. Careful examination on the carcass of study unit was made through palpation of the organs followed by incision as follows: the surface and substance of tongue was examined visually, followed by longitudinal ventral incision from the tip of the root. Extensive deep incision was made into external and internal muscles of masseters parallel to the plane of the jaw (parallel to the jaw bone from the lower jaw). Visual inspection and longitudinal incision of the myocardium from base to apex was made. The muscles of diaphragm were examined visually and by making incision. Examination of kidney, liver, and the lung was also conducted accordingly by visualization, palpation and incision. Cysts observed in these organs were carefully dissected and numbers and nature of cysts in each organ was recorded for each animal. The nature of the cyst was recorded as calcified and viable by visual observation of its appearance, as (Ashwani and Gebrehiwot, 2011) dead degenerated or calcified cysticerci clearly form identifiable spots of white and have fibrotic lesions, while the viable cysticerci are pinkish-red in colour.

### b) Questionnaire survey

Semi-structured questionnaire survey used to assess, potential risk factors and its public health importance. Questionnaire survey on the disease

occurrence and risk factor assessment was administered to 415 volunteer respondents who were selected based on convenience sampling method and the interview was conducted phase to phase. The potential risk factors of taeniasis considered in this study were :age, sex, religion, occupation, educational status, and habit of raw meat consumption. The awareness of the diseases, presence/absence of sanitation facilities like toilet and the drug used for treatment were also included in the questionnaire survey and at the end of interview some advices were provided to the respondents on how to control and prevents *T. Saginata* infection/cysticercosis.

### c) Sample Size Determination

The desired sample size was calculated using the standard formula described by Thrus field (Thrus field, 2005) for simple random sampling method. Since there was no similar previous study at this area, expected prevalence was considered 50%, 5%desired absolute precision and 95% confidence level were used to calculate the minimal sample size. Hence, the sample size required was 384 heads of cattle. But to increase the precision of the study the sample size were increased and a total of 430 heads of cattle were included in the current study. For questionnaire survey sample size was calculated by using the formula given by Arsham (2002) which is:

$$N = 0.25/SE^2,$$

When: N= sample size, SE (standard error) =5%),

The sample size required for the questionnaire survey as per the above formula is 100 for each site (urban and rural). However, to include different risk factors and increase the precision of the result the total number was increased to 415 individuals.

## IV. DATA MANAGEMENT AND ANALYSIS

The data collected were entered, recorded and stored in Microsoft excel spread sheets program version 2010. Descriptive and inferential analyzed was made by using SPSS version 20 software. Fisher's and Chi-square ( $X^2$ ) tests were used to determine the variation in prevalence of infection between different groups for abattoir and questionnaire survey respectively. Statistical significance level was set at  $P < 0.05$  at 95% confidence level to determine whether there are statistical significant differences between the parameters measured.

## V. RESULT

Prevalence of *C. bovis*: the study showed that from a total number of 430 carcasses inspected, 5(1.2%) were positive for bovine cysticercosis. The statistical analysis of the data revealed that no significant difference ( $p > 0.05$ ) was observed in the prevalence of cysticercosis in relation to the risk factors

like: sex, breed, body condition and ages. But high prevalence of *C. bovis* was observed in local breed, old age and female animals relative to cross breed, adult age and male animals respectively, while almost no difference was indicated between animals of good and medium body condition as shown in Table 1.

*Anatomical distribution of cysts*: frequency analysis of active abattoir survey revealed that nearly the there was the same distribution of *C. bovis* in the examined organs as follows: liver (0.9%), heart (0.7%), tongue (0.5%) and masseters muscle (0.5%). From the total number of 16 *C. bovis* observed on different organs, 7(43.8%), 5(31.3%), 2(12.5%) and 2(12.5%) were localized on the liver, heart, tongue and masseters muscle (table 2) and out of the total cyst observed 6 (37.5%) of them were viable with organ distribution of 50%, 33.3% and 16.7% on liver, heart and masseters muscle respectively as indicated in table 2.

*Questionnaire survey*: Of the total 415 interviewed respondents 44.3% (184/415) had contracted *T. saginata* infection at least once in their life time and the respondents confirmed that they were positive by witnessing they observed proglottids in their feces and/or under wear. Since due to religious purpose pork meat is not consumed in the study area, the proglottids observed were surely to be of *T. saginata*. From positive respondents 32% got taeniasis only before 2000E.C (2008G.C), while 6.7% got taeniasis after 2000 E.C (2008 G.C) and 5.5% were those remain positive both before and after 2000E.C. Out of those witnessed their positivity, 88.6% and 10.9% respondents used modern and traditional drugs for treatment respectively, while the remaining percent uses nothing for treatment as indicated by tables 4 respectively. Out of total respondents 90.6% (376/415) uses toilet, 73.5% (305/415) have awareness about human taeniasis.

*Association of risk factors with prevalence of taeniasis*: A statistical analysis showed that there was a highly significant variation between urban and rural, higher and lower age group, raw meat and cooked meat consumers, female and male, occupation group, and educational levels ( $p < 0.05$ ). High prevalence of human taeniasis was reported in rural area, male, higher age group, farmer, raw meat consumer and illiterate. But no statistical significance variation was seen among religion ( $p > 0.05$ ) as shown below by tables 5.

## VI. DISCUSSION

The prevalence of *C. bovis* among the carcasses inspected at Asella municipal abattoir was 1.2% which is comparable with the findings of (Birhanu, *et al.*, 2013) who reported prevalence of 2.58% from Bahir Dar Municipal Abattoir, 2.59% from Wolaita sodd municipal abattoir (Dawit, *et al.*, 2012), 3% from Zeway Municipal Abattoir (Bedu, 2011), 3.6% from Addis

Ababa abattoir (Nuraddis and Frew, 2012), 3.65% from Jimma municipal abattoir (Taresa, *et al.*, 2011), 3.11% indifferent agro climatic zones of Ethiopia (Tembo, 2001). But lower than the finding of (Dawit, 2004) 4.9% at Gondar, (Alula, 2010) 5.4% at Konbolcha, (Kebede, 2008) 18.49% in North West Ethiopia, (Abunna, 2008) 26.3% at Hawassa, (Hailu, 2005) 17.5% in East Shoa, (Fetene and Nibret, 2014) 5.1%, at Jimma municipal abattoir, (Belay, 2014) 5.2% at Municipal Abattoir of Shire, (Abunna, 2013) 12% at Yirgalem and (Lielt, 2015) 5.6 at Bishoftu/Elfora abattoir. This difference might be resulted from difference in the level of personal and environmental hygiene, habit of raw meat consumption, number of incision made at inspection site in the abattoir, and management type of the animals practiced. The main reason with low prevalence of bovine cysticercosis in the current study could be due to low number of organs inspected and low incision made at inspection site at the abattoir. In Asella municipal abattoir the commonly inspected organs for presence of *C. bovis* were internal organs (liver, heart, lung, tongue, and kidney) and masseters muscle while other predilection sites are rarely inspected due to multiple mutilation of carcass causes reduction in marketability of the meat and the owners not permit multiple incision of heavy muscles. This may in turn lead to omitting of infected animals as the sensitivity of detecting the parasite will decline with limited number of incisions (Wanzala, 2003) and experimental studies showed a 5-50 times higher prevalence will be achieved by complete slicing of the predilection sites (Minozzo, 2002).

But the current study's result was higher relative to the finding of (Zdolec *et al.*, 2012) and (Blessing *et al.*, 2011) who reported 0.11% and 0.2% from Croatia and South Africa. This could be due to strict application of meat inspection and public health extension rules and difference in hygiene measures in the study countries. It's known that sanitation facilities are better in the developed countries than in the developing countries in which poor environmental hygiene increase the prevalence of the diseases in the environment. More number of *C. bovis* was observed in the liver than other organs inspected. The reason is that absence of specific predilection site for *C. bovis* as stated by (Scandrett, 2009) so more number of larvae collected through mesenteric and portal veins residing in the liver.

Questionnaire survey indicated that human taeniasis was common in the study area with prevalence of 44.3%. This agree with the finding of (Mesfin and Nuraddis, 2012) 44% in Hawassa town and (Dawit and Temesgen, 2013) 44.44% in Shire Indasilassie district. But this result is lower relative to the finding of (Lielt *et al.*, 2015) 64% in Bishoftu, (Dawit, 2012) 62.5% in Wolaittasoddo, (Fetene and Nibret, 2014) 58%, (Abunna, 2013) 70% in Yirgalem, (Bedu *et al.*, 2011) 56.7% at Zeway, (Abunna *et al.*, 2008) 64.2% in Awassa

town and (Megersa *et al.*, 2010) 56.7% Jimma town. The reason for reporting lower prevalence of human taeniasis in the current study area could be due to the difference in the religious composition of the respondents, and sample size taken. Out of total respondents of the current study, 45.1% were Muslims that they have traditionally low habit of consuming raw meat than Christians and from the total respondents only 50.8% were raw meat consumers. Raw meat consumption is the only way of getting *T.saginata* infection, so as raw meat consumption decrease in the area the infection also decrease. The other is sample size difference and as sample size increase the precision will also increases. In the present study the sample size is very large (415) while in the above finding very small (not greater than 170). The other point is that some respondents shy to openly tell about taeniasis and this could also end up with low recovery of positive people in the study area.

It was revealed that *T.saginata* infection was more prevalent in the rural area than urban. This could be due to low level of personal and environmental sanitation facilities and absence of meat inspection in the rural area than urban area. This finding agrees with the statement of (Minozzo *et al.*, 2002) who stated that taeniasis are common where hygienic conditions are poor and the inhabitants traditionally eat raw or insufficiently cooked or sun-cured meat.

The current study indicates that the ages of respondents have strong association with the prevalence of *T.saginata* infection and high in higher age group (>35 years) than other age group (<18 and 18-35 years) (table 5) which is in agreement with the previous study of (Abunna *et al.*, 2008; Adugna *et al.*, 2013; Lielt *et al.*, 2015). This might be due to that the habit of raw meat consumption increase with age and the higher age group have better income to consume raw meat and more prone to *C.bovis*. But lower age groups are student that they have no sufficient access to raw meat from butcher's houses as commonly raw meat was eaten at this site and they are less invited than elder on different ceremony where raw meat consumption culturally practiced.

In this study *T. saginata* was more prevalent in men than in women which is a similar report with findings of (Hailu, 2005; Abunna, 2013; Lielt *et al.*, 2015) in other parts of Ethiopia. This may be due to cultural practice in Ethiopia that men not commonly prepare their dishes at home rather they frequently visiting but cheries and restaurants than women for beef consumption. In this study no statistically significant difference were observed between the proportion of taeniasis in Muslim and Christian community which is in consent with the previous reports of (Tembo, 2001; Abunna, 2007; Dawit *et al.*, 2012). The reason behind may be they share same culture and habit of raw meat

consumption in the study area regardless of their religion.

The current study revealed that raw beef consumers had contracted taeniasis infection more frequently than the non raw beef consumers which is inlined with the report of (Megersa, 2010; Fetene and Nibret, 2014). The reason is well known that in the consumption of raw meat the degree of ingesting *C.bovis* with meat is higher (Gajadhar *et al.*, 2006; Garcia *et al.*, 2007). *T. saginata* infection is highly prevalent in the illiterate than literate respondents and this report agrees with the finding of (Abunna *et al.*, 2008; Kebede *et al.*, 2009; Adugna *et al.*, 2013) and also the current study revealed that the farmers had contracted taeniasis than individuals with other occupational status. This difference might be from low level of awareness in the illiterate and farmers than literate individuals and other occupational status. The other reason for reporting high prevalence of taeniasis in the farmer community is that most of Ethiopian farmers are illiterate and from rural area where environmental hygiene is low and backyard slaughter was practiced with very low awareness about the diseases.

The questionnaire survey result showed that the prevalence of taeniasis in human population is decreasing and it also indicated that there was strong relationship between occurrence of *T.saginata* infection and residence area, age, sex, habit of raw meat consumption, occupational and educational status of the respondents. Therefore, continues public education should be provided to avoid consumption of raw meat and encourage use of latrines and improved standards of human hygiene and backyard slaughtering of cattle should be restricted and slaughter house which fulfills the necessary facilities and with qualified meat inspector should be constructed.

## VII. ACKNOWLEDGEMENTS

It is our proud privilege to express our sense of thankfulness to College of Agriculture and Veterinary Medicine, School of Veterinary Medicine for providing the financial aid to complete this project. We also highly thankful to Ato Haji Geleto the meat inspector of Asella municipal abattoir, for his valuable suggestions, experience share during the abattoir visits.

## RÉFÉRENCES

1. Central Statistical Agency (CSA). Agricultural Sample Survey, 2014/15, Volume II: Report on Livestock and livestock characteristics (Private peasant holdings). Statistical Bulletin 570. Central Statistical Agency (CSA), Federal Democratic Republic of Ethiopia, Addis Ababa. 2015.
2. Ethiopian Agricultural Research Organization (EARO). Beef Research Strategy. Ethiopian Agricultural Research Organization, Animal Science Directorate, Addis Ababa.2000: Pp. 1–16
3. GraceyJF, Collins, DS. Meat Hygiene, 9<sup>th</sup>edn.Balliere Tindal, London.1992.
4. Cabaret, J., Geerts, S., Madeline, M., Ballandonne, C. and Barbier, D. The use of urban sewage sludge on pastures: The cysticercosis threat. 2002; *Vet. Res.*, **33**, 575–597.
5. Dorny P, Praet N, Deckers., Gabriel S. Emerging food-born parasites. *Vet. Parasitol* 2009; **163**: 196-206.
6. World Health organization (WHO). Taeniasis/ cysticercosis. WHO Fact sheet N.376°. 2013.
7. Garcia LS. Parasitology: In *Manual of Clinical Microbiology* 8<sup>th</sup> ed. Washington, D.C.: ASM Press. 2003: Pp. 1895.
8. Garcia LS, Jimenez JA, Escalante H. Cestodes: In *Manual of Clinical Microbiology* 9<sup>th</sup> ed., Washington, D.C.: ASM Press. 2007: p 2166.
9. Yoder DR, Eblell ED, Hancock DD, Combs BA. Public Veterinary Medicine: Food Safety and Handling, Epidemiological Findings from an Out Break of Cysticecosis in Feed lot Cattle. *J.A.V.M.* A1994; **205**: 75-86.
10. Onyango-Abuje JA, Nginyi JM, Rugutt MK, Wright SH, Lumumba P, Hughes G, Harrison JS. Seroepidemiological survey of *T. saginata* cysticercosis in Kenya. *Vet. Parasitol* 1996; **64**: 177–185.
11. Giesecke WH. Prevalence and economic implications of taeniasis/cysticercosis in South Africa. Report on a Workshop held at the Onderstepoort Veterinary Institute, Onderstepoort, South Africa; 18-19August 1997. Pp. 19-70.
12. MinozzoJC, GussorLF, De CastroEA, Lago O,Socci VT. Experimental Bovine Infection with *Taeniasaginata* Eggs: Recovery Rates and cysticerci Location. *Braz. Arch. Biol. Technol* 2002; **45**: 451-455.
13. Pawlowski ZS. Helmenthic Zoonosis Affecting Humans in Africa. Veterinary Medicine, Impacts on Human Health and Nutrition in Africa. Proceedings of an International conference Lindberg. 1996: Pp. 50 - 71.
14. Jones TC, Hunt RD, Norval WK. Veterinary Pathology. 6<sup>th</sup> ed. Hong Kong: Lippincott Williams and Wilkings 1<sup>st</sup> ed. USA: Mosby. 1997; **431**: 651-654.
15. APEDO (Arsi Plan and Economic Development Office). Socio Economic on Arsi Zone; Assela, Arsi. 2007: 1-105.
16. Nicholson MJ, Butterworth MH.guide to body condition scoring zebu cattle international livestock center for Africa, Addis Ababa, Ethiopia.1986.
17. De-lahunta A, Habel R. Teeth apply Veterinary Anatomy W.B.Saunder Company. 1989: Pp. 4-6.

18. Ministry of Agriculture, Meat Inspection Regulations (MOA). Legal notice No. 428 Negarite Gazeta. Addis Ababa, Ethiopia.1972.
19. Ashwani K. and Gebrehiwot T. Bovine cysticercosis in Ethiopia: a review. *Ethiop. Vet. J.*, 2011;**15**: 15-35
20. Thrusfield M. *Veterinary epidemiology*. 3<sup>rd</sup> ed. Singapore, Black well Science. 2005: Pp: 233.
21. Arsham, H. Questionnaire Design and Surveys Sampling, SySurvey: The Online Survey Tool. <http://home.ubalt.edu/ntsbarsh/Business-stat>. 2002.
22. Birhanu M, Habtamu A, Dawit T. Study on Zoonotic Metacestodes of Cattle Slaughtered at Bahir Dar Municipal Abattoir, Northwest Ethiopia *Glob.Vet*. 2013; **10**: 592-598
23. Dawit T, Tewodros S, Tilaye D. Public Health and Economic Significance of Bovine Cysticercosis in Wolaita Sodo, Southern Ethiopia. *Glob. Vet* 2012; **9**: 557-563.
24. Bedu, H., Tafess, K., Shelima, B., Woldeyohannes, D. and Amare, B., Bovine Cysticercosis in Cattle Slaughtered at Zeway Municipal Abattoir: Prevalence and its Public Health Importance. *J. Vet. Sci. Technol* 2011; **2**: 108.
25. Nuraddis I, Frew Z. Prevalence of *Tania Saginata* cysticercosis in Cattle Slaughtered in Addis Ababa Municipal Abattoir, Ethiopia. *Glob. Vet* 2012; **8**: 467-471.
26. Taresa G, Melaku A, Basazenuw B, Chanie M. cyst viability, body site distribution and public health significance of Bovine Cysticercosis at Jimma south west Ethiopia. *Glob. Vet* 2011; **7**: 164-168.
27. Tembo A. Epidemiology of *Taenia saginata*, Taeniasis/Cysticercosis in three selected agro-climatic zones., Faculty of Veterinary Medicine Free University of Berlin, Berlin.2001;MSc. Thesis.
28. Dawit S. Epidemiology of *Taenia saginata*, taeniasis and cysticercosis in North Gondar zone Northwestern Ethiopia., Faculty of Veterinary Medicine, Addis Ababa University: DebreZeit, Ethiopia.2004:DVM thesis
29. Alula, A. Major Metacestodes in Cattle Slaughtered at Kombolcha ELFORA Abattoir, North East Ethiopia: prevalence, cyst viability organ distribution and socio economic implication. Faculty of Veterinary Medicine, Hawassa University,.Hawassa, Ethiopia. 2010; DVM Thesis.
30. Kebede N. Cysticercosis of slaughtered cattle in northwestern Ethiopia. *Research in Veterinary Science*. 2008; **85**: 522-526.
31. Abunna, F., Tilahun, G., Megersa, B., Kumsa, B., and Regassa, A. Bovine cysticercosis in cattle slaughtered at Awassa municipal abattoir, Ethiopia: Prevalence, cyst viability, distribution and its public health implication. *Zoon and Publ. Health* 2008; **55**: 82-88.
32. Hailu D. Prevalence and risk factors for *T. Saginata* cysticercosis in three selected areas of eastern Shoa. Addis Ababa University, Faculty of Veterinary Medicine, Debrezeit, Ethiopia.2005: M.Sc thesis.
33. Fetene F, Nibret M. Prevalence of Bovine Cysticercosis in Cattle and Zoonotic Significance in Jimma Town, Ethiopia. *Acta Parasitol Glob* 2014; **5**: 214-222.
34. Belay S. and Mekelle BA. Prevalence of *Cysticercus bovis* in Cattle at Municipal Abattoir of Shire. *J. Vet. Sci. Technol*. 2014; **5**: 196.
35. Abunna, F. Prevalence, organ distribution, viability and socioeconomic implication of *bovine cysticercosis/teniasis*, Ethiopia *Revue d'élevageet de médecinevétérinaire des pays tropicaux* 2013; **66**: 2530.
36. Lielt E, Desalew T, Tsegabirhan K, Teshale S, Yohannes H. Prevalence and significance of bovine cysticercosis at Elfora abattoir, Bishoftu, Ethiopia, *journal of public health and epidemiology*, 2015; **7**: 32-40.
37. Wanzala W, Onyango-Abuje J, Kang`Ethe E, Zessin K, Kyule N, Bauman M, Ochanda H, Harrison. Analysis of Post-Mortem Diagnosis of Bovine Cysticercosis in Kenyan Cattle, *Onderstepoort. J. Vet. Res* 2003; **1**: 28-31.
38. Zdolec N, Vujević I, Dobranić V, Juras M, Grgurević N, Ardalić D. and Njari B. Prevalence of *Cysticercus bovis* in slaughtered cattle determined by traditional meat inspection in Croatian abattoir from 2005 to 2010; *Inst.Parasitol. SAS, Košice, Helminthol* 2012; **49**: 229 – 232
39. Blessing MD, Ethel KS, Mphane MM, Lebogang E, Francis RB, Rendani VN, Mathew N. Prevalence of Bovine Cysticercosis in the North West Province of South Africa from 2000 to 2010, *J. Hum. Ecol* 2011; **36**: 9-12.
40. Scandrett B, Parker S, Forbes L, Gajadhar A, Dekumyoy P, Weikagul J, Haines D. Distribution of *Taeniasaginata* Cysticerci in Tissues of Experimentally Infected Cattle, *Vet. Parasitol* 2009; **164**: 223-231.
41. Mesfin B, Nuraddis I. Prevalence of *Cysticercus Bovis* in Hawassa Municipal Abattoir and its Public Health Implication, *American-Eurasian J. Sci. Res* 2012 **7**: 238-245.
42. Dawit G, Temesgen M. Risk Factors and Public Health Significance of Cysticercosis in Cattle and Human in Shire Indasilassie District, Northern Ethiopia *Adv. Biolog. Res* 2013; **7**: 282-287.
43. Megersa B, Tesfaye E, Regassa A, Abebe R, Abunna F. Bovine Cysticercosis In Cattle Slaughtered at Jimma Municipal Abattoir, South Western Ethiopia: Prevalence, Cyst Viability and Its Socio Economic Importance. *Vet. World* 2010; **3**: 257-262.



44. Adugna, T., Yacob, H.T., Dinka, A. and Getachew, T. Bovine cysticercosis and human taeniosis in South-west Shoa zone of Oromia Region, Ethiopia. *Ethiopia.Vet.Journal* 2013; **17**: 121-133
45. Abunna, F., Tilahun, G., Megersa, B. and Regassa, A. Taeniasis and its socio-economic Implication in Hawassa Town and its surrounding, Southern Ethiopia. *East Afric J. Public Health* 2007; **4**: 73-79.
46. Gajadhar AA, Scandrett WB, Forbes LB. Overview of food- and water-borne zoonotic parasites at the farm level. *Revue Scientifique ET Technique (International Office of Epizootics)* 2006; **25**: 595-606.
47. Kebede N, Tilahun G, Hailu A. Current Status of Bovine Cysticercosis of Slaughtered Cattle in Addis Ababa Abattoir, Ethiopia. *Tropical Animal Health and Production* 2009; **41**: 291-294.

**Table 1:** Prevalence of bovine cysticercosis in cattle based on breed, sex, body condition and age from Assella, Tiyo woreda, Ethiopia

Risk factor	Category	No. of examined	No. of affected	Prevalence (%)	Fisher's exact p- value
Breed	Local	418	5	1.2	0.868
	Cross	12	0	0	
Age	Old	58	2	3.45	0.136
	Adult	372	3	0.8	
Sex	Male	413	4	1	0.183
	Female	17	1	5.9	
BCS	Medium	281	3	1.1	0.564
	Good	149	2	1.3	
	Total	430	5	1.2	

**Table 2:** Frequency distribution of *C.bovis* in different organs examined and number of organs infected from cattle slaughtered at Assella, Tiyoworeda Abattoir

Organs inspected	No. of organs inspected	No. of positive organs	Prevalence (%)	Total No of cyst on organ	cyst viability (%) per organs
Tongue	430	2	0.5	2	0(0/6)
Masseters muscle	430	2	0.5	2	16.7(1/6)
Heart	430	3	0.7	5	33.3(2/6)
Liver	430	4	0.9	7	50(3/6)
Lung	430	0	0	0	0
Kidney	430	0	0	0	0
Total	1720	11	0.64	16	100

**Table 3:** Indicates status of taeniasis in study area in relation to time period considered

	Frequency	Percent
Taeniasis		
Positive before 2000 E.C	133	32.1
Positive after 2000 E.C	28	6.7
Positive both before and after 2000 E.C	23	5.5
Total positive	184	44.3
Total respondents	415	100.0

*Table 4:* Shows the drug used by Taeniasis positive respondents

Taeniasis positive individuals		
Drug used	Frequency	Percent
Modern drug	163	88.6
Traditional drug	19	10.3
Non drug users	2	1.1

*Table 5:* Prevalence of human taeniasis with risk factors (address, age, sex and religion of respondents) in and around Asella, Tiyo woreda, Ethiopia

Variables	Categories	No of interviewees	No infected	Prevalence%	X <sup>2</sup>	P –value
Residential area	Rural	307	151	49.2	11.2	0.001
	Urban	108	33	30.6		
Age	<18	92	19	20.7	67.1	0.000
	18-35	199	74	37.2		
	>35	124	91	73.4		
Sex	Male	216	113	52.3	11.6	0.000
	Female	199	71	35.7		
Religion	Christian	228	109	47.8	2.4	0.07
	Muslim	187	75	40.1		
Educational status	Illiterate	173	103	59.5	28.9	0.000
	Elementary	153	52	34		
	High school	54	20	37		
Occupational status	College/above	35	9	25	55.2	0.000
	Student	128	25	19.5		
	Farmer	205	125	61		
	Civil workers	25	10	40		
Raw meat	Other private workers	57	24	42	54.8	0.000
	Consumers	211	131	62.1		
	Non-consumer	204	53	26		
	Total	415	184	44.3		





GLOBAL JOURNAL OF MEDICAL RESEARCH: G  
VETERINARY SCIENCE AND VETERINARY MEDICINE  
Volume 16 Issue 3 Version 1.0 Year 2016  
Type: Double Blind Peer Reviewed International Research Journal  
Publisher: Global Journals Inc. (USA)  
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

## Epidemiological Study of Small Ruminant Diseases in Selected Districts of Kaffa and Bench- Maji Zone, Southern Nations Nationalities and Peoples Regional State (SNNPRs), Ethiopia

By Fisseha Mengstie & Amenay Assefa

*Southern Agricultural Research Institute*

**Abstract-** An epidemiological survey of small ruminant disease was conducted in Boka-Shuta, Konda-Zuriya and Debre-Work kebelles of Kaffa and Bench-Maji zone, Southern Nations, Nationalities and people's regional state (SNNPRs), Ethiopia from July 2012 to June 2013. The objective of the study was to assess diseases of small ruminants affecting production and productivity; their occurrence and distribution; and associated risk factors. For these, Semi structured questioner format was prepared and survey was conducted. The result revealed that, 90.6%, 76.0% and 65.6% of interviewed farmers describe the multi-factorial causes of respiratory diseases; systemic diseases; and Gastro-intestinal parasite respectively were the most prevalent diseases irrespective of agro-ecology difference. The type of disease conditions documented in each agro-ecology had differences. Bloating (accumulation of excess amount of fluid in their abdominal cavity and blood tinged internal organs observed when slaughtered) followed by head circling in Boka-Shuta (highland); gastro intestinal parasite and head circling in Debre-Work (midland); and bottle jaw and orf in Konda-Zuriya (lowland) were the most prevalent and had serious problems in small ruminant production and productivity.

**Keywords:** *agro-ecology, diseases, epidemiology, production, small ruminant, susceptible.*

**GJMR-G Classification:** *NLMC Code: WC 900*



*Strictly as per the compliance and regulations of:*



© 2016. Fisseha Mengstie & Amenay Assefa. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License (<http://creativecommons.org/licenses/by-nc/3.0/>), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

# Epidemiological Study of Small Ruminant Diseases in Selected Districts of Kaffa and Bench- Maji Zone, Southern Nations Nationalities and Peoples Regional State (SNNPRs), Ethiopia

Fisseha Mengstie<sup>α</sup> & Amenay Assefa<sup>σ</sup>

**Abstract-** An epidemiological survey of small ruminant disease was conducted in Boka-Shuta, Konda-Zuriya and Debre-Work kebelles of Kaffa and Bench-Maji zone, Southern Nations, Nationalities and people's regional state (SNNPRs), Ethiopia from July 2012 to June 2013. The objective of the study was to assess diseases of small ruminants affecting production and productivity; their occurrence and distribution; and associated risk factors. For these, Semi structured questioner format was prepared and survey was conducted. The result revealed that, 90.6%, 76.0% and 65.6% of interviewed farmers describe the multi-factorial causes of respiratory diseases; systemic diseases; and Gastro-intestinal parasite respectively were the most prevalent diseases irrespective of agro-ecology difference. The type of disease conditions documented in each agro-ecology had differences. Bloating (accumulation of excess amount of fluid in their abdominal cavity and blood tinged internal organs observed when slaughtered) followed by head circling in Boka-Shuta (highland); gastro intestinal parasite and head circling in Debre-Work (midland); and bottle jaw and orf in Konda-Zuriya (lowland) were the most prevalent and had serious problems in small ruminant production and productivity. Regarding the diseases occurrence and distribution, the highest cases were recorded in summer followed by autumn. Ewes/doe and lambs/kids were the most susceptible age group. Each specific disease conditions occurrence and distribution in this study had its own specific risk factor. In general, however, disease management methods and practices by the farmers, susceptible age groups and seasons of occurrence were important risk factors playing an important role. Thus, the production and productivity of small ruminant was highly determined by diseases conditions. The result presented here suggests that, ewes/doe and lambs/kids need special attention to prevent them from diseases to improve production and productivity. And also, the small ruminant flock should get appropriate vaccination prior to the disease occurrence. Thus, the farmers' disease management and practices method need to be improved through training on small ruminant disease and their management. In addition, further studies should be conducted to identify the specific diseases types and their associated risk factors.

**Keywords:** agro-ecology, diseases, epidemiology, production, small ruminant, susceptible.

## I. INTRODUCTION

Indigenous small ruminants constitute greater percentage of ruminant population in Africa (Lebbie *et al.*, 1994). These flocks of animals are commonly found in the rural areas where they are owned and managed under extensive system (Otchere, 1986). Small ruminants play an important role in the lives of most people especially rural farmers who livelihood entirely depend on them. They provide source of animal protein through their meat and milk (Fajemisin, 1991). Not with standing, they fetch a source of income when sold to meet some other family needs as well as play a vital social roles during ceremonies and festivals.

The importance of small ruminants (ie sheep and goats) to the socio-economic well being of people in developing countries in the tropics in terms of nutrition, income and intangible benefits (eg savings, insurance against emergencies, cultural and ceremonial purposes) cannot be overemphasized (Kosgey, 2004). Sheep and goats are important livestock species in developing countries because of their ability to convert forages, and crop and household residues into meat, fibre, skin and milk.

For an improved animal protein intake, there is need for improvement in the production of meat and other protein sources from the livestock industry. Sheep and goats offer a great potential in this respect due to their relative ease of breeding, management, ability to subsist on forages, hardiness, adaptation to a wide range of ecological zones and distribution among others. In recent times, sheep and goats production is becoming popular even among urban dwellers as result of the aforementioned merits (Umunna *et al.*, 2014).

Small ruminant management is seriously hindered by diseases in the tropics. Diseases are very important to farmers and affect the production of small ruminants in several ways. It increases cost of

Author <sup>α</sup>: Southern Agricultural Research Institute (SARI), Bonga Agricultural Research Center (BARC). Bonga, Ethiopia.  
e-mail: fissehamengstie@yahoo.com

production, lowers production level, reduces the quality and quantity of animal products and generally causes great loss to the farmer (Abdullahi *et al.*, 2013).

In Ethiopia, There are more than 38 millions of cattle and 30 million small ruminants (CSA, 2007). However, the country is not making use of this huge potential attributed to different constraints among which disease stands in the front line (Samson and Frehwot, 2010; Firew, 1999). Diseases of various origins (bacterial, viral, parasitic, etc.) directly or indirectly are among the numerous factors responsible for poor production and productivity (Firew, 1999; Feyesa *et al.*, 2010) which results the major barriers for the improvement of livestock production, reproduction and marketing. The annual total economic losses due to diseases, mortality and reduced productive and reproductive performance were estimated by 150 million USD (Berhanu, 2002).

In Ethiopia, Sheep and goats contribute 25% of the meat domestically consumed with a production surplus mainly being exported as live animals (Alemayehu and Fletcher, 1991; Tibbo, 2006). Both species also contribute 50% of the domestic needs in wool, about 40% of skins and 92% of the value of hides and skin exported (ILCA, 1993). The total income share of small ruminants tends to be inversely related to size of land-holding, suggesting that small ruminants are of particular importance for landless people. In some settings where, agriculture (crop production) provides only seasonal employment, rearing small ruminants would provide employment and income as a subsidiary occupation (Coppock *et al.*, 2006).

However in Ethiopia, Small ruminant contribution to food consumption, rural income and export economy is below the expected potential, because of their husbandry is constrained by compound effect of disease, poor feeding, and poor management (Chalachew, 2001). Among many factors which limit the economic return from small ruminant production diseases stands in the front line (Firew, 1999). As a result of small ruminant diseases, 5-7 million sheep and goats die each year (Sileshi and Lidetu, 2007) and 35% of sheep and 56% of goat skins rejected due to skin parasites (Bayou, 1998) which leads serious economic loss to small holder farmers, the tanning industry, and the country as whole (ESGPIP, 2009) in which Many of the diseases in Ethiopia are still uncontrolled and are causing devastating effects both to the producers and to the national economy (EARO, 1996).

Even if small ruminants play a very important socio-economic role, little is known about health problems of these animals in the study area. In order to design relevant disease control strategies, assessment of the existing small ruminant diseases in the area is vital to devise appropriate technological interventions as well as for further study. Therefore, the aim of this study

was, to assess diseases of small ruminant affecting production and productivity; their occurrence and distribution; and associated risk factors in selected districts of Kaffa and Bench Maji zone.

## II. MATERIALS AND METHODS

### a) Study Area and Study Design

A cross-sectional study to assess diseases of small ruminant affecting production and productivity; their occurrence and distribution; and associated risk factors was conducted from July 2012 to June 2013 in Boka-Shuta and Konda-Zuriya of Kaffa zone; and Debre-work of Bench-maji zone. The study areas were selected purposively based on altitude variations, accessibility and availability of small ruminant. Thus, Boka-Shuta (highland), Konda-Zuriya (lowland) and Debre-work (Midland). The interviewed households with in each study areas were selected using systematical random sampling technique.

### b) Sample Size Determination and Data Collection

Before conducting the formal survey, group discussion was made with experts of each respective zonal and district agriculture and rural development office on small ruminant production problems and diseases in particular. Semi structured questioner format was prepared. Then, it was pre-tested and modified as necessary and finally the formal survey was conducted and data was collected using single-visit-multiple-subject formal survey technique (ILCA, 1990). Accordingly, a total of 96 households were interviewed and data's were collected on major small ruminant disease and health problems; their occurrence and distribution; and associated risk factors.

### c) Data Management and Analysis

The data was analyzed using STATA version 11. Descriptive statistics was employed to determine the prevalence and Chi-square ( $X^2$ ) test was used to measure the effect of predisposing factors on the distribution of different diseases. A significance level ( $p < 0.05$ ) and confidence level (95%) was set to determine the presence or absence of statistically significant difference between the given parameters. And also, multivariate logistic regression was tested to confirm the stated factors.

## III. RESULTS AND DISCUSSIONS

### a) Importance of Livestock Species

The overall importance of livestock species irrespective of agro-ecology difference indicated that, cattle had the highest proportion (90.6%) followed by poultry (68.8%) (Figure 1). The different animal species are distributed throughout the three agro-ecologies of the study areas; and these could suggest that, the areas have a potential to favorably support live stock

production and productivity. However, the proportion of livestock species had differences amongst the different agro-ecologies. Thus, small ruminants had the highest level of proportion than other livestock species; goat (54.5%) in Konda-Zuriya (lowland) followed by sheep (49.2%) in Boka-Shuta (highland). These could indicate that, the production potential for selected species of

animals in different agro-ecologies (figure 2). The present finding was supported by Enwelu et al (2015) from Nigeria; the farmer's preferred goat than sheep due to a number of reasons. These were; prolificacy (54.2%); profitability (36.1%); longevity (long life span) (5.6%); and type of meat/animal dung (2.8%).




Figure 1: Overall proportion of livestock species importance in Boka-Shuta, Debre-Work and Konda-Zuriya of southern Ethiopia




Figure 2: Proportion of livestock species importance in Boka-Shuta, Debre-Work and Konda-Zuriya of southern Ethiopia

b) *Small Ruminant Diseases and Their Prevalence*

Farmers listed different types of small ruminant diseases conditions that could cause mortality and morbidity in the study areas during the questioner survey. Among these, respiratory disease, systemic diseases, diarrhea, swollen head, Gastro-intestinal parasite and head circling had most frequently recorded. 90.6% of the respondents mentioned, diseases of the respiratory system (with signs of

coughing and nasal discharge) which could be multifactorial caused followed by 76.0% for systemic diseases (with signs of erected hair, depression and inappetent) were the most common. The multi symptomatic gastro-intestinal parasitism (in the present case could be manifested in diarrhea, emaciation or bloating) had significant occurrence (Figure 3).

Different research reports in Ethiopia indicated that, diseases are the main constraints limiting small

ruminant production. Among these, internal parasitic infestation followed by occurrence of persistent nasal discharge and coughing had the highest incidence limiting small ruminant production and could cause serious problems in kids and lambs (Urgessa *et al.*, 2012; Solomon *et al.*, 1995; Yohannes *et al.*, 1995; Solomon and Gemed, 2000; Markos, 2006).

The study report by Abebe *et al* (2013) in north western Ethiopia confirmed that, Sheep diseases were one of the main constraints for sheep production and Foot rot, skin disease, pasteurellosis, orf and internal parasites were the main. The authors extended their findings and; sheep diseases were the major constraints (ranked first) amongst several constraints limiting small ruminant production. Other authors from South Sudan revealed that; internal parasite followed by pneumonia and external parasite were the most prevalent small ruminant diseases (Lado *et al.*, 2015).

The occurrence of small ruminant diseases conditions with respect to agro-ecologies indicated that; bloating (accumulation of excess amount of fluid in their abdominal cavity and blood tinged internal organs

observed when slaughtered) followed by head circling were the most prevalent in Boka-Shuta. Whereas, gastro intestinal parasite and head circling in Debre-Work; and bottle jaw and orf in Konda-Zuriya were the most prevalent and had serious problems in the small ruminant production and productivity (Figure 4).

And also, statistical significant differences of diseases occurrence among the study areas were observed. Thus, systemic diseases followed by gastro-intestinal parasite and head circling had highest in Midland; whereas, Orf followed by tick infestation in Lowland. Lowest proportion of diseases occurrence was observed in Highland (Boka-Shuta) and these could be due to several intervention activities in the area by Bonga Agricultural research center for controlling and preventing diseases occurrence in small ruminant (like regular vaccination against ovine pasteurellosis disease, anthrax, PPR, sheep and goat pox disease; routine treatment of cases; and regular deworming with broad spectrum antihelmentics), since it was one of Bonga sheep breed improvement community/cooperative (Table 1).




Figure 3: Overall Prevalence of small ruminant disease conditions of Boka-Shuta, Debre-Work and Konda-Zuriya, southern Ethiopia




Figure 4: Prevalence of small ruminant disease conditions with respect to agro-ecologies, southern Ethiopia

Table 1: Prevalence of small ruminant disease conditions having statistical significant difference with respect to agro-ecologies, southern Ethiopia

Health related problems	N	Study areas			Percentage (%)	Sig.
		Boka-shuta (Highland)	Debre-Work (Midland)	Konda-Zuriya (Lowland)		
Respiratory disease	87	25(28.7)	33(37.9)	29(33.3)	90.6	0.074
Systemic disease	73	20(27.4)	32(43.8)	21(28.8)	76.0	0.002
Diarrhea	66	19(28.8)	24(36.4)	23(34.8)	68.8	0.817
Head swelling	64	23(35.9)	22(34.4)	19(29.7)	66.7	0.145
Internal parasite	63	12(19.0)	32(50.8)	19(30.2)	65.6	0.000
Head circling	57	24(42.1)	25(43.9)	8(14.0)	59.4	0.000
Bottle jaw	38	1(2.6)	7(18.4)	30(78.9)	39.6	0.000
Emaciation	32	8(25.0)	11(34.4)	13(40.6)	33.3	0.671
Bloating	14	7(50.0)	1(7.1)	6(42.9)	14.6	0.052
Orf	13	3(23.1)	0(0.0)	10(76.9)	13.5	0.002
Foot rot	12	5(41.7)	2(16.7)	5(41.7)	12.5	0.368



Eye disease	11	4(36.4)	1(9.1)	6(54.5)	11.5	0.153
Tick infestation	8	2(25.0)	0(0.0)	6(75.0)	8.3	0.031
Orchitis	8	0(0.0)	0(0.0)	8(100)	8.3	0.000
Mastitis	2	2(100)	0(0.0)	0(0.0)	2.1	0.094
Total	96	29(30.2)	33(34.4)	34(35.4)	100.0	

c) *Small Ruminant Diseases Management and Practices by the Farmers*

In general, small ruminant disease management by modern treatment in veterinary clinics was significantly different among the study areas. For example, 77.1% the interviewed household in Konda-Zuriya had better trend and awareness for treating their sick animals in veterinary clinics (Table 2). But, in areas where veterinary clinics are remote from farmers' residences (like in case of Boka-Shuta out of the reach

of intervention areas by the research center); the farmers manage their sick animals either by treating with home remedy and herbs or slaughtered for skin. Few households were also mentioned that, they used drugs by buying from the market and shop to treat their animals by themselves. The present finding was strongly supported by the study from South Sudan that; inadequate knowledge of small ruminant management was the most common challenge facing small ruminant production (Lado *et al.*, 2015).

Table 2: Small ruminant diseases management and practices by the farmers

Variables		Study areas			Total (%)	Sig.
		Boka-shuta	Debre Work	Konda-Zuriya		
Small ruminant disease management (controlling and prevention)	Self treatment using home remedy and herbs	14(28.6)	17(34.7)	18(36.7)	49(51.0)	0.932
	Modern treatment in veterinary clinics	17(23.0)	24(32.4)	33(44.6)	74(77.1)	0.001
	Slaughtering diseased sheep for skin	8(61.5)	2(15.4)	3(23.1)	13(13.3)	0.029
	Selling to market	11(26.2)	21(50.0)	10(23.8)	42(43.8)	0.148
	Vaccination against small ruminant diseases	14(40.0)	19(54.3)	2(5.7)	35(36.5)	0.000
Pasture management for disease transmission	Separate grazing/browsing house hold herd and flock from the community	14(23.3)	19(31.7)	27(45.0)	60(62.5)	0.030
	Communal grazing/browsing house hold herd and flock from the community	17(34.7)	17(34.7)	15(30.6)	49(51.0)	0.516
Quarantine system	Free movement of animal within and across the areas	27(40.3)	24(35.8)	16(23.9)	67(69.8)	0.000
	Restricted movement of animal within and across the areas	2(6.9)	9(31.0)	18(62.1)	29(30.2)	0.000
Total		29(30.2)	33(34.4)	34(35.4)	96(100)	



d) Temporal Distribution of Small Ruminant diseases

The occurrence of small ruminant diseases conditions in different seasons of the year had significantly varied. Thus, the highest occurrence was documented in summer (50.0%) followed by autumn (21.9%). On the other hand, seasons with high risk of disease occurrence were identified; and thus, autumn and summer with 32.3%. For these, different factors were identified. Grazing on immature fresh grass

(22.9%) was the main predisposing factor followed by unknown reason (20.8%) (Table 3). The present finding was justified by Abebe et al (2013); Sheep death occurs mainly at the end of the rainy season. This may be due to feed shortage and the suitability of the environment for the disease causing organisms. These also further aggravated by the introduction of animals into the area from different places and markets which might introduce different diseases.

Table 3: Temporal distribution of small ruminant disease conditions, southern Ethiopia

Variables		Study Areas			Total (%)	Sig.
		Boka-shuta	Debre-Work	Konda-Zuriya		
		N(%)	N(%)	N(%)		
Occurrence of diseases in different seasons	All season	3(75.0)	0(0.0)	1(25.0)	4(4.2)	0.000
	Autumn	5(23.8)	14(66.7)	2(9.5)	21(21.9)	
	Spring	2(66.7)	0(0.0)	1(33.3)	3(3.1)	
	Summer	5(10.4)	17(35.4)	26(54.2)	48(50.0)	
	Winter	14(70.0)	2(10.0)	4(20.0)	20(20.8)	
Seasons with high risk of disease occurrence	All season	3(75.0)	0(0.0)	1(25.0)	4(4.2)	0.000
	Autumn	6(19.4)	16(51.6)	9(29.0)	31(32.3)	
	Autumn & Summer	1(20.0)	4(80.0)	0(0.0)	5(5.2)	
	Spring	3(75.0)	0(0.0)	1(25.0)	4(4.2)	
	Summer	4(12.9)	7(22.6)	20(64.5)	31(32.3)	
	Winter	12(66.7)	3(16.7)	3(16.7)	18(18.8)	
	Winter & Autumn	0(0.0)	3(100)	0(0.0)	3(3.1)	
Factors for seasons of high risk of disease occurrence	Communal grazing	0(0.0)	1(100)	0(0.0)	1(1.0)	0.000
	Feed shortage	1(14.3)	6(85.7)	0(0.0)	7(7.3)	
	Feed shortage and unfavorable weather	9(52.9)	0(0.0)	8(47.1)	17(17.7)	
	Grazing on meshy areas	0(0.0)	0(0.0)	1(100)	1(1.0)	
	Immature fresh grass	6(27.3)	14(63.6)	2(9.1)	22(22.9)	
	Immature fresh grass and biting flies	0(0.0)	1(11.1)	8(88.9)	9(9.4)	
	Not known	3(15.0)	7(35.0)	10(50.0)	20(20.8)	

	Unfavorable weather & dusts	5(100)	0(0.0)	0(0.0)	5(5.2)	
	Unfavorable weather	5(35.7)	4(28.6)	5(35.7)	14(14.6)	
	Total	29(30.2)	33(34.4)	34(35.4)	96(100)	

e) Occurrence of Disease Outbreak in Small Ruminant flock

Significant variations for the occurrence of small ruminant disease outbreak in different seasons were recorded. Thus, 33.3% were during winter followed by

25% in autumn. Diseases manifesting respiratory signs had 50% of probability for causing outbreak (Table 4). The present finding was justified by Abebe et al (2013) report; disease mainly occurs during feed shortage periods (dry periods).

Table 4: Occurrence of small ruminant diseases outbreak, southern Ethiopia

Occurrence of small ruminant disease outbreak		Study Areas			Total	Sig,
		Boka-shuta	Debre-Work	Konda-Zuriya		
		N(%)	N(%)	N(%)		
Occurrence of disease outbreak in small ruminant		14(38.9)	4(11.1)	18(50.0)	36(37.5)	
Seasonal distribution	Autumn	3(33.3)	3(33.3)	3(33.3)	9(25.0)	0.035
	In any of the seasons	0(0.0)	0(0.0)	8(100)	8(22.2)	
	Spring	3(75.0)	0(0.0)	1(25.0)	4(11.1)	
	Summer	2(66.7)	0(0.0)	1(33.3)	3(8.3)	
	Winter	6(50.0)	1(8.3)	5(41.7)	12(33.3)	
Factors for diseases outbreak	Anthrax	1(50.0)	1(50.0)	0(0.0)	2(5.6)	0.027
	Bloating	2(100)	0(0.0)	0(0.0)	2(5.6)	
	Head circling	2(100)	0(0.0)	0(0.0)	2(5.6)	
	Diarrhea	0(0.0)	1(50.0)	1(50.0)	2(5.6)	
	Respiratory disease	2(11.1)	2(11.1)	14(77.8)	18(50.0)	
	Respiratory disease with diarrhea	1(100)	0(0.0)	0(0.0)	1(2.8)	
	Swollen head	2(100)	0(0.0)	0(0.0)	2(5.6)	
	Swollen head with coughing	0(0.0)	0(0.0)	2(100)	2(5.6)	
	Swollen head with coughing and diarrhea	0(0.0)	0(0.0)	1(100)	1(2.8)	
	Swollen head with diarrhea	1(100)	0(0.0)	0(0.0)	1(2.8)	
	Unknown disease	3(100)	0(0.0)	0(0.0)	3(8.3)	

f) *Distribution of Small Ruminant Diseases Conditions in Different Age Groups*

The occurrence and distribution of small ruminant diseases conditions in different age groups

were significantly vary even if, 22.9% were documented in all age groups. Thus, the highest was in ewes/doe and lambs/kids (26%) as compared to other age groups (Table 5).

**Table 5:** The occurrence and distribution of small ruminant disease conditions in different age groups, southern Ethiopia

Diseases distribution in different small ruminant age group		Study area			Total (%)	Sig.
		Boka-Shuta	Debre-Work	Konda-Zuriya		
		N (%)	N (%)	N (%)		
Age group	Adult	0(0.0)	1(100)	0(0.0)	1(1.0)	0.000
	Adults and lambs/kids	8(66.7)	2(16.7)	2(16.7)	12(12.5)	
	All age	6(27.3)	2(9.1)	14(63.6)	22(22.9)	
	Ewes/doe	7(46.7)	5(33.3)	3(20.3)	15(15.6)	
	Ewes/doe and lambs/kids	7(28.0)	5(20.0)	13(52.0)	25(26.0)	
	Lambs/kids	1(4.8)	18(85.7)	2(9.5)	21(21.9)	
Total		29(30.2)	33(34.4)	34(35.4)	96(100)	

g) *Factors Associated with the Occurrence and Distribution of Small Ruminant Diseases*

Based on multivariate logistic regression analysis, different determinant factors were found significantly associated for the occurrence and distribution of small ruminant disease conditions (Table 6). According to the model, holding the effect of other variables constant, the risk of acquiring infection with disease conditions manifested by swelling of head was 6.9% higher for those not treated in veterinary clinic (didn't get Modern treatment) than treated. The risk of getting head circling was 13.6% higher for Ewes/doe and lambs/kids than Lambs/kids alone; and season of occurrence was 39.6% higher in summer than winter. For GIT parasite infestation, the risk was 0.9% lower in Boka-Shuta than Konda-Zuriya which is justified by the regular deworming intervention by the research center in the study area. Traditional treatment practice by the farmers in the study areas for GIT parasite infestation indicated that, the risk was 21.2% higher for those not treated by herbal remedy than treated. For the occurrence and distribution of disease conditions like emaciations and eye diseases, market was significantly associated as a source of infection where diseased animals could disseminate the disease causing organism to the areas.

**Table 6:** Multivariate logistic regression analysis result - lists of significant variables after insignificant factors were excluded

Diseases conditions	Variables	Categories	OR	95% CI for OR	P value
Swollen head	Disease management method and practice	Modern treatment (No vs. Yes)	6.882	1.191, 39.8	.031
	Susceptible age	all ages vs. Lambs/kids	21.037	2.088, 211.9	.010
Head Circling	Study area	Konda-Zuriya vs. Boka-Shuta	.001	.000, .033	.000
	Susceptible age	Ewes/doe and lambs/kids vs. Lambs/kids	13.578	.950, 194	.055
	Season of occurrence	Summer vs. Winter	39.632	1.508, 1041.4	.027
GIT parasites	Study area	Boka-Shuta vs. Konda-Zuriya	.009	.000, .604	.028
	Disease management method and practice	Traditional treatment (No vs. Yes)	21.164	1.691, 264.8	.018
	Season of occurrence	summer vs. winter	.010	.000, .407	.015
Emaciation	Disease management method and practice	Selling to market	.129	.022, .763	.024
Bottle jaw	Study area	Boka-Shuta vs. Konda-Zuriya	.001	.000, .062	.001
		Debre-Work vs. Konda-Zuriya	.009	.000, .231	.005
Eye diseases	Disease management method and practice	Selling to market	.002	.000, .635	.035

#### IV. CONCLUSIONS AND RECOMMENDATIONS

The distribution and production of small ruminant was highly determined by diseases conditions. Of which, multi-factorial cause of respiratory diseases and Gastro-intestinal parasite are most prevalently occurred. The occurrence and distribution of those disease conditions had differences in seasons where the highest was in summer followed by autumn; and ewes/doe and lambs/kids were most frequently affected. Each specific disease conditions occurrence and distribution in this study had its own specific risk factor. In general, however, disease management methods and practices by the farmers, susceptible age groups and seasons of occurrence were important risk factors playing an important role. Thus, the production and productivity of small ruminant was highly determined by

diseases conditions. The result presented here suggests that, ewes/doe and lambs/kids need special attention to prevent them from diseases to improve production and productivity. And also, the small ruminant flock should get appropriate vaccination prior to the disease occurrence. In addition, the farmers' disease management and practices method need to be improved through training to create awareness on small ruminant disease and their management. In addition, further studies should be conducted to identify the specific diseases types and their associated risk factors.

#### V. ACKNOWLEDGMENTS

The author gratefully acknowledges Southern Agricultural Research Institute (SARI), Bonga Center, Ethiopia for financial and logistic support. I would like to

appreciate the research center livestock department researchers and technical assistance for their contribution for the success of the study. Also, I like to express my deepest gratitude to the informants for unreservedly sharing their valuable indigenous knowledge genuinely.

### REFERENCES RÉFÉRENCES REFERENCIAS

1. Abdullahi R., Awwal M. and Sawaneh M. (2013): Indigenous Treatment Methods of Small Ruminant Livestock in the Tropics –A Case Study from Katsina State, Nigeria. *IOSR Journal of Agriculture and Veterinary Science (IOSR-JAVS)*. Volume 6, Issue 1.
2. Alemayehu Z, Fletcher I: Small ruminant productivity in the central Ethiopia mixed farming systems. Addis Ababa: IAR (Institute of Agricultural Research); 1991.
3. Berhanu, A., 2002. Welcome address: Animal health and poverty reduction strategies. In: proceedings of the 16<sup>th</sup> Annual Conference of the Ethiopian Veterinary Association (EVA) held 5-6 June, 2002, Ghion Hotel, Addis Ababa, Ethiopia, pp: 117-137.
4. Chalachew N (2001). Study on skin diseases of cattle, sheep, and goat in and around wolayta sodd, Southern Ethiopia. DVM thesis, Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia.
5. Coppock, D. L., S. Desta, S. Tezerra and G. Gebru (2006) An Innovation System in the Rangelands: Using Collective Action to Diversify Livelihoods among Settled Pastoral Women in Ethiopia. Paper presented at Innovation Africa Symposium November 21-23, 2006 held in Kampala, Uganda.
6. CSA (2007). Central statistical authority, Ethiopia agricultural sample enumeration, statistical report on livestock population, part 4, Addis Ababa, Ethiopia.
7. Daba Urgessa, Belay Duguma, Solomon Demeke and Taye Tolamariam. A Survey of Sheep and Goat Diseases in Ilu Abba Bora Zone of Oromia Regional State, Southwestern Ethiopia. *Global Veterinaria* 9 (5): 552-556, 2012.
8. EARO, 1996. Animal Health Research program strategy (draft document). Ethiopian Agricultural Research Organization, Addis Ababa, Ethiopia.
9. ESGPIP (2009). Ethiopian sheep and goat productivity improvement program. Common defects of Sheep and Goat skin in Ethiopia and their causes. *Tech. Bull.* p. 19.
10. Feyesa R., N. Misrak and Ts.Tefera. Sero-Prevalence of Contagious Caprine Pleuropneumonia in Goat at Selected Woredas of Afar Region, *Ethiopian veterinary journal*, 2010; 14(1), 83-89.
11. Firew, T., 1999. Livestock Production, Management and Utilization. Mekelle University, Mekelle, Ethiopia.
12. I.A. Enwelu, E.L. Ezeuko, and N.S. Machebe (2015). Challenges of smallholder sheep and goat keeping in rural communities of Aguata Agricultural Zone of Anambra State, Nigeria. *Indian J. Anim. Res.*, 49 (3). 373-377
13. ILCA (International Livestock Center for Africa): *Handbook of African Livestock Statistics*. Addis Ababa; 1993.
14. Kosgey IS, Breeding objectives and breeding strategies for small ruminants in the tropics. PhD Thesis, Wageningen University, The Netherlands, p. 272, 2004.
15. Lado, M.M.\*, Salah K. Jubarah and Erneo B. Ochi (2015). A Case Study on Major Constraints of Small Ruminants Management in Juba County Central Equatoria State South Sudan. *International Journal of Innovative Science, Engineering & Technology*, Vol. 2 Issue 12.
16. Samson L. and M. Frehwot, Prevalence of Small Ruminant Trypanosomosis and Tsetse Fly Challenge in Upper Didessa Valley, Ethiopia, *Global Veterinaria* 2010; 5 (4): 215-219.
17. Sileshi, Z. and D. Lidetu, 2007. Control of internal parasites in sheep and goats. In: technical bulletin no.3 Control of internal parasites in sheep and goats (Eds.) L. Dawson, R.C. Merkel and Alemu Yami. [www.esgPIP.org/PDF/Technical%20bulletin%20No.3.pdf](http://www.esgPIP.org/PDF/Technical%20bulletin%20No.3.pdf)
18. Tibbo M: Productivity and Health of Indegenous Sheep breeds and crossbreds in the Central Ethiopian Highlands. In *PhD Thesis*. Swedish University of Agricultural Sciences, Uppsala; 2006.
19. Umunna M. O., Olafadehan O. A. and Arowona A. (2014): Small Ruminant Production and Management Systems in Urban Area of Southern Guinea Savanna of Nigeria. *Asian Journal of Agriculture and Food Science*. Volume 02 – Issue 02.
20. Yenesew Abebe, Solomon Melaku, Azage Tegegne and Firew Tegegne (2013). Assessment of Sheep Production System In Burie District, North Western Ethiopia. *Global Journal of Agricultural Research* Vol.1, No.2, pp. 29-47.

This page is intentionally left blank





GLOBAL JOURNAL OF MEDICAL RESEARCH: G  
VETERINARY SCIENCE AND VETERINARY MEDICINE  
Volume 16 Issue 3 Version 1.0 Year 2016  
Type: Double Blind Peer Reviewed International Research Journal  
Publisher: Global Journals Inc. (USA)  
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

## Major Causes of Liver Condemnation and Associated Direct Financial Losses in Bovine Slaughtered at Assela Municipal Abattoir Arsi, South Eastern Ethiopia

By Hussein Aman, Seifudin Kassim, Gobu Boru, Hubado Hussien  
& Mukarim Abdurahaman

*Jimma University*

**Abstract-** A study was conducted from October, 2013 to March, 2014 on cattle slaughtered at Assela municipal abattoir with the aim of determining major causes of liver condemnation and to estimate the direct financial losses attributed to the condemned liver. Ante mortem and post mortem inspection procedures were followed throughout the study and abnormalities encountered were recorded. A total of 384 cattle were examined at ante mortem and 9 animals were found to have abnormalities/conditions like lameness, blindness, rough hair, amputated tail and branding. Out of 384 cattle slaughtered, 274 (71.4%) livers were totally condemned. The major causes of liver condemnation were due to hydatidosis (64.6%), fasciolosis (20.8%), calcification (7.7%), coinfection of hydatid cyst with fasciola (4.7%), cirrhosis (1.5%) and abscess (0.7%). Comparison of liver condemnation rate was carried out for different age, animal origin and body condition however, statistically significant difference was observed only between age groups.

**Keywords:** *abattoir, assela, cattle, condemnation, financial loss, liver.*

**GJMR-G Classification:** *NLMC Code: WI 141*



MAJOR CAUSES OF LIVER CONDEMNATION AND ASSOCIATED DIRECT FINANCIAL LOSSES IN BOVINE SLAUGHTERED AT ASSELA MUNICIPAL ABATTOIR ARSI SOUTH EASTERN ETHIOPIA

*Strictly as per the compliance and regulations of:*



© 2016. Hussein Aman, Seifudin Kassim, Gobu Boru, Hubado Hussien & Mukarim Abdurahaman. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License (<http://creativecommons.org/licenses/by-nc/3.0/>), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.



# Major Causes of Liver Condemnation and Associated Direct Financial Losses in Bovine Slaughtered at Assela Municipal Abattoir Arsi, South Eastern Ethiopia

Hussein Aman<sup>α</sup>, Seifudin Kassim<sup>σ</sup>, Gobu Boru<sup>ρ</sup>, Hubado Hussien<sup>ω</sup> & Mukarim Abdurahaman<sup>¥</sup>

**Abstract-** A study was conducted from October, 2013 to March, 2014 on cattle slaughtered at Assela municipal abattoir with the aim of determining major causes of liver condemnation and to estimate the direct financial losses attributed to the condemned liver. Ante mortem and post mortem inspection procedures were followed throughout the study and abnormalities encountered were recorded. A total of 384 cattle were examined at ante mortem and 9 animals were found to have abnormalities/conditions like lameness, blindness, rough hair, amputated tail and branding. Out of 384 cattle slaughtered, 274 (71.4%) livers were totally condemned. The major causes of liver condemnation were due to hydatidosis (64.6%), fasciolosis (20.8%), calcification (7.7%), coinfection of hydatid cyst with fasciola (4.7%), cirrhosis (1.5%) and abscess (0.7%). Comparison of liver condemnation rate was carried out for different age, animal origin and body condition however, statistically significant difference was observed only between age groups. The direct financial loss due to liver condemnation was calculated to be 418,761 ETB per annum, rendering them unfit for local market for eusthetic reason or considering their zoonotic risks to human beings. To this end, public education on the effects of animal disease, application of strict control measures accompanying epidemiological studies and exercising appropriate meat inspection procedures were forwarded to minimize liver condemnation and as well as to safeguard the welfare of the public.

**Keywords:** abattoir, assela, cattle, condemnation, financial loss, liver.

## I. INTRODUCTION

One of the losses from endemic disease is expressed in terms of organ condemnation. The most commonly affected organs are liver and lung due to fasciolosis (for liver) and hydatidosis (for both) (Teka, 1997). Each year significant loss results from death of animals, inferior weight gain, condemnation of edible organs and carcasses at slaughtered. This production loss to the livestock industry is estimated at more than 900 million USD annually (Abebe, 1995).

**Author α ¥:** Jimma University, College of Agriculture and Veterinary Medicine (JUCAVM). e-mails: hussaman@yahoo.com, gobuboru@yahoo.com, mukevet@yahoo.com

**Author σ ω:** Bale zone and Chole wereda Livestock and fisheries resource development office. e-mails: seifudinkassim@gmail.com, kedjajak@gmail.com

A through meat inspection procedure requires two-steps namely antemortem and postmortem inspection. The importance of ante mortem inspection in the abattoir has long been recognized in an attempt to avoid the introduction of clinically diseased animals into the slaughterhouse. Ante mortem inspection should be done within 24 hours of slaughter and repeated when slaughter has been delayed over a day (Teka, 1997; Mezgebu, 2003).

Postmortem inspection is screening or sorting process to separate the normal from abnormal. It is the center around which meat hygiene revolves since it provides information indispensable from the scientific evolution of clinical signs and pathological process that affect wholesomeness of meat. Routine post mortem inspection of carcass and organs should be carried out as soon as possible after completion of dressing (Gracey, 1986).

The final judgment to be taken with an organ and carcass or parts of a carcass is based on the total evidence produced by observation, palpation, incision, smell, and any ante mortem signs (Teka, 1997). It is necessary to be aware the extent to which the public is exposed to certain zoonotic diseases detected in abattoirs and financial losses attributed to condemnation of affected organs and carcass (Nfi and Alonge, 1987). This is due to the fact that, meat is the main source of particular importance to the public such as hydatidosis, fasciolosis, tuberculosis and cysticercosis (Sirak, 1991).

Bovine liver is one of the largest visceral organs in the animal body which performs numerous functions and very rich sources of vitamins and minerals (Radostitis *et al.*, 2007). The tissue is much sought by consumers due to its palatability and easy to consumption. However, it is one of the most commonly condemned visceral organs during routine meat inspection (Phiri, 2006).

Parasites in the tropics are responsible for greater losses to the meat industry than any other diseases (Jobre *et al.*, 1996). Similarly like many other tropical countries of Africa, it is well known that parasitic diseases are among the major factors responsible for

the low productivity of livestock in Ethiopia (Abebe, 1995; Jobre *et al.*, 1996).

Cystic echinococcosis/Hydatidosis is a parasitic infection caused by larval stage of *Echinococcus granulosus*, which is small tapeworm, for which dogs and other canids are typical definitive hosts. The adult parasite found in small intestine of carnivours while the metacestode (hydatid cyst) is found in different organs of a wide varieties of herbivours including (sheep, goats and cattle), pig, horse and man (Soulsby, 1982).

Fasciolosis in cattle is chronic wasting disease caused by the presence in the liver and bile ducts respectively of immature and adult Trematode of the genus *fasciola*. The disease is found in vast areas of the world with the smaller *fasciola hepatica* (3.5x1cm) in temperate countries and the large *fasciola gigantica* (7.5cm) in tropical regions (Andrews *et al.*, 1999).

Various researchers have undertaken studies at abattoir surveys to determine the prevalence and economic importance and cause of meat condemnation (hydatidosis, Fasciolosis, Cysticercosis) as these are mainly to be of major economic and public health importance in meat inspection (Jobre *et al.*, 1996). Therefore, the objective of this study was to identify the major causes of liver condemnation in cattle slaughtered in Assela municipal abattoir and to estimate the magnitude of direct economic losses attributed to this condemnation.

## II. MATERIALS AND METHODS

### a) Study area

The study was conducted at Assela municipal abattoir, from October 2013 to march 2014. Assela is located 175Km South East of Addis Ababa at an altitude of 2350-2400 meters above sea level and has a climatic condition of "Woynadega". The annual average rainfall is 1300-1350mm. A day and night temperature of the area ranges from 10-25°C and 10-20°C respectively. The area has a bimodal rainfall occurring from March to April (short rainy season) and from July to October (long rainy season) with mean annual rain fall of 1300-1350mm with the relative humidity of 43-60°C (CSA, 2009).

### b) Study animals

A total of 384 randomly selected cattle were inspected at Assela municipal abattoir. Out of which, 373 (97.14%) were males and only 11 (2.86%) were females. From the total of 384 cattle slaughtered, 348 were old and 36 were adults. Majorities of cattle came to Assela municipal abattoir for slaughters were originated from the market places of Sagure, Kersa and Assela.

### c) Study Design

A cross-sectional study was employed to identify the major cause of liver condemnation and to evaluate the direct financial losses. The study animals were selected using simple random sampling method

by taking the age, body condition and origin of the animals into consideration. The desired sample size for this study was calculated by using the formula given by Thrusfield (1995) with 95% confidence interval, 5% absolute precision and 50% expected prevalence

### d) Study Methodology

#### i. Ante-Mortem Examination

Ante-mortem examination was conducted on individual animals, while the animals were entering in to the lairage and in mass after they entered into the lairage. Both sides of the animals were inspected at rest and in motion. Moreover, the general behavior of the animals, sign of diseases and abnormality of any type were recorded according to the standard antemortem inspection procedures (Gracey, 1986). Following the judgment guideline by (FAO, 2003), animals fit for human consumption were allowed for slaughter.

#### ii. Post-mortem Examination

Post-mortem examination involved visual inspection, palpation and making systemic incision of liver to look for the presences of cysts, adult parasites and other abnormalities. Pathological lesions were differentiated and judged based on (FAO, 2003) guidelines on meat inspection for developing countries.

#### iii. Financial Loss Assessment

The direct financial losses due to liver condemnation from market were considered. The analysis was based on the annual slaughter capacity of the abattoir considering market demand, the current average price of one liver in Assela Butcherhouse and the rejection rate of liver. The direct financial loss incurred due to liver condemnation was estimated by using the formula indicated below (Ogunrinade and Adesoke, 1982).

$$ALC = CSR \times LC \times P$$

Whereas

ALC = Annual loss from liver condemnation

CSR = Mean annual cattle slaughtered at Assela municipal abattoir

LC = Mean cost of one liver at Assela

P = Liver condemnation rate

### e) Statistical Data Analysis

Data collected during the study were entered in to Excel spread sheet (Microsoft Excel 2007) and analyzed by statistical methods using SPSS 16.0 version. Descriptive statistics such as percentage was used to determine the level of liver condemnation rate. The association between condemnation rate of liver and the age, sex, body condition and origin of the animal assessed by Pearson chi-square ( $\chi^2$ ) and the *p*-value < 0.05 were considered significant.

### III. RESULTS

#### a) Ante-Mortem Examination

Of the total 384 cattle examined at ante-mortem, 9 cattle were found to have the abnormalities

listed below (Table 1). Some abnormalities encountered during antemortem inspection were rough hair coat, lameness, branding on skin, blindness and amputated tail.

**Table 1:** Abnormalities encountered during ante mortem examination

Abnormalities	No. of animals with disease condition	Judgment
Lameness	2	Passed for slaughter but with precaution
Blindness	1	Passed for slaughter but with precaution
Rough hair coat	4	Passed for slaughter but with precaution
Branding on skin	1	Passed for slaughter but with precaution
Amputated tail	1	Passed for slaughter but with precaution
Total	9	Passed for slaughter

#### b) Post-Mortem Examination

Out of 384 cattle slaughtered in Assela municipal abattoir, 274 livers were totally condemned due to various reasons (Table 2). The current study revealed the overall proportion of liver condemnation

rate due to various pathological findings was 71.4%. The occurrence of hydatidosis was the highest (64.6%); followed by fasciolosis (20.8%), calcification (7.7%), coinfection of hydatid cyst with fasciola (4.7%), cirrhosis (1.5%) and liver abscess (0.7%).

**Table 2:** Major causes of liver condemnation, Frequency and its proportion

Major cause of liver condemnation	Frequency	proportion
Hydatid cyst	177	64.6
Fasciolosis	57	20.8
Calcification	21	7.7
Coinfection of hydatid cyst with fasciola	13	4.7
Cirrhosis	4	1.5
Abscess	2	0.7
Total	274	100

Comparison of rejection rate of liver was carried out for different age, animal origin and sex groups. From the total of 384 sampled animals, 340 were from sagure while only 44 animals were from (Assela and near surrounding). The proportion of liver condemnation was 71.8% and 68.2% from sagure and (Assela & near surrounding) respectively. However, no statistically significance difference was observed in the liver condemnation rate between the two areas.

statistically significant difference was observed between the two sexes (Table 3). This study also showed that a highest liver condemnation rate in older age groups (> 5 years) than adults (< or = 5 years). The rejection rate was 73.3% in old and 52.8% in adult (Table 3). From the analysis, it was observed that as age increase, the rejection rate was also found to increase. This difference in the rejection rate between the age groups was significant (Table 3).

Regarding sex, the overall liver condemnation rate was 71.8% in male and 54.5% in female and no

**Table 3:** Liver condemnation rate based on Age, Sex, Animal origin and body condition

Variables		No. exam	No. (+)	%	$\chi^2$	p-value
Origin	Sagure	340	244	71.8	0.245	0.621
	A&S	44	30	68.2		
Sex	Male	373	268	71.8	1.565	0.211
	Female	11	6	54.5		
Age	old	348	255	73.3	6.707	0.010
	Young	36	19	52.8		
BCS	Good	250	184	73.6	1.768	0.184
	Medium	134	90	67.2		

A&S = Assela and near surrounding

No. exam = number of animals examined

No. (+) = number of positive animals

BCS = body condition score

c) *Annual Financial Losses Estimation*

The annual slaughter rate of the abattoir for the last three years were 6800, 6900 and 6998 in 2010, 2011 and 2012, respectively. So the direct annual financial loss due to rejection of liver was calculated based on

the price of a liver at Assela (Table 4). By using necessary information and formula, the annual direct financial loss incurred due to condemnation of liver was calculated to be 418,761 ETB or 22,040 US\$ per annum.

Table 4: Average annual estimated financial loss due to liver condemnation

Annual slaughter rate	Cause of liver condemnation	Annual rejection rate of liver	Average price of one liver
6900	Hydatid cyst	3181	85 ETB
	Fasciola	1024	
	Calcification	377	
	Hydatid cyst with fasciola	234	
	Cirrhosis	72	
	Abscess	36	

#### IV. DISCUSSION

In the present study, routine ante-mortem and post-mortem inspection was carried out to detect any abnormalities encountered in Assela municipal abattoir. Branding, lameness, rough hair, blindness and amputated tail were found in some animals during ante mortem examination and animals with such abnormalities were passed for slaughter by considering that the problems were localized and simply related to (rough or low management system) and not due to bad pathological condition.

From the total of 384 cattle slaughtered, 274 (71.4%) liver were condemned due to various causes. This finding is in agreement with one of previous study, where 66.55% liver was condemned in Kombolcha ELFORA industrial abattoir (Nurit *et al.*, 2012). But slightly higher than 53.7% that have been done in Kombolcha (Jemal, 2009) and 51.95% in Mekelle (Shegaw *et al.*, 2009). On the other hand, it is significantly higher than 31.1% and 17.61% reported by (Yifat *et al.*, 2011) in Gonder and (Mellau *et al.*, 2011) in Tanzania respectively.

In the present study, the major cause of liver condemnation were hydatidosis (64.6%) followed by fasciolosis (20.8%). In the present study, the rejection rate of liver due to hydatidosis is 64.6%, which is highly greater than the findings of (Nurit *et al.*, 2012) with 14.2% and (Jemal, 2009) with 9.2% in both are at Kombolcha ELFORA industrial abattoir, 4.2% in Tanzania (Mellau *et al.*, 2011) and (Yifat *et al.*, 2011) with 3.7% in Gondar.

The rejection rate due to faciolosis is high when it is compared with the rejection rate of 12.7, 14.05 and 8.6% by (Fufa *et al.*, 2009) at Welaita Sodo, (Swai and Ulicky, 2009) at Hawi and (Mellau *et al.*, 2011) at Tanzania, respectively. On the other hand it was slightly similar with 24.32% by (Gebretsadik *et al.*, 2009) at Mekelle, but significantly lower than 68.7% in Kombolcha (Jemal, 2009) and 86.4 % in Gondar (Yifat

*et al.*, 2011). This may be due to climate and ecological condition of the study areas considered.

The finding of 1.5% of liver condemnation due to cirrhosis was significantly lower than 16.06% reported by (Nurit *et al.*, 2012) at ELFORA and (Raji, *et al.*, 2010) with 10.4% at Zaria abattoir. However, it is almost similar with 1.1% reported by (Yifat *et al.*, 2011) at Gondar.

A 0.7% liver condemnation due to abscess was recorded in the present study. This is a little bit smaller than the report by (Cadmus and Adesokan, 2009) with 2.9% condemnation rate of liver due to hepatic abscess in western Nageria and (Ahmedullah *et al.*, 2007), who reported 3.8% liver condemnation rate in Bangladesh. On the other hand it was similar to the report of (Mellau and Nongaond, 2010) with 1.1% at Arusha abattoir.

Calcifications were also among the lesions which significantly contributed to the liver condemnation in the study abattoir. During this study, 7.7% of liver were condemned as a result of calcification, which is relatively higher than the finding of (Mellau and Nongaond, 2010) which is 1.9% at Arusha, Tanzania. On the other hand, it is similar with the finding of (Nurit *et al.*, 2012) with 8.18% done at ELFORA.

The analysis of the result on the bases of age indicated the total liver rejection rate was higher in older animals and a significant difference was observed between the two age groups. This may be due to most of liver diseases are chronic and the older animals are mostly affected by many diseases.

The direct financial loss incurred as result of condemnation of liver in the present study was 418,761 ETB or 22040 US\$ per annum. This is so much higher than 1800 US\$ per annum due to fasciolosis liver condemnation reported by (Mwabonimana, 2008) at Arusha abattoir, Tanzania.

#### V. CONCLUSION

In general, liver condemnations as have been reported in this study impact negatively on the economic status of the traders and the livestock industry at large. This constituted a substantial loss to the economy of the

slaughter stock owners under study as such an amount of money would have been harnessed into livelihood improvements. Though infected livers were condemned and rendered unfit for human consumption, there exist some public health threats from animals slaughtered at the abattoir due to the possibility of some missed cases as a result of poor cooperation between butchers and meat inspectors and other malpractices including hiding of infected meat from meat inspectors to avoid economic losses on their side. Indeed, the condemnation of cattle livers at slaughterhouse in Assela municipal abattoir represents a significant economic loss. Some of the conditions described however can be prevented. Cases of hydatidosis could be reduced by better control of stray dogs. Since most liver conditions were caused by parasites, deworming programmes coupled with good animal husbandry would likely be effective in lowering their incidence. Some of the limitations, however; encountered in this study included the use of only gross pathology in the diagnosis of the diseases, thus only those diseases with gross pathological lesions that are pathognomonic were likely to be diagnosed. In spite of the limitation mentioned, the public health implications of the quantity of infected livers condemned at Assela municipal abattoir on the consumers and the role which post-mortem inspection plays in safeguarding the health of the public cannot be overemphasized. Therefore, there is a need for adequate meat inspection in Assela municipal abattoir in order to reduce wastages, identify diseases and thereby minimize associated public health risks.

## VI. ACKNOWLEDGMENTS

The authors acknowledge Jimma University College of Agriculture and Veterinary Medicine for supporting the study

## REFERENCES RÉFÉRENCES REFERENCIAS

1. Abebe, G. (1995): Current status of vet. Education and Animal Health research in Ethiopia, in vet. Edu. Impact of human health and nutrition in Africa. Proceeding of an international IRLI, AA, and Pp 133-138.
2. Ahmedullah, F.M., M.G. Akbor, M.M. Kaider and K. Hossain, (2007): Pathological investigation of liver of the slaughtered buffaloes in Barisal district, Bangladesh. *J. Vet. Med.* **5**: 81-85.
3. Andrews, S.J. (1999): The life cycle of hepatica in fasciolosis Dalton, J.P.Ed. CABI publishing walling ford, UK, Pp: 1-30.
4. CSA, (2009): Central Statistical Authority livestock number of breed, age, sex and purpose. In: report on livestock Characteristics (private peasant holding) *statistical bulletin-vol II*, Addis Ababa.

5. FAO, (2003): Manual Diagnosis on Meat Inspection for Developing Country, FAO/UN. *Viale deile termai caralla*. Rome, Italy. Pp: 49.
6. Fufa, A., A. Loma, M. Bekele and R. Alemayehu, (2009). Bovine fasciolosis: coprological, abattoir survey and its economic impact due to liver condemnation at Soddo municipal abattoir, Southern Ethiopia. *Tropical Animal Health and Production.* **42**(2): 289-292.
7. Gebretsadik, B., B. Kassahun and T. Gebrehiwot, (2009): Prevalence and economic significance of fasciolosis in cattle in Mekelle Area of Ethiopia. *Tropical Animal Health and Production.* **41**(7): 1503-1504
8. Gracey, J.F. (1986): Meat hygiene. 8<sup>th</sup> ed. Baillere Tindal. London, Philadel Phia. Toronto. Pp: 223-260.
9. Jemal, T. (2009): Major causes of organ condemnation in cattle slaughter at Kombolcha ELFORA meat factory abattoir south Wollo, DVM thesis, Haramaya University, Ethiopia.
10. Jobre, Y., Labago, F., Tiruneh, R., Abebe, G. and Dorchies, PhD. (1996): Hydatidosis In three selected region in Ethiopia an assessment trial on its prevalence, Economic and public health importance. *Revne de medicine veterinary*, **147**(1): 797-804
11. Mellau, L.S. and E.D., Nongaond. (2010): Slaughter house survey liver lesions in slaughtered cattle, sheep and goats at Arasha, Tanzania. *Journal of Vet Sci*, **3**: 179-188
12. Mellau, B.L., H.E. Nonga and E.D. Karimuribo, (2011): Slaughter stock abattoir survey of carcasses and organ/offal condemnations in Arusha Region, northern Tanzania. *Tropical Animal Health and Production*, **43**: 857-864.
13. Mezgebu, Y. (2003): Major cause of organ condemnation in ruminants slaughtered at Gondar Abattoir, North western Ethiopia, DVM thesis, FVM, AAU, Debre-Zeit, Ethiopia, Pp17.
14. Mwabonimana, M.F. (2008): Cattle liver condemnation at Arusha meat company Ltd, Tanzania: causes and its financial implication. A Research Paper Submitted in Partial Fulfilment of the Requirements for the Degree of Master of Preventive Veterinary Medicine of Sokoine University of Agriculture, Morogoro, Tanzania. Pp: 10-30.
15. Nfi, A.N and Alonge, D.O. (1987): An economic survey of abattoir data in Fako division of south west, Cameroon, *Bulletin Animal Health and Production in Africa.* **35**(3): 239-242.
16. Nurit, M., Zerihun. H and Serkalem. M. (2012): Major Cause of Liver Condemnation and Associated Financial Loss at Kombolcha Elfora Abattoir, South Wollo, Ethiopia, *European Journal of Applied Sciences.* **4** (4): 140-145.
17. Ogunrinade, A. and G.O. Adesoke, (1982): Bovine fasciolosis in Nigeria, intercurrent parasitic and

- bacterial infection. *Trop. Alm. Health Prod*, **14**: 121-124.
18. Phiri, A.M., (2006): Common conditions leading to cattle carcass and offal condemnation at south west province, Cameroon. *Bull. Anium. Hlth. Prod. Afr.* **35**(3): 239-242.
  19. Radostits, O.M., Gay, C.C., Hinchcliff, K.W. and Constable, P.D. (2007): *Veterinary Medicine: A Textbook of the Diseases of Cattle, Horses, Sheep, Pigs and Goats*, 10<sup>th</sup> ed, Elsevier Health Sciences, Philadelphia, PA, USA, Pp: 1498-1506.
  20. Raji, M.A., S.O. Solam and J.A. Ameh, 2010. Pathological condition and lesions observed in slaughtered cattle at Zaria abattoir. *J. clinical pathology and Forensic Medicine*, **1**: 9-12.
  21. Sirak, A. (1991): Cause of organ condemnation at Bahir Dar abattoir. Proceeding of the fourth national livestock improvement Conference, Institute of Agricultural Research, Addis Ababa, Ethiopia, Pp: 25-36.
  22. Shegaw, S., K. Ashwani and A. Kassaw, (2009): Organ condemnation and economic loss at Mekelle municipal abattoir, Ethiopia. *Global veterinarian*, **48**: 17-22.
  23. Soulsby E.J.L. (1982): *Helminths, arthropods and Protozoa of domesticated animals*, 7<sup>th</sup> ed. Birilliere Tindall, London, UK, Pp: 40-52.
  24. Swai, E.S. and E. Ulicky, (2009): An evaluation of the economic losses resulting from condemnation of cattle livers and loss of carcass weight due to fasciolosis: A case study from Hai town abattoir, Kilimanjaro region, Tanzania. *Livestock Res. Rural Dev.*, **21**(11)
  25. Teka, G. (1997): *Meat Hygiene and Food Hygiene principles and methods of food borne disease control with special reference to Ethiopia* Pp: 99-133.
  26. Yifat, D., D. Gedefaw and S. Desie, (2011): Major cause of organ condemnation and financial significance of cattle slaughtered at Gondar ELLFORA abattoir. *Global Veterinarian* **7**(5): 487-490.

# GLOBAL JOURNALS INC. (US) GUIDELINES HANDBOOK 2016

---

[WWW.GLOBALJOURNALS.ORG](http://WWW.GLOBALJOURNALS.ORG)

# FELLOWS

## FELLOW OF ASSOCIATION OF RESEARCH SOCIETY IN MEDICAL (FARSM)

Global Journals Incorporate (USA) is accredited by Open Association of Research Society (OARS), U.S.A and in turn, awards "FARSM" title to individuals. The 'FARSM' title is accorded to a selected professional after the approval of the Editor-in-Chief/Editorial Board Members/Dean.



- The "FARSM" is a dignified title which is accorded to a person's name viz. Dr. John E. Hall, Ph.D., FARSS or William Walldroff, M.S., FARSM.

FARSM accrediting is an honor. It authenticates your research activities. After recognition as FARSM, you can add 'FARSM' title with your name as you use this recognition as additional suffix to your status. This will definitely enhance and add more value and reputation to your name. You may use it on your professional Counseling Materials such as CV, Resume, and Visiting Card etc.

*The following benefits can be availed by you only for next three years from the date of certification:*



FARSM designated members are entitled to avail a 40% discount while publishing their research papers (of a single author) with Global Journals Incorporation (USA), if the same is accepted by Editorial Board/Peer Reviewers. If you are a main author or co-author in case of multiple authors, you will be entitled to avail discount of 10%.

Once FARSM title is accorded, the Fellow is authorized to organize a symposium/seminar/conference on behalf of Global Journal Incorporation (USA). The Fellow can also participate in conference/seminar/symposium organized by another institution as representative of Global Journal. In both the cases, it is mandatory for him to discuss with us and obtain our consent.



You may join as member of the Editorial Board of Global Journals Incorporation (USA) after successful completion of three years as Fellow and as Peer Reviewer. In addition, it is also desirable that you should organize seminar/symposium/conference at least once.

We shall provide you intimation regarding launching of e-version of journal of your stream time to time. This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.








The FARSM can go through standards of OARS. You can also play vital role if you have any suggestions so that proper amendment can take place to improve the same for the benefit of entire research community.

As FARSM, you will be given a renowned, secure and free professional email address with 100 GB of space e.g. [johnhall@globaljournals.org](mailto:johnhall@globaljournals.org). This will include Webmail, Spam Assassin, Email Forwarders, Auto-Responders, Email Delivery Route tracing, etc.



The FARSM will be eligible for a free application of standardization of their researches. Standardization of research will be subject to acceptability within stipulated norms as the next step after publishing in a journal. We shall depute a team of specialized research professionals who will render their services for elevating your researches to next higher level, which is worldwide open standardization.

The FARSM member can apply for grading and certification of standards of their educational and Institutional Degrees to Open Association of Research, Society U.S.A. Once you are designated as FARSM, you may send us a scanned copy of all of your credentials. OARS will verify, grade and certify them. This will be based on your academic records, quality of research papers published by you, and some more criteria. After certification of all your credentials by OARS, they will be published on your Fellow Profile link on website <https://associationofresearch.org> which will be helpful to upgrade the dignity.



The FARSM members can avail the benefits of free research podcasting in Global Research Radio with their research documents. After publishing the work, (including published elsewhere worldwide with proper authorization) you can upload your research paper with your recorded voice or you can utilize chargeable services of our professional RJs to record your paper in their voice on request.



The FARSM member also entitled to get the benefits of free research podcasting of their research documents through video clips. We can also streamline your conference videos and display your slides/ online slides and online research video clips at reasonable charges, on request.





The FARSM is eligible to earn from sales proceeds of his/her researches/reference/review Books or literature, while publishing with Global Journals. The FARSS can decide whether he/she would like to publish his/her research in a closed manner. In this case, whenever readers purchase that individual research paper for reading, maximum 60% of its profit earned as royalty by Global Journals, will be credited to his/her bank account. The entire entitled amount will be credited to his/her bank account exceeding limit of minimum fixed balance. There is no minimum time limit for collection. The FARSM member can decide its price and we can help in making the right decision.

The FARSM member is eligible to join as a paid peer reviewer at Global Journals Incorporation (USA) and can get remuneration of 15% of author fees, taken from the author of a respective paper. After reviewing 5 or more papers you can request to transfer the amount to your bank account.



## MEMBER OF ASSOCIATION OF RESEARCH SOCIETY IN MEDICAL (MARSM)

The ' MARSM ' title is accorded to a selected professional after the approval of the Editor-in-Chief / Editorial Board Members/Dean.

The “MARSM” is a dignified ornament which is accorded to a person’s name viz. Dr. John E. Hall, Ph.D., MARSM or William Walldroff, M.S., MARSM.



MARSM accrediting is an honor. It authenticates your research activities. After becoming MARSM, you can add 'MARSM' title with your name as you use this recognition as additional suffix to your status. This will definitely enhance and add more value and repute to your name. You may use it on your professional Counseling Materials such as CV, Resume, Visiting Card and Name Plate etc.

*The following benefits can be availed by you only for next three years from the date of certification.*



MARSM designated members are entitled to avail a 25% discount while publishing their research papers (of a single author) in Global Journals Inc., if the same is accepted by our Editorial Board and Peer Reviewers. If you are a main author or co-author of a group of authors, you will get discount of 10%.

As MARSM, you will be given a renowned, secure and free professional email address with 30 GB of space e.g. [johnhall@globaljournals.org](mailto:johnhall@globaljournals.org). This will include Webmail, Spam Assassin, Email Forwarders, Auto-Responders, Email Delivery Route tracing, etc.





We shall provide you intimation regarding launching of e-version of journal of your stream time to time. This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.



The MARSM member can apply for approval, grading and certification of standards of their educational and Institutional Degrees to Open Association of Research, Society U.S.A.



Once you are designated as MARSM, you may send us a scanned copy of all of your credentials. OARS will verify, grade and certify them. This will be based on your academic records, quality of research papers published by you, and some more criteria.

It is mandatory to read all terms and conditions carefully.



# AUXILIARY MEMBERSHIPS

## Institutional Fellow of Open Association of Research Society (USA) - OARS (USA)

Global Journals Incorporation (USA) is accredited by Open Association of Research Society, U.S.A (OARS) and in turn, affiliates research institutions as “Institutional Fellow of Open Association of Research Society” (IFOARS).



The “FARSC” is a dignified title which is accorded to a person’s name viz. Dr. John E. Hall, Ph.D., FARSC or William Walldroff, M.S., FARSC.

The IFOARS institution is entitled to form a Board comprised of one Chairperson and three to five board members preferably from different streams. The Board will be recognized as “Institutional Board of Open Association of Research Society”-(IBOARS).

*The Institute will be entitled to following benefits:*



The IBOARS can initially review research papers of their institute and recommend them to publish with respective journal of Global Journals. It can also review the papers of other institutions after obtaining our consent. The second review will be done by peer reviewer of Global Journals Incorporation (USA) The Board is at liberty to appoint a peer reviewer with the approval of chairperson after consulting us.

The author fees of such paper may be waived off up to 40%.

The Global Journals Incorporation (USA) at its discretion can also refer double blind peer reviewed paper at their end to the board for the verification and to get recommendation for final stage of acceptance of publication.



The IBOARS can organize symposium/seminar/conference in their country on behalf of Global Journals Incorporation (USA)-OARS (USA). The terms and conditions can be discussed separately.


The Board can also play vital role by exploring and giving valuable suggestions regarding the Standards of “Open Association of Research Society, U.S.A (OARS)” so that proper amendment can take place for the benefit of entire research community. We shall provide details of particular standard only on receipt of request from the Board.



The board members can also join us as Individual Fellow with 40% discount on total fees applicable to Individual Fellow. They will be entitled to avail all the benefits as declared. Please visit Individual Fellow-sub menu of GlobalJournals.org to have more relevant details.



We shall provide you intimation regarding launching of e-version of journal of your stream time to time. This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.



After nomination of your institution as “Institutional Fellow” and constantly functioning successfully for one year, we can consider giving recognition to your institute to function as Regional/Zonal office on our behalf. The board can also take up the additional allied activities for betterment after our consultation.

**The following entitlements are applicable to individual Fellows:**

Open Association of Research Society, U.S.A (OARS) By-laws states that an individual Fellow may use the designations as applicable, or the corresponding initials. The Credentials of individual Fellow and Associate designations signify that the individual has gained knowledge of the fundamental concepts. One is magnanimous and proficient in an expertise course covering the professional code of conduct, and follows recognized standards of practice.



Open Association of Research Society (US)/ Global Journals Incorporation (USA), as described in Corporate Statements, are educational, research publishing and professional membership organizations. Achieving our individual Fellow or Associate status is based mainly on meeting stated educational research requirements.

Disbursement of 40% Royalty earned through Global Journals : Researcher = 50%, Peer Reviewer = 37.50%, Institution = 12.50% E.g. Out of 40%, the 20% benefit should be passed on to researcher, 15 % benefit towards remuneration should be given to a reviewer and remaining 5% is to be retained by the institution.



We shall provide print version of 12 issues of any three journals [as per your requirement] out of our 38 journals worth \$ 2376 USD.

**Other:**

**The individual Fellow and Associate designations accredited by Open Association of Research Society (US) credentials signify guarantees following achievements:**

- The professional accredited with Fellow honor, is entitled to various benefits viz. name, fame, honor, regular flow of income, secured bright future, social status etc.



- In addition to above, if one is single author, then entitled to 40% discount on publishing research paper and can get 10% discount if one is co-author or main author among group of authors.
- The Fellow can organize symposium/seminar/conference on behalf of Global Journals Incorporation (USA) and he/she can also attend the same organized by other institutes on behalf of Global Journals.
- The Fellow can become member of Editorial Board Member after completing 3yrs.
- The Fellow can earn 60% of sales proceeds from the sale of reference/review books/literature/publishing of research paper.
- Fellow can also join as paid peer reviewer and earn 15% remuneration of author charges and can also get an opportunity to join as member of the Editorial Board of Global Journals Incorporation (USA)
- • This individual has learned the basic methods of applying those concepts and techniques to common challenging situations. This individual has further demonstrated an in-depth understanding of the application of suitable techniques to a particular area of research practice.

**Note :**

//

- In future, if the board feels the necessity to change any board member, the same can be done with the consent of the chairperson along with anyone board member without our approval.
- In case, the chairperson needs to be replaced then consent of 2/3rd board members are required and they are also required to jointly pass the resolution copy of which should be sent to us. In such case, it will be compulsory to obtain our approval before replacement.
- In case of “Difference of Opinion [if any]” among the Board members, our decision will be final and binding to everyone.

//



## PROCESS OF SUBMISSION OF RESEARCH PAPER

---

The Area or field of specialization may or may not be of any category as mentioned in 'Scope of Journal' menu of the GlobalJournals.org website. There are 37 Research Journal categorized with Six parental Journals GJCST, GJMR, GJRE, GJMBR, GJSFR, GJHSS. For Authors should prefer the mentioned categories. There are three widely used systems UDC, DDC and LCC. The details are available as 'Knowledge Abstract' at Home page. The major advantage of this coding is that, the research work will be exposed to and shared with all over the world as we are being abstracted and indexed worldwide.

The paper should be in proper format. The format can be downloaded from first page of 'Author Guideline' Menu. The Author is expected to follow the general rules as mentioned in this menu. The paper should be written in MS-Word Format (\*.DOC,\*.DOCX).

The Author can submit the paper either online or offline. The authors should prefer online submission.Online Submission: There are three ways to submit your paper:

**(A) (I) First, register yourself using top right corner of Home page then Login. If you are already registered, then login using your username and password.**

**(II) Choose corresponding Journal.**

**(III) Click 'Submit Manuscript'. Fill required information and Upload the paper.**

**(B) If you are using Internet Explorer, then Direct Submission through Homepage is also available.**

**(C) If these two are not convenient, and then email the paper directly to dean@globaljournals.org.**

Offline Submission: Author can send the typed form of paper by Post. However, online submission should be preferred.



# PREFERRED AUTHOR GUIDELINES

## MANUSCRIPT STYLE INSTRUCTION (Must be strictly followed)

Page Size: 8.27" X 11"

- Left Margin: 0.65
- Right Margin: 0.65
- Top Margin: 0.75
- Bottom Margin: 0.75
- Font type of all text should be Swis 721 Lt BT.
- Paper Title should be of Font Size 24 with one Column section.
- Author Name in Font Size of 11 with one column as of Title.
- Abstract Font size of 9 Bold, "Abstract" word in Italic Bold.
- Main Text: Font size 10 with justified two columns section
- Two Column with Equal Column with of 3.38 and Gaping of .2
- First Character must be three lines Drop capped.
- Paragraph before Spacing of 1 pt and After of 0 pt.
- Line Spacing of 1 pt
- Large Images must be in One Column
- Numbering of First Main Headings (Heading 1) must be in Roman Letters, Capital Letter, and Font Size of 10.
- Numbering of Second Main Headings (Heading 2) must be in Alphabets, Italic, and Font Size of 10.

**You can use your own standard format also.**

### Author Guidelines:

1. General,
2. Ethical Guidelines,
3. Submission of Manuscripts,
4. Manuscript's Category,
5. Structure and Format of Manuscript,
6. After Acceptance.

### 1. GENERAL

Before submitting your research paper, one is advised to go through the details as mentioned in following heads. It will be beneficial, while peer reviewer justify your paper for publication.

### Scope

The Global Journals Inc. (US) welcome the submission of original paper, review paper, survey article relevant to the all the streams of Philosophy and knowledge. The Global Journals Inc. (US) is parental platform for Global Journal of Computer Science and Technology, Researches in Engineering, Medical Research, Science Frontier Research, Human Social Science, Management, and Business organization. The choice of specific field can be done otherwise as following in Abstracting and Indexing Page on this Website. As the all Global



Journals Inc. (US) are being abstracted and indexed (in process) by most of the reputed organizations. Topics of only narrow interest will not be accepted unless they have wider potential or consequences.

## 2. ETHICAL GUIDELINES

Authors should follow the ethical guidelines as mentioned below for publication of research paper and research activities.

Papers are accepted on strict understanding that the material in whole or in part has not been, nor is being, considered for publication elsewhere. If the paper once accepted by Global Journals Inc. (US) and Editorial Board, will become the copyright of the Global Journals Inc. (US).

**Authorship: The authors and coauthors should have active contribution to conception design, analysis and interpretation of findings. They should critically review the contents and drafting of the paper. All should approve the final version of the paper before submission**

The Global Journals Inc. (US) follows the definition of authorship set up by the Global Academy of Research and Development. According to the Global Academy of R&D authorship, criteria must be based on:

- 1) Substantial contributions to conception and acquisition of data, analysis and interpretation of the findings.
- 2) Drafting the paper and revising it critically regarding important academic content.
- 3) Final approval of the version of the paper to be published.

All authors should have been credited according to their appropriate contribution in research activity and preparing paper. Contributors who do not match the criteria as authors may be mentioned under Acknowledgement.

Acknowledgements: Contributors to the research other than authors credited should be mentioned under acknowledgement. The specifications of the source of funding for the research if appropriate can be included. Suppliers of resources may be mentioned along with address.

**Appeal of Decision: The Editorial Board's decision on publication of the paper is final and cannot be appealed elsewhere.**

**Permissions: It is the author's responsibility to have prior permission if all or parts of earlier published illustrations are used in this paper.**

Please mention proper reference and appropriate acknowledgements wherever expected.

If all or parts of previously published illustrations are used, permission must be taken from the copyright holder concerned. It is the author's responsibility to take these in writing.

Approval for reproduction/modification of any information (including figures and tables) published elsewhere must be obtained by the authors/copyright holders before submission of the manuscript. Contributors (Authors) are responsible for any copyright fee involved.

## 3. SUBMISSION OF MANUSCRIPTS

Manuscripts should be uploaded via this online submission page. The online submission is most efficient method for submission of papers, as it enables rapid distribution of manuscripts and consequently speeds up the review procedure. It also enables authors to know the status of their own manuscripts by emailing us. Complete instructions for submitting a paper is available below.

Manuscript submission is a systematic procedure and little preparation is required beyond having all parts of your manuscript in a given format and a computer with an Internet connection and a Web browser. Full help and instructions are provided on-screen. As an author, you will be prompted for login and manuscript details as Field of Paper and then to upload your manuscript file(s) according to the instructions.



To avoid postal delays, all transaction is preferred by e-mail. A finished manuscript submission is confirmed by e-mail immediately and your paper enters the editorial process with no postal delays. When a conclusion is made about the publication of your paper by our Editorial Board, revisions can be submitted online with the same procedure, with an occasion to view and respond to all comments.

Complete support for both authors and co-author is provided.

#### 4. MANUSCRIPT'S CATEGORY

Based on potential and nature, the manuscript can be categorized under the following heads:

Original research paper: Such papers are reports of high-level significant original research work.

Review papers: These are concise, significant but helpful and decisive topics for young researchers.

Research articles: These are handled with small investigation and applications

Research letters: The letters are small and concise comments on previously published matters.

#### 5. STRUCTURE AND FORMAT OF MANUSCRIPT

The recommended size of original research paper is less than seven thousand words, review papers fewer than seven thousands words also. Preparation of research paper or how to write research paper, are major hurdle, while writing manuscript. The research articles and research letters should be fewer than three thousand words, the structure original research paper; sometime review paper should be as follows:

**Papers:** These are reports of significant research (typically less than 7000 words equivalent, including tables, figures, references), and comprise:

(a) Title should be relevant and commensurate with the theme of the paper.

(b) A brief Summary, "Abstract" (less than 150 words) containing the major results and conclusions.

(c) Up to ten keywords, that precisely identifies the paper's subject, purpose, and focus.

(d) An Introduction, giving necessary background excluding subheadings; objectives must be clearly declared.

(e) Resources and techniques with sufficient complete experimental details (wherever possible by reference) to permit repetition; sources of information must be given and numerical methods must be specified by reference, unless non-standard.

(f) Results should be presented concisely, by well-designed tables and/or figures; the same data may not be used in both; suitable statistical data should be given. All data must be obtained with attention to numerical detail in the planning stage. As reproduced design has been recognized to be important to experiments for a considerable time, the Editor has decided that any paper that appears not to have adequate numerical treatments of the data will be returned un-refereed;

(g) Discussion should cover the implications and consequences, not just recapitulating the results; conclusions should be summarizing.

(h) Brief Acknowledgements.

(i) References in the proper form.

Authors should very cautiously consider the preparation of papers to ensure that they communicate efficiently. Papers are much more likely to be accepted, if they are cautiously designed and laid out, contain few or no errors, are summarizing, and be conventional to the approach and instructions. They will in addition, be published with much less delays than those that require much technical and editorial correction.



The Editorial Board reserves the right to make literary corrections and to make suggestions to improve brevity.

It is vital, that authors take care in submitting a manuscript that is written in simple language and adheres to published guidelines.

## Format

*Language: The language of publication is UK English. Authors, for whom English is a second language, must have their manuscript efficiently edited by an English-speaking person before submission to make sure that, the English is of high excellence. It is preferable, that manuscripts should be professionally edited.*

Standard Usage, Abbreviations, and Units: Spelling and hyphenation should be conventional to The Concise Oxford English Dictionary. Statistics and measurements should at all times be given in figures, e.g. 16 min, except for when the number begins a sentence. When the number does not refer to a unit of measurement it should be spelt in full unless, it is 160 or greater.

Abbreviations supposed to be used carefully. The abbreviated name or expression is supposed to be cited in full at first usage, followed by the conventional abbreviation in parentheses.

Metric SI units are supposed to generally be used excluding where they conflict with current practice or are confusing. For illustration, 1.4 l rather than  $1.4 \times 10^{-3} \text{ m}^3$ , or 4 mm somewhat than  $4 \times 10^{-3} \text{ m}$ . Chemical formula and solutions must identify the form used, e.g. anhydrous or hydrated, and the concentration must be in clearly defined units. Common species names should be followed by underlines at the first mention. For following use the generic name should be constricted to a single letter, if it is clear.

## Structure

All manuscripts submitted to Global Journals Inc. (US), ought to include:

Title: The title page must carry an instructive title that reflects the content, a running title (less than 45 characters together with spaces), names of the authors and co-authors, and the place(s) wherever the work was carried out. The full postal address in addition with the e-mail address of related author must be given. Up to eleven keywords or very brief phrases have to be given to help data retrieval, mining and indexing.

*Abstract, used in Original Papers and Reviews:*

### Optimizing Abstract for Search Engines

Many researchers searching for information online will use search engines such as Google, Yahoo or similar. By optimizing your paper for search engines, you will amplify the chance of someone finding it. This in turn will make it more likely to be viewed and/or cited in a further work. Global Journals Inc. (US) have compiled these guidelines to facilitate you to maximize the web-friendliness of the most public part of your paper.

### Key Words

A major linchpin in research work for the writing research paper is the keyword search, which one will employ to find both library and Internet resources.

One must be persistent and creative in using keywords. An effective keyword search requires a strategy and planning a list of possible keywords and phrases to try.

Search engines for most searches, use Boolean searching, which is somewhat different from Internet searches. The Boolean search uses "operators," words (and, or, not, and near) that enable you to expand or narrow your affords. Tips for research paper while preparing research paper are very helpful guideline of research paper.

Choice of key words is first tool of tips to write research paper. Research paper writing is an art. A few tips for deciding as strategically as possible about keyword search:



- One should start brainstorming lists of possible keywords before even begin searching. Think about the most important concepts related to research work. Ask, "What words would a source have to include to be truly valuable in research paper?" Then consider synonyms for the important words.
- It may take the discovery of only one relevant paper to let steer in the right keyword direction because in most databases, the keywords under which a research paper is abstracted are listed with the paper.
- One should avoid outdated words.

Keywords are the key that opens a door to research work sources. Keyword searching is an art in which researcher's skills are bound to improve with experience and time.

Numerical Methods: Numerical methods used should be clear and, where appropriate, supported by references.

*Acknowledgements: Please make these as concise as possible.*

#### References

References follow the Harvard scheme of referencing. References in the text should cite the authors' names followed by the time of their publication, unless there are three or more authors when simply the first author's name is quoted followed by et al. unpublished work has to only be cited where necessary, and only in the text. Copies of references in press in other journals have to be supplied with submitted typescripts. It is necessary that all citations and references be carefully checked before submission, as mistakes or omissions will cause delays.

References to information on the World Wide Web can be given, but only if the information is available without charge to readers on an official site. Wikipedia and Similar websites are not allowed where anyone can change the information. Authors will be asked to make available electronic copies of the cited information for inclusion on the Global Journals Inc. (US) homepage at the judgment of the Editorial Board.

The Editorial Board and Global Journals Inc. (US) recommend that, citation of online-published papers and other material should be done via a DOI (digital object identifier). If an author cites anything, which does not have a DOI, they run the risk of the cited material not being noticeable.

The Editorial Board and Global Journals Inc. (US) recommend the use of a tool such as Reference Manager for reference management and formatting.

#### Tables, Figures and Figure Legends

*Tables: Tables should be few in number, cautiously designed, uncrowned, and include only essential data. Each must have an Arabic number, e.g. Table 4, a self-explanatory caption and be on a separate sheet. Vertical lines should not be used.*

*Figures: Figures are supposed to be submitted as separate files. Always take in a citation in the text for each figure using Arabic numbers, e.g. Fig. 4. Artwork must be submitted online in electronic form by e-mailing them.*

#### Preparation of Electronic Figures for Publication

Even though low quality images are sufficient for review purposes, print publication requires high quality images to prevent the final product being blurred or fuzzy. Submit (or e-mail) EPS (line art) or TIFF (halftone/photographs) files only. MS PowerPoint and Word Graphics are unsuitable for printed pictures. Do not use pixel-oriented software. Scans (TIFF only) should have a resolution of at least 350 dpi (halftone) or 700 to 1100 dpi (line drawings) in relation to the imitation size. Please give the data for figures in black and white or submit a Color Work Agreement Form. EPS files must be saved with fonts embedded (and with a TIFF preview, if possible).

For scanned images, the scanning resolution (at final image size) ought to be as follows to ensure good reproduction: line art: >650 dpi; halftones (including gel photographs) : >350 dpi; figures containing both halftone and line images: >650 dpi.



Color Charges: It is the rule of the Global Journals Inc. (US) for authors to pay the full cost for the reproduction of their color artwork. Hence, please note that, if there is color artwork in your manuscript when it is accepted for publication, we would require you to complete and return a color work agreement form before your paper can be published.

*Figure Legends: Self-explanatory legends of all figures should be incorporated separately under the heading 'Legends to Figures'. In the full-text online edition of the journal, figure legends may possibly be truncated in abbreviated links to the full screen version. Therefore, the first 100 characters of any legend should notify the reader, about the key aspects of the figure.*

## **6. AFTER ACCEPTANCE**

Upon approval of a paper for publication, the manuscript will be forwarded to the dean, who is responsible for the publication of the Global Journals Inc. (US).

### **6.1 Proof Corrections**

The corresponding author will receive an e-mail alert containing a link to a website or will be attached. A working e-mail address must therefore be provided for the related author.

Acrobat Reader will be required in order to read this file. This software can be downloaded

(Free of charge) from the following website:

[www.adobe.com/products/acrobat/readstep2.html](http://www.adobe.com/products/acrobat/readstep2.html). This will facilitate the file to be opened, read on screen, and printed out in order for any corrections to be added. Further instructions will be sent with the proof.

Proofs must be returned to the dean at [dean@globaljournals.org](mailto:dean@globaljournals.org) within three days of receipt.

As changes to proofs are costly, we inquire that you only correct typesetting errors. All illustrations are retained by the publisher. Please note that the authors are responsible for all statements made in their work, including changes made by the copy editor.

### **6.2 Early View of Global Journals Inc. (US) (Publication Prior to Print)**

The Global Journals Inc. (US) are enclosed by our publishing's Early View service. Early View articles are complete full-text articles sent in advance of their publication. Early View articles are absolute and final. They have been completely reviewed, revised and edited for publication, and the authors' final corrections have been incorporated. Because they are in final form, no changes can be made after sending them. The nature of Early View articles means that they do not yet have volume, issue or page numbers, so Early View articles cannot be cited in the conventional way.

### **6.3 Author Services**

Online production tracking is available for your article through Author Services. Author Services enables authors to track their article - once it has been accepted - through the production process to publication online and in print. Authors can check the status of their articles online and choose to receive automated e-mails at key stages of production. The authors will receive an e-mail with a unique link that enables them to register and have their article automatically added to the system. Please ensure that a complete e-mail address is provided when submitting the manuscript.

### **6.4 Author Material Archive Policy**

Please note that if not specifically requested, publisher will dispose off hardcopy & electronic information submitted, after the two months of publication. If you require the return of any information submitted, please inform the Editorial Board or dean as soon as possible.

### **6.5 Offprint and Extra Copies**

A PDF offprint of the online-published article will be provided free of charge to the related author, and may be distributed according to the Publisher's terms and conditions. Additional paper offprint may be ordered by emailing us at: [editor@globaljournals.org](mailto:editor@globaljournals.org) .



Before start writing a good quality Computer Science Research Paper, let us first understand what is Computer Science Research Paper? So, Computer Science Research Paper is the paper which is written by professionals or scientists who are associated to Computer Science and Information Technology, or doing research study in these areas. If you are novel to this field then you can consult about this field from your supervisor or guide.

#### TECHNIQUES FOR WRITING A GOOD QUALITY RESEARCH PAPER:

**1. Choosing the topic:** In most cases, the topic is searched by the interest of author but it can be also suggested by the guides. You can have several topics and then you can judge that in which topic or subject you are finding yourself most comfortable. This can be done by asking several questions to yourself, like Will I be able to carry our search in this area? Will I find all necessary recourses to accomplish the search? Will I be able to find all information in this field area? If the answer of these types of questions will be "Yes" then you can choose that topic. In most of the cases, you may have to conduct the surveys and have to visit several places because this field is related to Computer Science and Information Technology. Also, you may have to do a lot of work to find all rise and falls regarding the various data of that subject. Sometimes, detailed information plays a vital role, instead of short information.

**2. Evaluators are human:** First thing to remember that evaluators are also human being. They are not only meant for rejecting a paper. They are here to evaluate your paper. So, present your Best.

**3. Think Like Evaluators:** If you are in a confusion or getting demotivated that your paper will be accepted by evaluators or not, then think and try to evaluate your paper like an Evaluator. Try to understand that what an evaluator wants in your research paper and automatically you will have your answer.

**4. Make blueprints of paper:** The outline is the plan or framework that will help you to arrange your thoughts. It will make your paper logical. But remember that all points of your outline must be related to the topic you have chosen.

**5. Ask your Guides:** If you are having any difficulty in your research, then do not hesitate to share your difficulty to your guide (if you have any). They will surely help you out and resolve your doubts. If you can't clarify what exactly you require for your work then ask the supervisor to help you with the alternative. He might also provide you the list of essential readings.

**6. Use of computer is recommended:** As you are doing research in the field of Computer Science, then this point is quite obvious.

**7. Use right software:** Always use good quality software packages. If you are not capable to judge good software then you can lose quality of your paper unknowingly. There are various software programs available to help you, which you can get through Internet.

**8. Use the Internet for help:** An excellent start for your paper can be by using the Google. It is an excellent search engine, where you can have your doubts resolved. You may also read some answers for the frequent question how to write my research paper or find model research paper. From the internet library you can download books. If you have all required books make important reading selecting and analyzing the specified information. Then put together research paper sketch out.

**9. Use and get big pictures:** Always use encyclopedias, Wikipedia to get pictures so that you can go into the depth.

**10. Bookmarks are useful:** When you read any book or magazine, you generally use bookmarks, right! It is a good habit, which helps to not to lose your continuity. You should always use bookmarks while searching on Internet also, which will make your search easier.

**11. Revise what you wrote:** When you write anything, always read it, summarize it and then finalize it.



**12. Make all efforts:** Make all efforts to mention what you are going to write in your paper. That means always have a good start. Try to mention everything in introduction, that what is the need of a particular research paper. Polish your work by good skill of writing and always give an evaluator, what he wants.

**13. Have backups:** When you are going to do any important thing like making research paper, you should always have backup copies of it either in your computer or in paper. This will help you to not to lose any of your important.

**14. Produce good diagrams of your own:** Always try to include good charts or diagrams in your paper to improve quality. Using several and unnecessary diagrams will degrade the quality of your paper by creating "hotchpotch." So always, try to make and include those diagrams, which are made by your own to improve readability and understandability of your paper.

**15. Use of direct quotes:** When you do research relevant to literature, history or current affairs then use of quotes become essential but if study is relevant to science then use of quotes is not preferable.

**16. Use proper verb tense:** Use proper verb tenses in your paper. Use past tense, to present those events that happened. Use present tense to indicate events that are going on. Use future tense to indicate future happening events. Use of improper and wrong tenses will confuse the evaluator. Avoid the sentences that are incomplete.

**17. Never use online paper:** If you are getting any paper on Internet, then never use it as your research paper because it might be possible that evaluator has already seen it or maybe it is outdated version.

**18. Pick a good study spot:** To do your research studies always try to pick a spot, which is quiet. Every spot is not for studies. Spot that suits you choose it and proceed further.

**19. Know what you know:** Always try to know, what you know by making objectives. Else, you will be confused and cannot achieve your target.

**20. Use good quality grammar:** Always use a good quality grammar and use words that will throw positive impact on evaluator. Use of good quality grammar does not mean to use tough words, that for each word the evaluator has to go through dictionary. Do not start sentence with a conjunction. Do not fragment sentences. Eliminate one-word sentences. Ignore passive voice. Do not ever use a big word when a diminutive one would suffice. Verbs have to be in agreement with their subjects. Prepositions are not expressions to finish sentences with. It is incorrect to ever divide an infinitive. Avoid clichés like the disease. Also, always shun irritating alliteration. Use language that is simple and straight forward. put together a neat summary.

**21. Arrangement of information:** Each section of the main body should start with an opening sentence and there should be a changeover at the end of the section. Give only valid and powerful arguments to your topic. You may also maintain your arguments with records.

**22. Never start in last minute:** Always start at right time and give enough time to research work. Leaving everything to the last minute will degrade your paper and spoil your work.

**23. Multitasking in research is not good:** Doing several things at the same time proves bad habit in case of research activity. Research is an area, where everything has a particular time slot. Divide your research work in parts and do particular part in particular time slot.

**24. Never copy others' work:** Never copy others' work and give it your name because if evaluator has seen it anywhere you will be in trouble.

**25. Take proper rest and food:** No matter how many hours you spend for your research activity, if you are not taking care of your health then all your efforts will be in vain. For a quality research, study is must, and this can be done by taking proper rest and food.

**26. Go for seminars:** Attend seminars if the topic is relevant to your research area. Utilize all your resources.



**27. Refresh your mind after intervals:** Try to give rest to your mind by listening to soft music or by sleeping in intervals. This will also improve your memory.

**28. Make colleagues:** Always try to make colleagues. No matter how sharper or intelligent you are, if you make colleagues you can have several ideas, which will be helpful for your research.

**29. Think technically:** Always think technically. If anything happens, then search its reasons, its benefits, and demerits.

**30. Think and then print:** When you will go to print your paper, notice that tables are not be split, headings are not detached from their descriptions, and page sequence is maintained.

**31. Adding unnecessary information:** Do not add unnecessary information, like, I have used MS Excel to draw graph. Do not add irrelevant and inappropriate material. These all will create superfluous. Foreign terminology and phrases are not apropos. One should NEVER take a broad view. Analogy in script is like feathers on a snake. Not at all use a large word when a very small one would be sufficient. Use words properly, regardless of how others use them. Remove quotations. Puns are for kids, not grunt readers. Amplification is a billion times of inferior quality than sarcasm.

**32. Never oversimplify everything:** To add material in your research paper, never go for oversimplification. This will definitely irritate the evaluator. Be more or less specific. Also too, by no means, ever use rhythmic redundancies. Contractions aren't essential and shouldn't be there used. Comparisons are as terrible as clichés. Give up ampersands and abbreviations, and so on. Remove commas, that are, not necessary. Parenthetical words however should be together with this in commas. Understatement is all the time the complete best way to put onward earth-shaking thoughts. Give a detailed literary review.

**33. Report concluded results:** Use concluded results. From raw data, filter the results and then conclude your studies based on measurements and observations taken. Significant figures and appropriate number of decimal places should be used. Parenthetical remarks are prohibitive. Proofread carefully at final stage. In the end give outline to your arguments. Spot out perspectives of further study of this subject. Justify your conclusion by at the bottom of them with sufficient justifications and examples.

**34. After conclusion:** Once you have concluded your research, the next most important step is to present your findings. Presentation is extremely important as it is the definite medium through which your research is going to be in print to the rest of the crowd. Care should be taken to categorize your thoughts well and present them in a logical and neat manner. A good quality research paper format is essential because it serves to highlight your research paper and bring to light all necessary aspects in your research.

## INFORMAL GUIDELINES OF RESEARCH PAPER WRITING

### Key points to remember:

- Submit all work in its final form.
- Write your paper in the form, which is presented in the guidelines using the template.
- Please note the criterion for grading the final paper by peer-reviewers.

### Final Points:

A purpose of organizing a research paper is to let people to interpret your effort selectively. The journal requires the following sections, submitted in the order listed, each section to start on a new page.

The introduction will be compiled from reference matter and will reflect the design processes or outline of basis that direct you to make study. As you will carry out the process of study, the method and process section will be constructed as like that. The result segment will show related statistics in nearly sequential order and will direct the reviewers next to the similar intellectual paths throughout the data that you took to carry out your study. The discussion section will provide understanding of the data and projections as to the implication of the results. The use of good quality references all through the paper will give the effort trustworthiness by representing an alertness of prior workings.





Writing a research paper is not an easy job no matter how trouble-free the actual research or concept. Practice, excellent preparation, and controlled record keeping are the only means to make straightforward the progression.

**General style:**

Specific editorial column necessities for compliance of a manuscript will always take over from directions in these general guidelines.

To make a paper clear

- Adhere to recommended page limits

Mistakes to evade

- Insertion a title at the foot of a page with the subsequent text on the next page
- Separating a table/chart or figure - impound each figure/table to a single page
- Submitting a manuscript with pages out of sequence

In every sections of your document

- Use standard writing style including articles ("a", "the," etc.)
- Keep on paying attention on the research topic of the paper
- Use paragraphs to split each significant point (excluding for the abstract)
- Align the primary line of each section
- Present your points in sound order
- Use present tense to report well accepted
- Use past tense to describe specific results
- Shun familiar wording, don't address the reviewer directly, and don't use slang, slang language, or superlatives
- Shun use of extra pictures - include only those figures essential to presenting results

**Title Page:**

Choose a revealing title. It should be short. It should not have non-standard acronyms or abbreviations. It should not exceed two printed lines. It should include the name(s) and address (es) of all authors.



## Abstract:

The summary should be two hundred words or less. It should briefly and clearly explain the key findings reported in the manuscript-- must have precise statistics. It should not have abnormal acronyms or abbreviations. It should be logical in itself. Shun citing references at this point.

An abstract is a brief distinct paragraph summary of finished work or work in development. In a minute or less a reviewer can be taught the foundation behind the study, common approach to the problem, relevant results, and significant conclusions or new questions.

Write your summary when your paper is completed because how can you write the summary of anything which is not yet written? Wealth of terminology is very essential in abstract. Yet, use comprehensive sentences and do not let go readability for brevity. You can maintain it succinct by phrasing sentences so that they provide more than lone rationale. The author can at this moment go straight to shortening the outcome. Sum up the study, with the subsequent elements in any summary. Try to maintain the initial two items to no more than one ruling each.

- Reason of the study - theory, overall issue, purpose
- Fundamental goal
- To the point depiction of the research
- Consequences, including definite statistics - if the consequences are quantitative in nature, account quantitative data; results of any numerical analysis should be reported
- Significant conclusions or questions that track from the research(es)

## Approach:

- Single section, and succinct
- As an outline of job done, it is always written in past tense
- A conceptual should situate on its own, and not submit to any other part of the paper such as a form or table
- Center on shortening results - bound background information to a verdict or two, if completely necessary
- What you account in an abstract must be regular with what you reported in the manuscript
- Exact spelling, clearness of sentences and phrases, and appropriate reporting of quantities (proper units, important statistics) are just as significant in an abstract as they are anywhere else

## Introduction:

The **Introduction** should "introduce" the manuscript. The reviewer should be presented with sufficient background information to be capable to comprehend and calculate the purpose of your study without having to submit to other works. The basis for the study should be offered. Give most important references but shun difficult to make a comprehensive appraisal of the topic. In the introduction, describe the problem visibly. If the problem is not acknowledged in a logical, reasonable way, the reviewer will have no attention in your result. Speak in common terms about techniques used to explain the problem, if needed, but do not present any particulars about the protocols here. Following approach can create a valuable beginning:

- Explain the value (significance) of the study
- Shield the model - why did you employ this particular system or method? What is its compensation? You strength remark on its appropriateness from a abstract point of vision as well as point out sensible reasons for using it.
- Present a justification. Status your particular theory (es) or aim(s), and describe the logic that led you to choose them.
- Very for a short time explain the tentative propose and how it skilled the declared objectives.

## Approach:

- Use past tense except for when referring to recognized facts. After all, the manuscript will be submitted after the entire job is done.
- Sort out your thoughts; manufacture one key point with every section. If you make the four points listed above, you will need a least of four paragraphs.



- Present surroundings information only as desirable in order hold up a situation. The reviewer does not desire to read the whole thing you know about a topic.
- Shape the theory/purpose specifically - do not take a broad view.
- As always, give awareness to spelling, simplicity and correctness of sentences and phrases.

#### **Procedures (Methods and Materials):**

This part is supposed to be the easiest to carve if you have good skills. A sound written Procedures segment allows a capable scientist to replacement your results. Present precise information about your supplies. The suppliers and clarity of reagents can be helpful bits of information. Present methods in sequential order but linked methodologies can be grouped as a segment. Be concise when relating the protocols. Attempt for the least amount of information that would permit another capable scientist to spare your outcome but be cautious that vital information is integrated. The use of subheadings is suggested and ought to be synchronized with the results section. When a technique is used that has been well described in another object, mention the specific item describing a way but draw the basic principle while stating the situation. The purpose is to text all particular resources and broad procedures, so that another person may use some or all of the methods in one more study or referee the scientific value of your work. It is not to be a step by step report of the whole thing you did, nor is a methods section a set of orders.

#### **Materials:**

- Explain materials individually only if the study is so complex that it saves liberty this way.
- Embrace particular materials, and any tools or provisions that are not frequently found in laboratories.
- Do not take in frequently found.
- If use of a definite type of tools.
- Materials may be reported in a part section or else they may be recognized along with your measures.

#### **Methods:**

- Report the method (not particulars of each process that engaged the same methodology)
- Describe the method entirely
- To be succinct, present methods under headings dedicated to specific dealings or groups of measures
- Simplify - details how procedures were completed not how they were exclusively performed on a particular day.
- If well known procedures were used, account the procedure by name, possibly with reference, and that's all.

#### **Approach:**

- It is embarrassed or not possible to use vigorous voice when documenting methods with no using first person, which would focus the reviewer's interest on the researcher rather than the job. As a result when script up the methods most authors use third person passive voice.
- Use standard style in this and in every other part of the paper - avoid familiar lists, and use full sentences.

#### **What to keep away from**

- Resources and methods are not a set of information.
- Skip all descriptive information and surroundings - save it for the argument.
- Leave out information that is immaterial to a third party.

#### **Results:**

The principle of a results segment is to present and demonstrate your conclusion. Create this part a entirely objective details of the outcome, and save all understanding for the discussion.

The page length of this segment is set by the sum and types of data to be reported. Carry on to be to the point, by means of statistics and tables, if suitable, to present consequences most efficiently. You must obviously differentiate material that would usually be incorporated in a study editorial from any unprocessed data or additional appendix matter that would not be available. In fact, such matter should not be submitted at all except requested by the instructor.



## Content

- Sum up your conclusion in text and demonstrate them, if suitable, with figures and tables.
- In manuscript, explain each of your consequences, point the reader to remarks that are most appropriate.
- Present a background, such as by describing the question that was addressed by creation an exacting study.
- Explain results of control experiments and comprise remarks that are not accessible in a prescribed figure or table, if appropriate.
- Examine your data, then prepare the analyzed (transformed) data in the form of a figure (graph), table, or in manuscript form.

### What to stay away from

- Do not discuss or infer your outcome, report surroundings information, or try to explain anything.
- Not at all, take in raw data or intermediate calculations in a research manuscript.
- Do not present the similar data more than once.
- Manuscript should complement any figures or tables, not duplicate the identical information.
- Never confuse figures with tables - there is a difference.

### Approach

- As forever, use past tense when you submit to your results, and put the whole thing in a reasonable order.
- Put figures and tables, appropriately numbered, in order at the end of the report
- If you desire, you may place your figures and tables properly within the text of your results part.

### Figures and tables

- If you put figures and tables at the end of the details, make certain that they are visibly distinguished from any attach appendix materials, such as raw facts
- Despite of position, each figure must be numbered one after the other and complete with subtitle
- In spite of position, each table must be titled, numbered one after the other and complete with heading
- All figure and table must be adequately complete that it could situate on its own, divide from text

### Discussion:

The Discussion is expected the trickiest segment to write and describe. A lot of papers submitted for journal are discarded based on problems with the Discussion. There is no head of state for how long a argument should be. Position your understanding of the outcome visibly to lead the reviewer through your conclusions, and then finish the paper with a summing up of the implication of the study. The purpose here is to offer an understanding of your results and hold up for all of your conclusions, using facts from your research and generally accepted information, if suitable. The implication of result should be visibly described. Infer your data in the conversation in suitable depth. This means that when you clarify an observable fact you must explain mechanisms that may account for the observation. If your results vary from your prospect, make clear why that may have happened. If your results agree, then explain the theory that the proof supported. It is never suitable to just state that the data approved with prospect, and let it drop at that.

- Make a decision if each premise is supported, discarded, or if you cannot make a conclusion with assurance. Do not just dismiss a study or part of a study as "uncertain."
- Research papers are not acknowledged if the work is imperfect. Draw what conclusions you can based upon the results that you have, and take care of the study as a finished work
- You may propose future guidelines, such as how the experiment might be personalized to accomplish a new idea.
- Give details all of your remarks as much as possible, focus on mechanisms.
- Make a decision if the tentative design sufficiently addressed the theory, and whether or not it was correctly restricted.
- Try to present substitute explanations if sensible alternatives be present.
- One research will not counter an overall question, so maintain the large picture in mind, where do you go next? The best studies unlock new avenues of study. What questions remain?
- Recommendations for detailed papers will offer supplementary suggestions.

### Approach:

- When you refer to information, differentiate data generated by your own studies from available information
- Submit to work done by specific persons (including you) in past tense.
- Submit to generally acknowledged facts and main beliefs in present tense.



## THE ADMINISTRATION RULES

Please carefully note down following rules and regulation before submitting your Research Paper to Global Journals Inc. (US):

**Segment Draft and Final Research Paper:** You have to strictly follow the template of research paper. If it is not done your paper may get rejected.

- The **major constraint** is that you must independently make all content, tables, graphs, and facts that are offered in the paper. You must write each part of the paper wholly on your own. The Peer-reviewers need to identify your own perceptives of the concepts in your own terms. NEVER extract straight from any foundation, and never rephrase someone else's analysis.
- Do not give permission to anyone else to "PROOFREAD" your manuscript.
- **Methods to avoid Plagiarism is applied by us on every paper, if found guilty, you will be blacklisted by all of our collaborated research groups, your institution will be informed for this and strict legal actions will be taken immediately.)**
- To guard yourself and others from possible illegal use please do not permit anyone right to use to your paper and files.



CRITERION FOR GRADING A RESEARCH PAPER (COMPILATION)  
BY GLOBAL JOURNALS INC. (US)

Please note that following table is only a Grading of "Paper Compilation" and not on "Performed/Stated Research" whose grading solely depends on Individual Assigned Peer Reviewer and Editorial Board Member. These can be available only on request and after decision of Paper. This report will be the property of Global Journals Inc. (US).

Topics	Grades		
	A-B	C-D	E-F
<i>Abstract</i>	Clear and concise with appropriate content, Correct format. 200 words or below	Unclear summary and no specific data, Incorrect form  Above 200 words	No specific data with ambiguous information  Above 250 words
<i>Introduction</i>	Containing all background details with clear goal and appropriate details, flow specification, no grammar and spelling mistake, well organized sentence and paragraph, reference cited	Unclear and confusing data, appropriate format, grammar and spelling errors with unorganized matter	Out of place depth and content, hazy format
<i>Methods and Procedures</i>	Clear and to the point with well arranged paragraph, precision and accuracy of facts and figures, well organized subheads	Difficult to comprehend with embarrassed text, too much explanation but completed	Incorrect and unorganized structure with hazy meaning
<i>Result</i>	Well organized, Clear and specific, Correct units with precision, correct data, well structuring of paragraph, no grammar and spelling mistake	Complete and embarrassed text, difficult to comprehend	Irregular format with wrong facts and figures
<i>Discussion</i>	Well organized, meaningful specification, sound conclusion, logical and concise explanation, highly structured paragraph reference cited	Wordy, unclear conclusion, spurious	Conclusion is not cited, unorganized, difficult to comprehend
<i>References</i>	Complete and correct format, well organized	Beside the point, Incomplete	Wrong format and structuring



# INDEX

---

---

## **A**

Agroclimatic · 4  
Antemortem · 15, 16, 17

---

## **C**

Camelus · 1  
Cysticercosis · 1, 2, 3, 4, 5, 6, 7, 15

---

## **D**

Deconinck · 1, 3  
Dromedarius · 1

---

## **E**

Epidemiological · 1, 2, 9, 10, 5, 2

---

## **F**

Fasciola · 1, 2, 3, 6, 7, 9, 10, 11, 18  
Flukicides · 8  
Frehwot · 3, 13

---

## **G**

Gesellsfür · 2  
Gigantica · 1, 2, 3, 6, 9, 10, 11

---

## **H**

Hydatidosis · 9, 15, 16, 17, 18, 19

---

## **L**

Lidetu · 3, 13

---

## **M**

Metacercariae · 4

---

## **O**

Oocysts · 1, 3, 2

---

## **P**

Proglottids · 3

---

## **T**

Taeniasis · 1, 5, 6, 7, 2  
Toxoplasma · 1, 2, 3  
Trematode · 1, 5, 10




save our planet



# Global Journal of Medical Research

Visit us on the Web at [www.GlobalJournals.org](http://www.GlobalJournals.org) | [www.MedicalResearchJournal.org](http://www.MedicalResearchJournal.org)  
or email us at [helpdesk@globaljournals.org](mailto:helpdesk@globaljournals.org)

ISSN 9755896



© Global Journals