

# GLOBAL JOURNAL

OF MEDICAL RESEARCH: E

## Gynecology & Obstetrics

Massive Ascites Complicating

Incidence & Fetomaternal Outcome

Highlights

Biochemical Markers and Uterine

Incidence, Clinical Characteristics

Discovering Thoughts, Inventing Future

VOLUME 17    ISSUE 2    VERSION 1.0



GLOBAL JOURNAL OF MEDICAL RESEARCH: E  
GYNECOLOGY AND OBSTETRICS

---



GLOBAL JOURNAL OF MEDICAL RESEARCH: E  
GYNECOLOGY AND OBSTETRICS

---

VOLUME 17 ISSUE 2 (VER. 1.0)

© Global Journal of Medical Research. 2017.

All rights reserved.

This is a special issue published in version 1.0 of "Global Journal of Medical Research." By Global Journals Inc.

All articles are open access articles distributed under "Global Journal of Medical Research"

Reading License, which permits restricted use. Entire contents are copyright by of "Global Journal of Medical Research" unless otherwise noted on specific articles.

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopy, recording, or any information storage and retrieval system, without written permission.

The opinions and statements made in this book are those of the authors concerned. Ultraculture has not verified and neither confirms nor denies any of the foregoing and no warranty or fitness is implied.

Engage with the contents herein at your own risk.

The use of this journal, and the terms and conditions for our providing information, is governed by our Disclaimer, Terms and Conditions and Privacy Policy given on our website <http://globaljournals.us/terms-and-condition/menu-id-1463/>

By referring / using / reading / any type of association / referencing this journal, this signifies and you acknowledge that you have read them and that you accept and will be bound by the terms thereof.

All information, journals, this journal, activities undertaken, materials, services and our website, terms and conditions, privacy policy, and this journal is subject to change anytime without any prior notice.

Incorporation No.: 0423089  
License No.: 42125/022010/1186  
Registration No.: 430374  
Import-Export Code: 1109007027  
Employer Identification Number (EIN):  
USA Tax ID: 98-0673427

## Global Journals Inc.

(A Delaware USA Incorporation with "Good Standing"; Reg. Number: 0423089)  
Sponsors: Open Association of Research Society  
Open Scientific Standards

### *Publisher's Headquarters office*

Global Journals® Headquarters  
945th Concord Streets,  
Framingham Massachusetts Pin: 01701,  
United States of America  
USA Toll Free: +001-888-839-7392  
USA Toll Free Fax: +001-888-839-7392

### *Offset Typesetting*

Global Journals Incorporated  
2nd, Lansdowne, Lansdowne Rd., Croydon-Surrey,  
Pin: CR9 2ER, United Kingdom

### *Packaging & Continental Dispatching*

Global Journals Pvt. Ltd.  
E-3130 Sudama Nagar, Near Gopur Square,  
Indore, M.P., Pin: 452009, India

### *Find a correspondence nodal officer near you*

To find nodal officer of your country, please  
email us at [local@globaljournals.org](mailto:local@globaljournals.org)

### *eContacts*

Press Inquiries: [press@globaljournals.org](mailto:press@globaljournals.org)  
Investor Inquiries: [investors@globaljournals.org](mailto:investors@globaljournals.org)  
Technical Support: [technology@globaljournals.org](mailto:technology@globaljournals.org)  
Media & Releases: [media@globaljournals.org](mailto:media@globaljournals.org)

### *Pricing (Including by Air Parcel Charges):*

#### *For Authors:*

22 USD (B/W) & 50 USD (Color)  
Yearly Subscription (Personal & Institutional):  
200 USD (B/W) & 250 USD (Color)

# EDITORIAL BOARD

GLOBAL JOURNAL OF MEDICAL RESEARCH

## *Dr. Apostolos Ch. Zarros*

DM, Degree (Ptychio) holder in Medicine,  
National and Kapodistrian University of Athens  
MRes, Master of Research in Molecular Functions in  
Disease, University of Glasgow FRNS, Fellow, Royal  
Numismatic Society Member, European Society for  
Neurochemistry Member, Royal Institute of Philosophy  
Scotland, United Kingdom

## *Dr. Alfio Ferlito*

Professor Department of Surgical Sciences  
University of Udine School of Medicine, Italy

## *Dr. Jixin Zhong*

Department of Medicine, Affiliated Hospital of  
Guangdong Medical College, Zhanjiang, China, Davis  
Heart and Lung Research Institute, The Ohio State  
University, Columbus, OH 43210, US

## *Rama Rao Ganga*

MBBS  
MS (University of Health Sciences, Vijayawada, India)  
MRCS (Royal College of Surgeons of Edinburgh, UK)  
United States

## *Dr. Izzet Yavuz*

MSc, Ph.D., D Ped Dent.  
Associate Professor, Pediatric Dentistry Faculty of  
Dentistry, University of Dicle Diyarbakir, Turkey

## *Sanguansak Rerksuppaphol*

Department of Pediatrics Faculty of Medicine  
Srinakharinwirot University  
NakornNayok, Thailand

## *Dr. William Chi-shing Cho*

Ph.D.,  
Department of Clinical Oncology  
Queen Elizabeth Hospital  
Hong Kong

## *Dr. Michael Wink*

Ph.D., Technical University Braunschweig, Germany  
Head of Department Institute of Pharmacy and Molecular  
Biotechnology, Heidelberg University, Germany

## *Dr. Pejic Ana*

Assistant Medical Faculty Department of Periodontology  
and Oral Medicine University of Nis, Serbia

## *Dr. Ivandro Soares Monteiro*

M.Sc., Ph.D. in Psychology Clinic, Professor University of  
Minho, Portugal

## *Dr. Sanjay Dixit, M.D.*

Director, EP Laboratories, Philadelphia VA Medical Center  
Cardiovascular Medicine - Cardiac Arrhythmia  
Univ of Penn School of Medicine  
Web: [pennmedicine.org/wagform/MainPage.aspx?](http://pennmedicine.org/wagform/MainPage.aspx?)

## *Antonio Simone Laganà*

M.D. Unit of Gynecology and Obstetrics  
Department of Human Pathology in Adulthood and  
Childhood "G. Barresi" University of Messina, Italy

*Dr. Han-Xiang Deng*

MD., Ph.D  
Associate Professor and Research Department  
Division of Neuromuscular Medicine  
Davee Department of Neurology and Clinical  
Neurosciences  
Northwestern University Feinberg School of Medicine  
Web: [neurology.northwestern.edu/faculty/deng.html](http://neurology.northwestern.edu/faculty/deng.html)

*Dr. Roberto Sanchez*

Associate Professor  
Department of Structural and Chemical Biology  
Mount Sinai School of Medicine  
Ph.D., The Rockefeller University  
Web: [mountsinai.org/](http://mountsinai.org/)

*Dr. Feng Feng*

Boston University  
Microbiology  
72 East Concord Street R702  
Duke University  
United States of America

*Dr. Pina C. Sanelli*

Associate Professor of Radiology  
Associate Professor of Public Health  
Weill Cornell Medical College  
Associate Attending Radiologist  
NewYork-Presbyterian Hospital  
MRI, MRA, CT, and CTA  
Neuroradiology and Diagnostic Radiology  
M.D., State University of New York at Buffalo,  
School of Medicine and Biomedical Sciences  
Web: [weillcornell.org/pinasanelli/](http://weillcornell.org/pinasanelli/)

*Dr. Michael R. Rudnick*

M.D., FACP  
Associate Professor of Medicine  
Chief, Renal Electrolyte and Hypertension Division (PMC)  
Penn Medicine, University of Pennsylvania  
Presbyterian Medical Center, Philadelphia  
Nephrology and Internal Medicine  
Certified by the American Board of Internal Medicine  
Web: [uphs.upenn.edu/](http://uphs.upenn.edu/)

*Dr. Seung-Yup Ku*

M.D., Ph.D., Seoul National University Medical College,  
Seoul, Korea Department of Obstetrics and Gynecology  
Seoul National University Hospital, Seoul, Korea

## CONTENTS OF THE ISSUE

---

- i. Copyright Notice
- ii. Editorial Board Members
- iii. Chief Author and Dean
- iv. Contents of the Issue
  1. Incidence and Fetomaternal Outcome of Eclampsia in a Tertiary Medical College Hospital in Bangladesh. **1-4**
  2. Successful Outcome of Pregnancy after Cystectomy for Twisted Ovarian Cyst : A Case Report. **5-7**
  3. Biochemical Markers and Uterine Artery Doppler Study for the Prediction and the Severity of the Hypertensive Disorders during Pregnancy. **9-13**
  4. Awareness of Signs of Obstetric Complications amongst Married Couples: Implications for Decision-Making towards Care-Seeking. A Qualitative Study of Selected Districts in Ghana. **15-25**
  5. Massive Ascites Complicating Severe Preeclampsia : A Case Report. **27-28**
  6. Incidence, Clinical Characteristics and Outcome of Unexpected Uterine Sarcoma after Hysterectomy for Uterine Mass: A Retrospective Study of 774 Cases. **29-32**
  7. Effect of Pregnancy and Mode of Delivery on HPV Infection and on Cervical Cytological Changes. **33-39**
- v. Fellows
- vi. Auxiliary Memberships
- vii. Process of Submission of Research Paper
- viii. Preferred Author Guidelines
- ix. Index



GLOBAL JOURNAL OF MEDICAL RESEARCH: E  
GYNECOLOGY AND OBSTETRICS  
Volume 17 Issue 2 Version 1.0 Year 2017  
Type: Double Blind Peer Reviewed International Research Journal  
Publisher: Global Journals Inc. (USA)  
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

## Incidence and Fetomaternal Outcome of Eclampsia in a Tertiary Medical College Hospital in Bangladesh

By Kamrun Nessa, Sanjida khan, Selina Begum, Ferdowsi Sultana & Tania Akbar

*Enam Medical College & Hospital*

**Abstract- Background:** In day to day obstetric practice, the incidence of eclampsia and its complication is high. It is one of the leading causes of maternal mortality in Bangladesh accounting about 20%. This is also associated with poor perinatal outcome.

**Objectives:** To determine the incidence of eclampsia and associated feto-maternal outcome.

**Materials and Methods:** A retrospective study was conducted on the diagnosed patients of eclampsia admitted in Enam Medical College and hospital, Dhaka, Bangladesh, during the period of January 2015 to December 2016. During these years total 2295 patients were admitted in obstetrics unit among them 35 patient were admitted with eclampsia. Analysis done regarding age of women, parity, type of eclampsia, gestational age of delivery, mode of delivery, maternal and fetal outcome. Patients with convulsion and or coma due to other causes were excluded. Study was ethically approved by Ethical Review committee of Enam Medical College and Hospital. All data were entered in SPSS16 and also analyzed through it.

**Keywords:** *eclampsia, perinatal death, preterm delivery.*

**GJMR-E Classification:** NLMC Code: WQ 215



*Strictly as per the compliance and regulations of:*



© 2017. Kamrun Nessa, Sanjida khan, Selina Begum, Ferdowsi Sultana & Tania Akbar. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License <http://creativecommons.org/licenses/by-nc/3.0/>), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.



# Incidence and Fetomaternal Outcome of Eclampsia in a Tertiary Medical College Hospital in Bangladesh

Kamrun Nessa <sup>α</sup>, Sanjida khan <sup>σ</sup>, Selina Begum <sup>ρ</sup>, Ferdowsi Sultana <sup>ω</sup> & Tania Akbar <sup>¥</sup>

**Abstract- Background:** In day to day obstetric practice, the incidence of eclampsia and its complication is high. It is one of the leading causes of maternal mortality in Bangladesh accounting about 20%. This is also associated with poor perinatal outcome.

**Objectives:** To determine the incidence of eclampsia and associated fetomaternal outcome.

**Materials and Methods:** A retrospective study was conducted on the diagnosed patients of eclampsia admitted in Enam Medical College and hospital, Dhaka, Bangladesh, during the period of January 2015 to December 2016. During these years total 2295 patients were admitted in obstetrics unit among them 35 patient were admitted with eclampsia. Analysis done regarding age of women, parity, type of eclampsia, gestational age of delivery, mode of delivery, maternal and fetal outcome. Patients with convulsion and or coma due to other causes were excluded. Study was ethically approved by Ethical Review committee of Enam Medical College and Hospital. All data were entered in SPSS16 and also analyzed through it.

**Results:** Total admitted patients in obstetrics unit during the given time period were 2295. Among them 35(1.52%) were eclampsia. Among eclampsia patients more than half that is 19(54.28%) were in between age of 21 to 30 years of age. Most 21(60%) were nulliparous and 14(40%) were multiparous. 15 (42.85%) were antepartum eclampsia whereas 20(57.14%) were postpartum. Maximum 22(62.85%) delivered preterm. Mode of delivery by caesarean section 24(68.57%) is more than double of vaginal delivery. Maternal outcome was good after treatment in eclampsia ward in 18 (51.42%) cases, whereas 14 (40%) needed ICU support and 3 (8.57%) died in hospital. Most of the babies were born preterm low birth weight (37.14%) and perinatal deaths were 20%. Only 6 babies were term and healthy.

**Conclusion:** Our observation states postpartum eclampsia is more common in nulliparous between 21-30 years of age with preterm delivery and poor fetomaternal outcome.

**Author α:** Professor, Department of Obstetrics & Gynecology, Enam Medical College & Hospital, Dhaka, Bangladesh.  
e-mail: knessa3@gmail.com

**Author σ:** Assistant Professor, Department of Obstetrics & Gynecology, Enam Medical College & Hospital, Dhaka, Bangladesh.  
e-mail: sanjida.jimch@yahoo.com

**Author ρ:** Classified specialist, Department of Obstetrics & Gynecology, Combined Military Hospital, Comilla, Bangladesh.  
e-mail: selinabegum101101@yahoo.com

**Author ω:** Registrar, Department of Obstetrics & Gynecology, Enam Medical College & Hospital, Dhaka, Bangladesh.  
e-mail: drfspoly@gmail.com

**Author ¥:** Assistant Professor, Department of Obstetrics & Gynecology, Enam Medical College & Hospital, Dhaka, Bangladesh.  
e-mail: taniaakbarfcp@gmail.com

**Keywords:** eclampsia, perinatal death, preterm delivery.

## I. INTRODUCTION

Now a days Eclampsia is one of the leading cause of maternal and perinatal mortality as well as morbidity throughout the world<sup>1,2</sup>. Pre-eclampsia when complicated with generalized tonic-clonic convulsions and/or coma is called eclampsia<sup>3</sup>. Since eclampsia is a severe form of pre-eclampsia, early detection of risk factors, symptoms and signs by good antenatal checkup and initiation therapy will prevent occurrence of eclampsia. Unfortunately, eclampsia still complicates much larger number of pregnancies in the world. It is estimated that every year eclampsia is associated with about 50, 000 maternal death worldwide, most of which occur in developing countries<sup>4</sup>. The incidence of eclampsia has been reduced to. 2%-5 % of all deliveries in developed countries. But in Bangladesh about 5% of the total pregnancies develop eclampsia<sup>5</sup>. There are approximately 3.6 million births per year in Bangladesh and over 10,000 women develop eclampsia each year<sup>6</sup>. It is one of the common cause of maternal mortality and responsible for 20% of maternal death<sup>7</sup>. Incidence of eclampsia varies inversely with the quality of antenatal care.

## II. METHODOLOGY

This retrospective study was conducted on the diagnosed patients of eclampsia admitted in Enam medical college hospital, Savar, Dhaka, Bangladesh, during the period of January 2015 to December 2016. During these months total 2295 patients were admitted in obstetric ward and among them 35 patients were eclampsia. Age of patients, parity, type of eclampsia, gestational age, mode of delivery, maternal and fetal outcome were noted from medical records of patients. Patients with diagnosis of convulsion and coma of other causes were excluded.

All patients of eclampsia were followed up thoroughly from admission to discharge. Study was ethically approved by Ethical Review committee of Enam Medical College and Hospital. All data were entered in SPSS16 and also analyzed through it.

### III. RESULTS

35 cases of eclampsia were recorded out of 2295 obstetric patients admitted during the study period which gives an incidence of 1.52%. 19(54.28%) cases were in between age of 21 to 30 years, while 12(34.28%) were less than 21 years and 4(11.42%) were more than 30 years as illustrated in Table 1.

Table 1: Age distribution of patients of eclampsia

| Age            | Frequency | Percentage |
|----------------|-----------|------------|
| < 21 Years     | 12        | 34.28      |
| 21 to 30 years | 19        | 54.28      |
| > 30 years     | 4         | 11.42      |

Majority (60%) were nulliparous, while 40% were multiparous which is shown in Fig 1.




Fig. 1: Distribution of study patients according to parity

Most (62.85%) of the cases delivered preterm and 37.14% at term. Caesarean section were common mode of delivery being 24 (68.57%) cases, whereas delivered by vaginally only 13 (37.14%) cases. 15 (42.85%) had convulsion before delivery and 20 (57.14%) thereafter as illustrated in Fig 2, 3 and 4.




Fig. 2: Distribution of study group according to mode of delivery




Fig. 3: Distribution of study group according to duration of pregnancy




Fig. 4: Distribution of study population according to type of eclampsia




Fig. 5: Distribution of study group according to maternal outcome

**Table 2:** Distribution of study subjects according to fetal outcome

| Fetal Outcome   | Frequency | Percentage |
|-----------------|-----------|------------|
| Term Healthy    | 6         | 17.14      |
| IUGR            | 9         | 25.71      |
| Preterm LBW     | 13        | 37.14      |
| Perinatal Death | 7         | 20         |

After meticulous treatment 18(51.42%) patients improved in eclampsia ward, while 14 (40%) patients needed ICU support and 3(8.57%) died. Among the babies only 6(17.14%) were term and healthy. Most (37.14%) of the babies were preterm low birth weight, 9 (25.71%) were IUGR and 7 (20%) were perinatal death which is shown in Fig 5 and Table 2.

#### IV. DISCUSSION

The incidence of eclampsia in our hospital was 1.52% of total obstetric admission.. Onuh in Benin Nigeria reported 1.32% and Okafor recently reported an incidence of 0.82% in Abuja, Nigeria<sup>8</sup>. A high incidence of eclampsia is common in developing countries where most patients have no antenatal care which would allow for early recognition and treatment of eclampsia. Majority of the patients were between age 21 to 30 years in this study which is contrary to the report in the developed world where severe preeclampsia with severe features and eclampsia is significantly commoner among women older than 40 years<sup>9</sup>. Nulliparity strongly associated with eclampsia in this study is supported by previous reports in other centres<sup>10-12</sup>.

Antepartum eclampsia accounted for 42.85% in this study is higher than 36.8% reported in Lagos but lower than 61.6%,84% and 85% reported in Ethiopia, Enugu and Ibadan respectively<sup>11,13-15</sup>. In our study post partum eclampsia was more common, about 57.14% which correlate with another study on similar topic<sup>16</sup>.

The majority of antepartum cases in our study had eclampsia before term and caesarean section was a leading mode of delivery, which is comparable to other studies<sup>17,18</sup>. In this study patients underwent caesarean section due to an unfavourable cervix remote from delivery, but the decision to perform a caesarean delivery was based on multiple factors which included gestational age, foetal condition, stage of labour and Bishop scoring of cervix<sup>19</sup>.

Maternal mortality rate of 8.57% reported in this study was higher than 7.9%, 8% and 9% reported respectively from Tanzania, India and Ibadan, Nigeria<sup>14,20,21</sup>.

Hypertensive disorders are a common cause of preterm labour, perinatal death and intrauterine growth restriction<sup>22</sup>. The 20% perinatal mortality rate in this study is higher than 10% reported from Ibadan, but lower than 29% and 40.9% reported from Ethiopia and Kaduna,

Nigeria respectively<sup>23,14,24</sup>. A significant percentage (37.14%) of low birth weight neonates might have been the result of the high number of preterm deliveries among the eclamptic patients. Similar findings have been reported in the literature that links the incidence of low birth weight infants with preterm deliveries in eclamptic patients<sup>17,25,26</sup>.

#### V. CONCLUSION

The incidence of eclampsia remains high in our hospital. Fetomaternal morbidity and mortality are in alarming rate. Hence, eclampsia remains a continuing problem in developing countries and leading cause of fetal-maternal mortality and morbidity. Careful antenatal supervision, early detection and management of high risk cases can reduce this dreadful disease.

#### REFERENCES RÉFÉRENCES REFERENCIAS

1. Tuffnell D.J., Jankowicz D., Lindow S.W., Lyons G., Mason G.C., Rusell L.F., Walker J.J. (2005). Outcomes of severe pre-eclampsia/ eclampsia in Yorkshire 1999/2003, *BJOG: An International Journal of Obstetrics and Gynaecology*, 112, 875-880. doi:10.1111/j.14710528.2005.00565.x.
2. Ikechebelu,JI, Okoli, C.C.(2002). Review of eclampsia at the Nnamdi Azikiwe University Teaching Hospital, Nnewi. (January 1996-December 2000), *Journal of Obstetrics and Gynaecology* 22(3), 287-280. doi: 10.1080/01443610220130580.
3. Dutta, D.C. Text Book of Obstetrics. 7th Edition, 2011; 230. Retrieved from: <https://www.meripustak.com/DCDutta-Textbook-Of-Obstetrics-7th-Edition/Obstetrics-and-Gynecology-Books/Books/pid-105443>.
4. Tukur, J. (2009).The use of magnesium sulphate for the treatment of severe pre-eclampsia and eclampsia. *Annals of African Medicine*, 8(2), 76-80. doi: 10.4103/1596-3519.56232.
5. Khatun, S. (1994). Perinatal outcome in eclampsia, Bangladesh. *The Journal of Obstetrics & Gynecology Research*, 9, 53-60.
6. Nahar, S., Rasul, C.H., Sayed, A. & Azim, A.K. (2004). Utility of misoprostol for labour induction in severe pre-eclampsia and eclampsia. *The Journal of Obstetrics & Gynecology Research*, 30(5), 349-53. doi:10.1111/j.1447-0756.2004.00207.x BMMS 2010.
7. Efetie,R.E., Okafor, U.V. (2007). Maternal outcome in Eclamptic patients in Abuja, Nigeria-5 years review. *Nigeria J clin Pract*, 10(3), 309-313, Retrieved from: <https://www.ncbi.nlm.nih.gov/pubmed/18293641>
8. Duley, L. (2003). Pre-eclampsia and hypertensive disorder in pregnancy. *Br Med Bull*, 67, 161-76. Retrieved from: Pub Med
9. Okogbenin, S.A., Eigbefoh, J.O., Omorogbe, F., Okogbo, F., Okonto, P.I. & Ohihoin, A.G. (2010).

- Eclampsia in Irrua apecialist teaching hospital: a five-year review. *Niger J Clin Pract*, 13(2), 149-53. Retrieved From: PubMed
10. Akinola, O.I., Fabanwa, A.O., Gbagesin, A., Ottun, T.A. & Kusemiju OA (2008). Improving the critical outcome in cases of eclampsia: The experience t lagos state university teaching hospital, ikeja. *The Internal Journal of Third World Medicine*, 6(2), 2.
  11. Simon, E., Caille, A., Perrotin, F. & Giraudeau, B. (2013). Mixing Nulliparous and multiparous women in randomized control trials of preeclampsia prevention is debatable: evidence from a systemetic review. *PLoS ONE*, 8(6): e66677. doi: 10.1371/journal.pone.0066677
  12. Ozumbia, B.C., Ibe, A.I. (1993). Eclampsia in Enugu, estern Nigeria. *Acta Obstet Gynecol Scand*, 72(3), 189-92. Retrieved from: PubMed
  13. Oladokun, A., Okewole, A.I., Adewole I.F. & Babarinsa, I.A. (2000). Evaluation of cases of eclampsia in university college Hospital Ibadan over a 10 year period. *West A frJ Med*, 19(3), 192-4. Retrieved from: <https://www.ncbi.nlm.nih.gov/pubmed/11126082>
  14. Abate, M.M., Lakew, Z. (2006). Eclampsia. A 5 years retrospective review of 216 cases managed in 2 teaching hospitals in addis abada. *Ethiop Med j*, 44(1), 27-31. Retrieved from: PubMed
  15. Zwart, J.J., Richters, A., Ory, F., de Vries, J.I., Bloemenkamp, K.W. & Van, R.J. (2008). Eclampsia in Netherlands. *Obstet Gynaecol*, 112(4):820-7. doi: 10.1097/AOG.0b013e3181875eb3
  16. Onuh, S.O., Aisien, A.O. (2004). Maternal and fetal outcome in eclamptic patients in Bebin city, Nigeria. *J Obstet Gynaecol*, 24(7), 765-8. doi: 10.1080/01443610400009451
  17. Noraihan, M.N., Sharda, P., Jammal, A.B. (2005). Report of %0 cases of eclampsia. *J Obstet Gynaecol Res*. 31 (4), 302-9.
  18. Sibai, B.M. (2005). Diagnosis, prevention and management of eclampsia. *Obstet Gynaecol*, 105(2), 402-10. doi: 10.1097/01.AOG.0000152351.13671.99
  19. Ndaboine, E.M., Kihunrwa, A., Rumanyika, R., Im, H.B., Massinde, A.N. (2012). Maternal and perinatal outcomes among eclamptic patients admitted to bugando medical centre, mwanza, Tanzania. *Afr J Reprod Health*, 16(1), 35-41. Retrieved from: PubMed
  20. Singhal, S.R., Deepika, Anshu, Nanda, S. (2009). Maternal and perinatal outcome in severe preeclampsia and eclampsia. *Journal of South Asian Federation of Obstetrics and Gynaecology*. 1 (3), 25-28. Retrieved from: [www.jaypeejournals.com/eJournals/ShowText.aspx?ID=387&Type=FREE&TY=TOP&IN=\\_eJournals/images/JPLGO.gif&IID=41&isPDF=YES](http://www.jaypeejournals.com/eJournals/ShowText.aspx?ID=387&Type=FREE&TY=TOP&IN=_eJournals/images/JPLGO.gif&IID=41&isPDF=YES) Maternal and perinatal outcome in severe preeclampsia and eclampsia.
  21. Duley, L. (2009). Global impact of preeclampsia and eclampsia. *Semin perinatal*, 33(3), 130-7. doi: 10.1053/j.semperi.2009.02.010
  22. Onwuhafua, P.I., Onwuhafua, A., Adze, J., Mairami, Z. (2001) Eclampsia in Kaduna state of Nigeria. A proposal for better outcome. *Niger J Med*, 10(2), 81-4. Retrieved from: PubMed
  23. Endeshaw, G., Berhan, Y. (2015). Perinatal outcome in women with hypertensive disorders of pregnancy: a retrospective cohort study. *International Scholarly Research Notices*, 2015, 1-8. doi: <http://dx.doi.org/10.1155/2015/208043>
  24. Urassa, D.P., Carlstedt, A., Nystrom, L., Massawe, S.N. & Lindmark, G. (2006). Eclampsia in Dar es Salaam, Tanzania-incidence, outcome, and the role of antenatal care. *Acta Obstet Gynecol Scand*, 85(5), 571-8. doi: 10.1080/00016340600604880
  25. Sibai, B.M. (1990). Eclampsia. VI. maternal-perinatal outcome in 254 consecutive cases. *American Journal of Obstetrics and Gynecology*, 163 (3), 1049-1054. doi: [https://doi.org/10.1016/0002-9378\(90\)91123-T](https://doi.org/10.1016/0002-9378(90)91123-T).



GLOBAL JOURNAL OF MEDICAL RESEARCH: E  
GYNECOLOGY AND OBSTETRICS  
Volume 17 Issue 2 Version 1.0 Year 2017  
Type: Double Blind Peer Reviewed International Research Journal  
Publisher: Global Journals Inc. (USA)  
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

## Successful Outcome of Pregnancy after Cystectomy for Twisted Ovarian Cyst : A Case Report

By Dr. Amrita Jain & Dr. Rekha Daver

*Grant Government Medical College*

**Abstract- Aim & Background:** Though ovarian masses during pregnancy are relatively common, its complication can pose a significant risk to pregnant woman and her fetus.

Two most common complications of ovarian mass are torsion and hemorrhage. Ovarian torsion encountered during pregnancy carries a significant risk to pregnant woman and intrauterine foetus. We present an interesting case of complication of an ovarian cyst in 2nd trimester of pregnancy and how it was managed with a successful antenatal outcome.

**Case Description:** In this case we report a 28 -year-old second gravida with 26 weeks of pregnancy presenting with torsion of the right ovarian cyst. She presented to the emergency with acute pain in abdomen and vomiting. She was diagnosed to have torsion of ovarian cyst with 26 wks live pregnancy. Emergency exploratory laparotomy followed by right ovarian cystectomy was done. Histopathology report showed a haemorrhagic ovarian cyst. Her pregnancy was followed up and was uneventful.

**Keywords:** ovarian cyst torsion, pregnancy, corpus luteum cyst.

**GJMR-E Classification:** NLMC Code: WQ 200



*Strictly as per the compliance and regulations of:*



© 2017. Dr. Amrita Jain & Dr. Rekha Daver. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License (<http://creativecommons.org/licenses/by-nc/3.0/>), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

# Successful Outcome of Pregnancy after Cystectomy for Twisted Ovarian Cyst : A Case Report

Dr. Amrita Jain <sup>α</sup> & Dr. Rekha Daver <sup>ο</sup>

**Abstract- Aim & Background:** Though ovarian masses during pregnancy are relatively common, its complication can pose a significant risk to pregnant woman and her fetus.

Two most common complications of ovarian mass are torsion and hemorrhage. Ovarian torsion encountered during pregnancy carries a significant risk to pregnant woman and intrauterine foetus. We present an interesting case of complication of an ovarian cyst in 2nd trimester of pregnancy and how it was managed with a successful antenatal outcome.

**Case Description:** In this case we report a 28 -year-old second gravida with 26 weeks of pregnancy presenting with torsion of the right ovarian cyst. She presented to the emergency with acute pain in abdomen and vomiting. She was diagnosed to have torsion of ovarian cyst with 26 wks live pregnancy. Emergency exploratory laparotomy followed by right ovarian cystectomy was done. Histopathology report showed a haemorrhagic ovarian cyst. Her pregnancy was followed up and was uneventful.

**Conclusion:** Ovarian torsion in second trimester of pregnancy, though is an extremely rare problem in pregnancy, adnexal torsion should be taken into consideration in the differential diagnosis of acute abdominal pain and urgent Ultrasound aids in diagnosis.

**Keywords:** ovarian cyst torsion, pregnancy, corpus luteum cyst.

## CASE REPORT

### I. BACKGROUND

The incidence of adnexal masses in pregnancy ranges from 1 in 81 to 1 in 8000 pregnancies<sup>1</sup>. The most frequent types of ovarian masses in pregnancy are corpus luteum cysts, endometriomas, benign cystadenomas, and mature cystic teratomas called dermoids. Because pregnant women are usually young, malignant tumors and those of low malignant potential are proportionately uncommon and vary from 4 to 13 percent<sup>2</sup>.

Most ovarian masses are asymptomatic in pregnant women. Some cause pressure or chronic pain however acute abdominal pain may be due to torsion, rupture, or hemorrhage. Torsion of ovary is the total or partial rotation of the adnexa around its vascular axis or pedicle. Minimal or early twisting may compromise only venous flow, thus leaving arterial supply intact.

*Author α:* Asst Prof, Grant Government Medical College, Mumbai.

*e-mail:* amritadby1901@gmail.com

*Author ο:* (Prof & HOD).

Complete torsion causes total blockade of venous and lymphatic supply that leads to venous congestion, haemorrhage and necrosis, subsequently cyst becomes tense and may rupture.

The risk of ovarian torsion rises by 5 fold during pregnancy. Incidence is 5 per 10,000 pregnancies<sup>3,4</sup>. Torsion of ovarian tumors occurs predominantly in the reproductive age group. The majority of the cases presented in pregnant (22.7%) than in non-pregnant (6.1%) women<sup>5</sup>.

Patient usually presents with acute pain in lower abdomen that frequently is accompanied by nausea and vomiting. Pelvic examination may reveal a tender cystic mass separate from the uterus.

Sonography aids in diagnosis and with Color Doppler, presence of ovarian mass with absent flow strongly correlates with torsion. Here, we report a case of torsion of ovarian cyst during second trimester of pregnancy with no known predisposing factors.

### II. CASE DESCRIPTION

A 28 year old second gravida who delivered her first child vaginally 2 years back presented in the emergency on 26/8/2016 with history of 6 months pregnancy with dull aching pain since 1 month & acute pain in abdomen and vomiting for one day. Pt gave history of abdominal pain since 1 month for which pt had shown in private where she was started on tocolytic agents.

Now she had acute pain since morning associated with nausea. On examination she was afebrile with pulse of 100/min and BP-120/80 mm Hg.

On examination her fundal height was corresponding to 26 weeks of gestation with Fetal heart sounds 150/m. Per abdomen tenderness was present in right iliac fossa region. There was no guarding/rigidity. Pelvic examination revealed cervical motion tenderness and right adnexal fullness.

Urgent USG done showed a single viable fetus of 26 weeks maturity and a septated cyst (10.6 ×8.4) cm in right lumbar region extending upto umbilicus probably originating from right ovary with absent flow. All other investigation reports were within normal limit. Patient was counselled about the risk of preterm labor

and informed consent for exploratory laparotomy with possible need for salpingoophorectomy taken.

During laparotomy (10 ×8) cm black gangrenous cyst was found in the right adnexa which was twisted around its pedicle 3 times (Figure 1 & 2).

Right fallopian tube was found gangrenous, attached to the cyst. No free intraperitoneal fluid was found. Uterus was 26-28 weeks in size. Left sided fallopian tube and ovary were healthy. Right salpingoophorectomy was done without untwisting its pedicle.



Figure 1: Twisted rt ovarian cyst with 26 wks pregnant uterus.



Figure 2: Specimen of 10 x 8 cm twisted rt ovarian cyst.

Post op she was started on injection Duvadilan. Her post operative period was uneventful and she was discharged on 7th postoperative day.

Histopathological examination of the specimen showed haemorrhagic ovarian cyst.

She was followed up in the antenatal clinic where rest of her antenatal period was essentially uneventful. Her follow up USG scans showed normal growth and development of the fetus.

On 11/12 2016, she was admitted at 40 wks in prelabour and she delivered vaginally 2.9 kg male baby.

### III. DISCUSSION

The management of ovarian tumors during pregnancy can be challenging because of the risk of fetal wastage and the possibility of surgery-related complications, or a delayed diagnosis of a possibly lethal disease or malignancy. Ovarian masses should be managed conservatively, because the majority of

ovarian tumors in pregnant women spontaneously disappear during follow up<sup>6</sup>.

Most ovarian masses are detected during routine prenatal sonography or during imaging done for other indications. The typical sonographic appearance of these masses are as - A simple anechoic cyst with smooth wall is characteristic of a physiological corpus luteum cyst or benign cystadenoma, Cystic structure with diffuse internal low level echoes suggestive of an endometrioma or hemorrhagic corpus luteum cyst. In some instance MR imaging can be used to evaluate complicated anatomy<sup>2</sup>. CA 125 levels are frequently elevated with ovarian malignancy.

If a cystic benign appearing mass is <5cm, it often requires no additional antepartum surveillance. Tumor between 5 and 10 cm should be carefully evaluated by sonography along with color Doppler and possibly MR imaging. For cyst >10cm, because of substantial risk of malignancy, torsion or labor obstruction surgical removal is reasonable.

Complications of the ovarian cysts in pregnancy are torsion of the cyst, rupture, infection, malignancy, impaction of cyst in pelvis causing retention of urine, malpresentation of foetus and during labour obstructed labour<sup>6</sup>. Early diagnosis is essential as it makes a conservative approach possible. When diagnosis is made earlier, simple detorsion is possible with good functional results.

The diagnosis of twisted ovarian cyst, which is an acute abdominal emergency, can be made in the majority of cases, although the symptoms are nonspecific for ovarian torsion. The usual symptoms of torsion of an ovarian cyst are acute, colicky pain in the lower abdomen, with vomiting and tenderness. If torsion is suspected, laparoscopy or laparotomy is warranted. If the adnexa is healthy, there are two options. First, neoplasms are resected, ovarian cystectomy done if ischemic and oedematous ovary as it may be technically difficult, and adnexectomy may be necessary. Second unilateral or bilateral oophorectomy has been described to minimize the risk of repeated torsion<sup>2</sup>.

Evacuation by needle aspiration is not recommended in pregnancy<sup>7</sup>.

Differential diagnosis of ovarian cyst torsion in pregnancy includes: uterine leiomyomas, non pregnant horn of bicornuate uterus, appendiceal abscess, diverticular abscess, pelvic kidney, retroperitoneal tumours, ectopic pregnancy and retroverted gravid uterus<sup>5</sup>.

#### IV. CONCLUSION

The management of ovarian cyst in pregnancy is usually conservative with serial ultrasound monitoring. The nature of an ovarian cyst should be studied in detail and malignancy has to be ruled out. Ovarian torsion in second trimester of pregnancy, though is an extremely rare problem in pregnancy, adnexal torsion should be taken into consideration in the differential diagnosis of abdominal pain. Treatment options are limited to surgery, either by laparoscopy or laparotomy. In our case we performed a laparotomy with Pfannenstiel incision, and did not attempt to untwist the adnexa because of widespread necrosis.

#### *Clinical Significance*

Most common cause of ovarian cyst in pregnancy is corpus luteum cyst which regresses by 12 to 16 weeks<sup>8</sup>. Ovarian torsion, therefore, occurs most frequently in the first trimester, occasionally in second, and rarely in third<sup>8</sup>. If the ovarian cyst is diagnosed in the first trimester, it is better to wait till 16 wks when the implantation of pregnancy is more secure and also the cyst may disappear spontaneously. Persisting tumours are treated by cystectomy or ovariectomy as indicated. Ovarian tumour or cyst can be easily removed till 28 wks of gestation thereafter not only it becomes hard to

access but also operation may precipitate preterm labour.

Previously untwisting of the pedicle was avoided to prevent emboli and toxic substances related to hypoxia, from entering peripheral circulation, but recently, re-establishing ovarian circulation by untwisting, has shown to improve circulation in viable ovarian tissue with no systemic complications<sup>3,4</sup>. Obviously in cases where ovarian cysts have undergone gangrenous changes, untwisting are not tried, which was present in the present case. Thus each case has to be individualized, and the management should be done accordingly.

#### REFERENCES RÉFÉRENCES REFERENCIAS

1. Whitecar MP, Turner S, Higby MK. Adnexal masses in pregnancy: a review of 130 cases undergoing surgical management. *Am J Obstet Gynecol* 1999; 181: 19-24.
2. William obs/ 24th Edition/chapter 63/pg 1226-29.
3. Kolluru V, Gurumurthy R, Vellanki V, Gururaj D. Torsion of ovarian cyst during pregnancy: a case report. *Cases Journal* 2009;2:9405. <http://dx.doi.org/10.1186/1757-1626-2-9405> PMID: 20090873 PMCID: 2809077
4. Ventolini G, Hunter L, Drollinger D, Hurd WW: Ovarian torsion during pregnancy. [http://www.residentandstaff.com/issues/articles/2005-09\\_04.asp](http://www.residentandstaff.com/issues/articles/2005-09_04.asp)].
5. Lee CH, Raman S, Sivanesaratnam V: Torsion of ovarian tumors: a clinicopathological study. *Int J Gynaecol Obstet* 1989, 28: 21-25.
6. Wang PH, Chang WH, Cheng MH, Horng HC. Management of adnexal masses during pregnancy. *J Obstet Gynaecol Res* 2009; 35: 597-8.
7. Goffinet F. Ovarian cysts and pregnancy. *J Gynecol Obstet Biol Reprod (Paris)* 2001; 30(1 Suppl): S100-8.
8. Hibbard LT: Adnexal torsion. *Am J Obstet Gynecol* 1985,152: 456-461.





This page is intentionally left blank



## Biochemical Markers and Uterine Artery Doppler Study for the Prediction and the Severity of the Hypertensive Disorders during Pregnancy

By Saptarshi Chakraborty, Veena Acharya & Surabhi Saharan

*Mahatma Gandhi Medical College & Hospital*

**Abstract-** Hypertensive disorder in pregnancy (HDP) is one of the major cause of maternal and perinatal mortality & morbidity worldwide particularly in developing countries. In developed countries, maternal mortality rate varies from 4-40 per 1 lakh live birth. In developing countries, it varies from 100-700 with India having 178 per 1 lakh live birth.

**Objective:** Study of biochemical markers and uterine artery Doppler for the prediction of hypertensive disorders and its severity. Sensitivity and Specificity of biochemical markers and Uterine Artery Doppler and their comparison for the prediction and the severity of hypertensive disorders during pregnancy.

**Material & Methods:** This is a prospective study. Approximately hundred patients with hypertensive disorders during pregnancy attending the OPD (Out patients Department) and IPD (In patient Department) in Obstetrics & Gynecology department will be included.

**Keywords:** hypertension, hypertensive disorder in pregnancy, biochemical markers, uterine artery doppler.

**GJMR-E Classification:** NLMC Code: WQ 270



*Strictly as per the compliance and regulations of:*



# Biochemical Markers and Uterine Artery Doppler Study for the Prediction and the Severity of the Hypertensive Disorders during Pregnancy

Saptarshi Chakraborty <sup>α</sup>, Veena Acharya <sup>σ</sup> & Surabhi Saharan <sup>ρ</sup>

**Abstract-** Hypertensive disorder in pregnancy (HDP) is one of the major cause of maternal and perinatal mortality & morbidity worldwide particularly in developing countries. In developed countries, maternal mortality rate varies from 4-40 per 1 lakh live birth. In developing countries, it varies from 100-700 with India having 178 per 1 lakh live birth.

**Objective:** Study of biochemical markers and uterine artery Doppler for the prediction of hypertensive disorders and its severity. Sensitivity and Specificity of biochemical markers and Uterine Artery Doppler and their comparison for the prediction and the severity of hypertensive disorders during pregnancy.

**Material & Methods:** This is a prospective study. Approximately hundred patients with hypertensive disorders during pregnancy attending the OPD (Out patients Department) and IPD (In patient Department) in Obstetrics & Gynecology department will be included.

All the patients in this study group will be subjected to biochemical markers tests and Ultrasonographic evaluation of the pregnancy along with the Arterial Doppler of both the Uterine Artery and Umbilical Artery will be done.

**Conclusion:** Using biochemical markers (Inhibin-A and PAPP-A) and Uterine Artery Doppler Study in combination is significantly useful in early prediction of PIH having specificity and sensitivity of Inhibin-A as 88.89% and 83.33%, PAPP-A as 89.29% and 71.43%, Uterine artery Doppler study – Pi Index as 91.67% and 85.71 %, Ri Index as 87.5% and 71.43% and diastolic notch as 94.44% and 92.85 % respectively. The use of biochemical markers and uterine artery Doppler Study as an important tool for early prediction of PIH and has a lot of prognostic value.

**Keywords:** hypertension, hypertensive disorder in pregnancy, biochemical markers, uterine artery doppler.

## I. INTRODUCTION

Hypertensive disorders of pregnancy are one of the major cause of maternal and perinatal mortality and morbidity worldwide particularly in developing countries. Hypertensive disorder of pregnancy is a sign of an underlying pathology which may be pre existing or appear for the first time during pregnancy. The identification of this clinical problem & effective management plays a significant role in the prevention of the adverse effects on pregnancy outcome [1].

Hypertensive Disorders of Pregnancy is a multi-system disorder, which is characterized by new onset hypertension (systolic and diastolic blood pressure of  $\geq 140$  and  $90$  mm Hg, respectively, on two occasions, at least 6 hours apart) and proteinuria (protein excretion of  $\geq 300$  mg in a 24 h urine collection, or a dipstick of  $\geq 2+$ ), that develop after 20 weeks of gestation in previously Normotensive women [2].

Hypertensive Disorders of Pregnancy can have an early onset (preeclampsia starting before 34 weeks of gestation) or late onset (preeclampsia starting after 34 weeks of gestation), can show mild or severe symptoms (systolic blood pressure  $\geq 160$  mmHg or diastolic blood pressure  $\geq 110$  mmHg, proteinuria  $>5$  g/24 hours, oliguria, neurological symptoms, other clinical symptoms such as deranged liver function, thrombocytopenia  $< 100\ 000$  mm<sup>3</sup>, HELLP syndrome), and can evolve in eclampsia in the most severe cases whereas Eclampsia is the occurrence of one or more convulsions superimposed on the syndrome of pre-eclampsia [3].

Dependent on the systemic involvement along with several other symptoms, such as edema, disturbance of hemostasis, renal or liver failure, and the HELLP syndrome (hemolysis, elevated liver enzymes, and low platelet counts) also complicates the clinical picture.

Most theories on the etiology of hypertensive disorders of pregnancy suggests that the disease is a cascade triggered by combination of abnormal maternal inflammatory response, endothelial cell activation/damage with deranged hemodynamic milieu, and deranged immunity [4,5].

Numerous patho-physiological mechanisms, alone or in combination, have been suggested to be responsible for the diverse subsets of hypertensive disorders of pregnancy. They include impaired vascular remodeling of the maternal-fetal interface, excessive immune response to paternal antigens, systemic inflammatory response, and dysfunctional placental or endothelial response, all of these processes being modulated by genetic and environmental parameters. Such heterogeneity of potential processes leading to, or resulting from, hypertensive disorders of pregnancy has contributed to the lack of diagnostic means for identification of women susceptible to developing pre-

Author <sup>α</sup> <sup>σ</sup> <sup>ρ</sup>: Department of Obstetrics & Gynaecology, Mahatma Gandhi Medical College & Hospital, Sitapura, Jaipur, Rajasthan, India.  
e-mail: saptarshichakraborty8@gmail.com

eclampsia, resulting in delayed recognition and severe complications and impeding evaluation of new preventive interventions [6].

Since there is an involvement of various organ systems, the potential markers would include inhibin A, PAPP-A, Activin A, angiotensin 11 sensitivity, urine albumin excretion, uric acid and microproteinuria, urinary inhibin-a, urinary kallikrein –calcium/kallikrein ratio, maternal serum alpha fetoprotein; coagulation and platelet activation [7].

*Risk factors for preeclampsia:* Include nulliparity, multifetal gestations, previous history of preeclampsia, obesity, diabetes mellitus, vascular and connective tissue disorders like systemic lupus erythematosus and antiphospholipid antibodies, age >35 years at first pregnancy, smoking, and African American race. Ultrasonography Color Doppler study may also help in the prediction of hypertensive disorder. Persistence of high impedance to blood flow in the uterine arteries of women with hypertension, which is one of the indirect evidence of abnormal placentation also helps in early identification of disease[8]. The changes are increased systolic to diastolic ratio, absent diastolic flow, reverse diastolic flow depending upon severity of the disease[9]. With the help of Uterine Artery Doppler Velocity waveform analysis we can measure resistance index, pulsatility index, systolic/diastolic ratio, diastolic/systolic ratio, presence of any diastolic notch [10].

The specificity and sensitivity of the Uterine Artery Color Doppler Study for PI Index are 91.67% and 85.71%, RI Index 87.5% and 71.43% and for Diastolic Notch are 94.44% and 92.86% which is very useful in early detection of PIH [11]. Therefore, Ultrasonography Color Doppler study may also help in the prediction of hypertensive disorder. Persistence of high impedance to blood flow in the uterine arteries of women with hypertension, which is one of the indirect evidence of abnormal placentation also helps in early identification of disease[12]. The changes are increased systolic to diastolic ratio, absent diastolic flow, reverse diastolic flow depending upon severity of the disease[13,14].

WHO, UNFPA, UNICEF, IPPFF, the population council and other national & international agencies concerned with safe motherhood concluded that it is possible to reduce maternal mortality significantly with investigation and its effective management [15].

Reducing maternal mortality by 75% between 1990 and 2015 has been considered as part of the millennium development goals of the World Health Organization (WHO) Nations [15].

## II. MATERIALS AND METHODS

This is a prospective study done at Mahatma Gandhi Medical College & Hospital, Sitapura, Jaipur. Approximately 100 patients, during pregnancy attending the OPD and IPD in Obst. & Gynae department will be

included. Patient's detailed clinical history, personal history, significant medical history, obstetric history and menstrual history will be taken.

General examination of all the patients will be done and Pulse, Blood Pressure, Temperature and Respiratory Rate will be noted.

Systemic examination including heart, lungs and other systems will be examined in detail.

All the patients in this study group will be subjected to all the routine blood tests and special tests, including biochemical markers. A detailed Ultrasonographic evaluation of the pregnancy along with the Arterial Doppler of both the Uterine Artery and Umbilical Artery will be done. All patients will be kept in the regular follow up and in the end, their maternal and fetal outcome will also be noted.

*Special blood tests will be including:*

- 1) Complete blood count (specially hemoglobin and platelet count),
- 2) Liver function tests including SGOT, SGPT, Alkaline Phosphatase Level, LDL Level and PT/INR.
- 3) Lipid profile – LDL, HDL, Triglyceride, Total Cholesterol, VLDL.

*Biochemical Markers:* Serum Uric Acid, Blood Urea, Serum Creatinine, Activin-A, Pregnancy Associated Plasma Protein-A (PAPP-A)

*Urinary Tests:* Urinary Micro Albumin Level, Urine Protein/Creatinine Ratio, Urinary Calcium Creatinine Ratio.

In the Doppler study following parameters will be assessed:

*Pulsatility Index:* A measure of the variability of the blood velocity in a vessel equal to the difference between the peak systolic and minimum diastolic velocities divided by the mean velocity during the cardiac cycle.

$$\text{Calculated by} = \frac{\text{Peak Systolic Velocity} - \text{End Diastolic Velocity}}{\text{Mean Velocity}}$$

*Resistance Index:* Is a measure of pulsatile blood flow that reflects the resistance to blood flow caused by microvascular bed distal to the site of measurement.

$$\text{Calculated by} = \frac{\text{Peak Systolic Velocity} - \text{End Diastolic Velocity}}{\text{Peak Systolic Velocity}}$$

*Systolic to Diastolic Ratio:* Determinations of blood flow velocities that reflects intrinsic resistance in an arterial blood vessel.

Presence and absence of the diastolic in both the uterine arteries.

Using Statistical test i.e. Z-test where the distribution of the test statistic under the null hypothesis can be approximated by a normal distribution, using expected value  $\theta$  of T under the null hypothesis, and then obtaining an estimate S of the standard deviation of

T, the standard score  $Z = (T - \theta) / s$  is calculated, will be applied wherever needed.

With the help of these statistical methods, the sensitivity and the specificity of the biological markers and the Uterine Artery Doppler study will be calculated.

### III. RESULTS

Pregnancy Induced Hypertension is the most common obstetrical disorder world-wide and on a national scale in India also. It is one of the major causes of maternal and fetal mortality.

**Table 1:** Distribution of the cases according to Age

| Age in Years | No. of patients | Percentage |
|--------------|-----------------|------------|
| 20 -24       | 48              | 48%        |
| 25 – 29      | 27              | 27%        |
| 29- 33       | 17              | 17%        |
| 34 – 38      | 7               | 7%         |
| 39 – 40      | 1               | 1%         |
| Total        | 100             | 100%       |

Table 1 showed the age distribution of 100 patients of my study group that maximum patients 38% were in the age of 20 – 24 years and second most common were in the age group in between 25 – 29 years that was 27%. While only 8 patients were above 35 years.

**Table 2:** Distribution of the cases according to Parity

| Parity | No. of patients | Percentage |
|--------|-----------------|------------|
| Para 1 | 22              | 81 %       |
| Para 2 | 32              | 12 %       |
| Para 3 | 20              | 3%         |
| Para 4 | 18              | 2%         |
| Para 5 | 7               | 2%         |
| Total  | 100             | 100%       |

In this present study of 100 patients, maximum number of patients was primigravida (81%), while multiparous women were 19%.

**Table 3:** Distribution of the cases according to occupation

| Wife Occupation | Number | Percentage |
|-----------------|--------|------------|
| Farmer          | 6      | 6          |
| Housewife       | 84     | 84         |
| Labour          | 1      | 1          |
| Nurse           | 3      | 3          |
| Self Employed   | 1      | 1          |
| Sweeper         | 1      | 1          |
| Teacher         | 4      | 4          |

Out of 100 patients, maximum number of patients 84% was from house wife while 16 had their own jobs (4 were teacher, 3 had nursing job, 6 were working in the field etc.)

**Table 4:** Distribution of the cases according to Residence

| Location | Number | Percentage |
|----------|--------|------------|
| Rural    | 56     | 56         |
| Urban    | 44     | 44         |
| Total    | 100    | 100        |

Out of 100 patients, 56 patients were residing in rural area while 44% in urban area.

**Table 5:** Distribution of the cases according to Socioeconomic Status (Kuppuswami Scale)

| SE. Status   | Number | Percentage |
|--------------|--------|------------|
| Lower Class  | 65     | 65         |
| Middle Class | 31     | 31         |
| Upper Class  | 4      | 4          |
| Total        | 100    | 100        |

In this study of 100 patients, 65 patients were from lower socioeconomic group, while only 4 patients belong to upper class and 34 were from middle class.

*Distribution according to age group:* In my study, 55% were present between 20- 35 years, 7% 34 – 38yrs and 1% 39 – 40yrs.

Same age groups were studied by MANJUSHA SAJITH, VANDANA NIMBARGI et. al. in their study had 1330 patients, out of which 104 were between the age group of 18-32 years.

*Distribution according to Parity:* In my study on the basis of parity showed that maximum number of patients were from the Primigravidae group comprising about 81% of the total patients.

Mary Esien Kooffreh, Mabel Ekott, Dorcas O Ekpoudom in their study on the prevalence of pre-eclampsia among pregnant women including 8524, showed that majority of the patients between the age group of 25-29 years that is about 104 patients were Primigravida.

*Distribution according to Occupation:* In my study out of 100 patients 84% were housewives and 16% were working women.

Swati Singh, Ekele Bissallah Ahmed, Shehu Constance Egundu and Nwobodo Emmanuel Ikechukwu in their study on Hypertensive disorders in pregnancy among pregnant women showed that maximum number of patients that is 88.13% were housewives.

*Distribution according to Residence:* In my study maximum numbers of patients were from rural area i.e.56%.

Shikha Saxena, Prem Chandra Srivastava, K. V. Thimmaraju et. al in their study on Socio-demographic Profile of Pregnancy Induced Hypertension in a Tertiary

Care Centre showed that maximum number of patients that is 77.14% were residing in rural area.

*Distribution of the cases according to Socioeconomic Status:* In my study maximum number of patients i.e. 65% were from lower socioeconomic status.

Parveen M. Aabidha, Anne G. Cherian, Emmanuel Paul and Jasmin Helan in their study on Maternal and fetal outcome in pre-eclampsia in a secondary care hospital in South India showed that maximum number of patients 61% were from lower socioeconomic group.

*Uterine Artery Doppler Study specificity and sensitivity in the prediction of PIH*

It showed that Doppler Study of Uterine Artery depicting 100 patients. In this table out of 100 patients, Pi Index was normal in 66 patients and elevated in 6 patients, out of 72 normotensive patients, whereas in Ri Index 63 patients were normal and 6 elevated, out of 72 normotensive patients and in diastolic notch 68 patients were having absent notch and 4 patients had notch present out of 72 patients who were normotensive on follow-up. Whereas in case of 28 PIH patients, Pi Index was elevated in 24 patients and 4 patients were in normal value, whereas in Ri Index 20 patients were having elevated Ri Index and 8 patients were having normal value and in diastolic notch 26 patients had present notch and 2 had absent. This table showed the specificity and sensitivity of Pi Index, Ri Index and Diastolic Notch as 91.67%, 87.5% and 94.44% and 85.71%, 71.43% and 92.86% respectively along with Pi Index PPV 80%, NPV 94.29% and accuracy of 90%, whereas Ri Index PPV 68.97%, NPV 88.73% and accuracy of 83% and diastolic notch PPV 86.67%, NPV 97.14% and accuracy 94% showing Pi Index is having higher specificity and sensitivity along with other parameters after diastolic notch.

#### IV. CONCLUSION

I from my study infer that using biochemical markers (Inhibin-A and PAPP-A) and Uterine Artery Doppler Study in combination is significantly useful in early prediction of PIH having specificity and sensitivity of Inhibin-A as 88.89% and 83.33%, PAPP-A as 89.29% and 71.43%, Uterine artery Doppler study – Pi Index as 91.67% and 85.71 %, Ri Index as 87.5% and 71.43% and diastolic notch as 94.44% and 92.85 % respectively.

#### V. DISCUSSION

My Study on Early prediction of Hypertensive Disorders of Pregnancy using biochemical markers and uterine artery Doppler study was conducted on 100 pregnant women attending the antenatal clinic of Mahatma Gandhi Medical College, Jaipur. Out of 100 patients, 72 patients were normotensive and 28 patients were of PIH.

In my study group maximum patients were in the age group of 20-29 years comprising 75% and second most common were in the age group in between 29 – 33 years that was 17%, While only 8 patients were above 35 years with mean age 26.7 years.

Maximum number of patients were primigravida (81%), while multiparous women were 19%. Maximum number of patients was housewives 84%, while 16 had their own jobs. 56% patients were from rural area while 44% in urban area. 65% patients were from lower socioeconomic group, while only 4 patients belonged to upper class and 34 were from middle class.

9% patients were between 16-20 weeks, 26% between 20-24 weeks and 65% patients were between 25-29 weeks.

77% showed normal range of blood urea 20-24mg/dl, while 23% showed slightly high level of urea, while 93% patients had slightly raised serum bilirubin, while serum creatinine were normal in all the patients.

Out of 72 normotensive patients, 64 patients had normal level of Inhibin-A 514-890 pg/ml, while 8 patients had elevated level of Inhibin-A (891-1021 pg/ml) and out of 28 PIH patients, 3 patients had an Inhibin-A level within normal range and 25 patients had elevated Inhibin-A level showing the specificity and sensitivity of Inhibin-A as 88.89%, 83.33%, PPV 75.76%, NPV 95.52% and accuracy of 89%.

Out of 72 normotensive patients, 60 patients showed normal level of PAPP-A 0.90-0.60, while 12 patients had decreased level of PAPP-A (0.50-0.40) and out of 28 PIH patients, 8 patients had normal PAPP-A level and 20 patients had decreased PAPP-A level showing the specificity and sensitivity of Inhibin-A as 89.29% and 71.43%, PAPP-A PPV 62.5%, NPV 88.24% and accuracy of 80%.

12. Out of 72 normotensive patients, 66 patients had normal Pi Index while 6 patients had slightly elevated level. Out of 28 PIH patients 24 patients had elevated Pi Index and 4 had normal Pi Index. This result showed the specificity and sensitivity of Pi Index as 91.67% and 85.71 % and having a PPV of 80%, NPV of 94.29% and accuracy of 90%.

Out of 72 patients who remained normotensive, 63 patients had normal Ri Index, while 9 had slight elevated level and out of 28 patients who developed PIH on follow-up, 20 patients showed elevated level of Ri Index, while 8 patients had normal Ri Index, showing the specificity and sensitivity of Ri Index are respectively 87.5% and 71.43%, which is followed by 68.87% of PPV, 88.73% of NPV and accuracy of 83%.

Out of 72 normotensive patients, 57 patients had absent diastolic notch, while 15 patients had diastolic notch present and Out of 28 patients who developed PIH on follow-up, 21 patients had diastolic notch present, while 7 patients had absent diastolic notch showing the specificity and sensitivity of 94.44%

and 92.85 % respectively while PPV was 94.44%, NPV 86.67% and accuracy of 94%.

## VI. ACKNOWLEDGEMENTS

I would like to acknowledge:

### Abbreviations:

HDP –Hypertensive Disorder in Pregnancy,  
OPD – Out Patient Department,  
IPD – In Patient Department,  
PAPP-A – Pregnancy Associated Plasma Protein-A.

## REFERENCES RÉFÉRENCES REFERENCIAS

1. National Journal of Community Medicine: 2014, 5(1),
2. Fisher SJ. The placental problem: Linking abnormal Cytotrophoblast differentiation to the maternal symptoms of Reprod Biol Endocrinol 2004.
3. Steegers EA, von Dadelszen P, Duvekot JJ, Pijnenborg R. Pre-eclampsia. The Lancet. 2010 Aug 27; 376(9741): 631-44.
4. Aloizos S, Seretis C, Liakos N, Aravosita P, Mystakelli C, Kanna E, Gourgiotis S. HELLP syndrome: understanding and management of a pregnancy-specific disease. Journal of Obstetrics and Gynaecology. 2013 May 1; 33(4): 331-7.
5. Mele C, Remuzzi G, Noris M. Hemolytic uremic syndrome. In Seminars in immunopathology, Springer Berlin Heidelberg. 2014, 36, 4, 399-420.
6. Maler J, Manor D, Itskovitz J, et al. Changes in uterine blood flow during human pregnancy. Am J Obstet Gynecol 1990.
7. Conde-Agudelo AG, Romero RO, Lindheimer MD. Tests to predict preeclampsia. Chesley's hypertensive disorders in pregnancy. Amsterdam: Academic Press, Elsevier. 2009 Apr 16:189-211.
8. Jeyabalan A. Epidemiology of preeclampsia: impact of obesity. Nutrition reviews. 2013 Oct 1; 71(suppl 1): S18-25.
9. Trudinger BJ, Giles WB, Cook Cm, Bombardieri J, Collins LE. Fetal umbilical artery flow velocity waveforms and placental resistance: clinical significance. BJOG: An International Journal of Obstetrics & Gynaecology. 1985 Jan 1; 92(1): 23-30.
10. Fay RA, Ellwood DA, Bruce S, Turner A. Color Doppler imaging of the uteroplacental circulation in the middle trimester: observations on the development of a low-resistance circulation. Ultrasound in Obstetrics and Gynecology. 1994 Sep 1; 4(5): 391-5.
11. Harrington KD, Cooper D, Lees C, Hecher K, Campbell S. Doppler ultrasound of the uterine arteries: the importance of bilateral notching in the prediction of pre-eclampsia, placental abruption or delivery of a small-for-gestational-age baby. Ultrasound in Obstetrics and Gynecology. 1996 Mar 1; 7(3): 182-8.
12. Role of color doppler in pregnancy induced hypertension (a study of 100 cases) J Arora, CJ Bhatt, MS Shah Department of Radiology and Imaging, VS General Hospital, Ahmedabad, India, Indian Journal of Radiology and Imaging, 2003, 13(4), 417-420.
13. Jeltsje S. Cnossen, Rachel K. Morris et. al. Use of uterine artery Doppler ultrasonography to predict pre-eclampsia and intrauterine growth restriction: a systematic review and bivariable meta-analysis.
14. Uterine artery Doppler study in second trimester of pregnancy Olufemi Adebari Oloyede, Faye Iketubosin Fetal Medicine Unit, Department of Obstetrics & Gynaecology, Lagos State University Teaching Hospital, Ikeja, Lagos, Nigeria, Assisted Conception Unit, Georges Memorial Medical Centre, Lagos, Nigeria.
15. AbouZahr C. Cautious champions: international agency efforts to get safe motherhood onto the agenda. Safe motherhood strategies: A review of the evidence. 2001.



This page is intentionally left blank







## Awareness of Signs of Obstetric Complications amongst Married Couples: Implications for Decision-Making towards Care-Seeking. A Qualitative Study of Selected Districts in Ghana

By Bougangué Bassoumah & Mpawenimana Abdallah Saidi

*University Malaysia Sarawak*

**Abstract-** This paper examined how couples' awareness of signs of obstetric complications inform intra-household decision-making towards care-seeking. The data were generated from twelve focus groups and twenty-four individual interviews involving women who were receiving maternity care and their husbands in the Yendi Municipality, Awutu-Senya West and Chereponi Districts in Ghana. Quota sampling and snowballing were used to reach the participants. The study classified couples as knowledgeable or less knowledgeable based on their level of awareness of key signs of obstetric complications. Most knowledgeable couples took joint decisions devoid of cultural and gender influences. Their decisions were reached based on their knowledge about signs/symptoms of obstetric complications and their implications for maternal and neonatal health. There were some level of gender equality and female autonomy amongst the knowledgeable in decision-making. However, decisions made by less knowledgeable couples were male-dominated, culture- and gender-driven embedded in health beliefs and traditional gender norms.

**Keywords:** awareness, maternity, signs, intra-household, decision-making, clinic attendance, needs assessment.

**GJMR-E Classification:** NLMC Code: WQ 400



*Strictly as per the compliance and regulations of:*



# Awareness of Signs of Obstetric Complications amongst Married Couples: Implications for Decision-Making towards Care-Seeking. A Qualitative Study of Selected Districts in Ghana

Bougangue Bassoumah <sup>α</sup> & Mpawenimana Abdallah Saidi <sup>σ</sup>

**Abstract-** This paper examined how couples' awareness of signs of obstetric complications inform intra-household decision-making towards care-seeking. The data were generated from twelve focus groups and twenty-four individual interviews involving women who were receiving maternity care and their husbands in the Yendi Municipality, Awutu-Senya West and Chereponi Districts in Ghana. Quota sampling and snowballing were used to reach the participants. The study classified couples as knowledgeable or less knowledgeable based on their level of awareness of key signs of obstetric complications. Most knowledgeable couples took joint decisions devoid of cultural and gender influences. Their decisions were reached based on their knowledge about signs/symptoms of obstetric complications and their implications for maternal and neonatal health. There were some level of gender equality and female autonomy amongst the knowledgeable in decision-making. However, decisions made by less knowledgeable couples were male-dominated, culture- and gender-driven embedded in health beliefs and traditional gender norms. Decision-making by the less knowledgeable was associated with delays due to socio-cultural structures within the household and community. Though decisions by knowledgeable couples showed positive attitudes towards care-seeking as compared with the less knowledgeable, the data indicated that awareness of maternity conditions did not necessarily translate into positive clinic attendance because it may be influenced by cultural beliefs and gender roles to determine the need for clinical care as well as the type of care-seeking decision taken by couples. Healthcare practitioners should consider carrying out need-assessment for maternal counselling messaging for couples. Education on pregnancy and related complications as well as their implications for maternal and neonatal health is essential to facilitate informed decisions and prompt maternal clinic attendance for positive health outcomes.

**Keywords:** awareness, maternity, signs, intra-household, decision-making, clinic attendance, needs assessment.

## 1. BACKGROUND

Studies on maternity have shown that intra-household decision-making is an essential component of maternal healthcare service utilisation (Tsikata, 2007; Babalola & Fatusi, 2009;

*Author α σ: (PhD), Faculty of Social Sciences, University Malaysia Sarawak, Malaysia. e-mails: wepyjadam@gmail.com, mpawenimanaas@gmail.com*

Hagman, 2013; Ghana Statistical Service [GSS], 2008, 2009, 2015; International Centre for Research on Women [ICRW], 2008, 2010; Bougangue, 2017). Informed decision is a necessary requirement to ensure appropriate timing and choice of source of healthcare. However, reaching an informed decision depends on the couple or the care-giver's awareness of signs of obstetric complications as well as their health implications (GSS, 2008; ICRW, 2010). This is because awareness of signs and complications arising from pregnancy and childbirth facilitates early recognition of danger signs/symptoms associated with maternity, which forms the basis of care-seeking decision-making (Waiswa et al., 2010; Comb Thorsen et al., 2012). This requires active involvement of women, for appropriate clinical diagnosis is largely dependent on the information that is given to professionals by the women or their care-givers (WHO, 2002, 2003).

Research has established that, apart from bleeding, most obstetric conditions cannot easily be recognised by non-professionals until the situation aggravates, whilst some other complications occur suddenly at the onset and quickly progress to become life-threatening (WHO, 2003; GSS, 2015). Based on this, the WHO enjoins all member countries to educate women on maternity issues with the view to increasing their knowledge and understanding of pregnancy and childbirth as a pre-condition for adequate preparation and sound decision-making towards pre-natal, intra-partum and post-partum care (WHO, 2002; 2013). Some couples may be aware of key danger signs of obstetric conditions but lack of knowledge and understanding of the implications of these signs for pregnancy outcomes could influence their care-seeking decisions. Sometimes, early signs and symptoms may be ignored by the pregnant woman or her spouse because the signs may be considered as 'normal' (WHO, 2001; Senah, 2003; Okolocha et al., 1998 in WHO, 2003 pg.45).

Female autonomy and joint decision-making are recommended for couples to ensure informed decisions for positive maternal and child health outcomes (ICRW, 2008, 2010; UNFPA/ICRW, 2014). However, this usually



becomes impossible, particularly in patriarchal societies where decision-making is considered as a men's space (Nwokocha, 2007; Bougangue, 2017). This act of gender discrimination, which creates inequality in decision-making is more entrenched in rural communities and amongst people who are not well informed about maternity and its associated complications (Bougangue & Ling, 2017). It is women who carry pregnancy and experience the symptoms of its complications, and, therefore, they, in particular, can best determine the severity of the conditions and the need for healthcare. However, most women are marginalised at the household level in decision-making towards maternity care-seeking (Hagman, 2013; Bougangue, 2017). In some patriarchal communities in Ghana, decision-making is regarded as an act of protection for women, with spirituality attached as a restriction to the females (Bougangue, 2017). This situation compromises quality decision and positive maternal healthcare behaviour towards care-seeking, with an implication for health outcome.

Though some men may engage their spouses in intra-household decision-making, the final decision on care-seeking is usually reached based on the man's own assessment of the severity of the woman's condition and the need for care (Ampim, 2013; Bougangue, 2017). Meanwhile, studies show that some men are not aware of maternity signs and complications, which suggests that men may lack the knowledge to make informed decisions for their spouses. This presupposes that male-dominated decisions are most likely to have far-reaching implications for maternal and neonatal healthcare. Thus, active involvement of women in intra-household decision-making is a crucial strategy for positive maternal outcomes (ICRW, 2008, 2010). Moreover, female autonomy at the household level facilitates quicker decision-making and promotes positive behaviour towards maternity care-seeking, which is a necessary pre-condition for improved maternal and neonatal health (Babalola & Fatusi, 2009; Hagman, 2013).

This paper assessed how couples' awareness of maternity signs and complications influences female spousal involvement in intra-household decision-making towards maternity care-seeking vis-a-vis clinic attendance. The paper provides insight into how healthcare decision-making affects women's health during the pregnancy-postpartum period. This study is motivated by the fact that gender inequality and women's low social status and disempowerment, relative to men, significantly impact women's healthcare decisions, their health and the demand for maternal healthcare services (African Union, 2006; Nwokocha, 2007; Bougangue & Ling, 2017).

## II. METHODOLOGY

### a) Study setting

The Awutu-Senya West and the Chereponi Districts are newly created districts in the Central and Northern Regions of Ghana, with Awutu-Beraku and Chereponi as the administrative capitals respectively. The Yendi Municipality is located in the Northern Region, with Yendi, the paramouncy of Dagban Traditional Area as the capital town. Both the Yendi Municipality and the Chereponi District are Muslim-dominated settings, whilst the Awutu-Senya West District is dominated by Christians. The Dagombas, from the Mole-Dagbani ethnic group, form the largest proportion in the Yendi Municipality whilst the Awutus and the Chokosis, from the Guan and Akan ethnic groups, are the main ethnic groups in the Awutu-Senya and Chereponi districts respectively (Awutu-Senya West District Assembly, 2015; Yendi Municipal Assembly, 2015; Chereponi District Assembly, 2015). Most of the households in Yendi Municipality and the Chereponi District are headed by men. The compound house system forms the predominant housing system in these areas, where couples live together with members of the extended family, usually relatives of the husbands (Chereponi District Assembly, 2015; Yendi Municipal Assembly, 2015).

### b) Research design

The study adopted a qualitative design, using culturally appropriate methods in the data collection and analysis. This design was adopted to provide space for married couples to share their experiences/knowledge about signs of pregnancy/labour-related complications through focus group and individual interviews. The questions used in this study to assess the awareness of couples of the complications related to pregnancy and childbirth were adapted from the 2008 Ghana Demographic and Health Survey and the 2007 Ghana Maternal Health Survey. Spouses were made to identify pregnancy-related signs/symptoms, complications and their implications individually. Couples who exhibited high level of awareness were categorised as *knowledgeable* whilst those with low level of awareness were classified as *less knowledgeable*. The research organised separate focus group interviews for the two categories of couples. Also, some of the spouses from both the knowledgeable and less knowledgeable categories of couples were individually interviewed.

### c) Sampling and data collection

The data were solely collected by the lead author who was the principal researcher in this study. Women who were receiving maternity care between August 2015 to December 2015 and their spouses were purposively selected for the study and reached through quota sampling and snowballing. Based on thematic data saturation, twelve FGDs and twenty-four IDIs were

conducted in the three settings. The districts were put into urban and rural zones. Two FGDs and four IDIs involving *knowledgeable* and *less knowledgeable* couples were conducted in each zone in each district. The use of the IDI was to validate the FGD data by providing privacy to participants to some extent for them to express themselves freely and candidly (Milena et al., 2008). This allowed the research to delve deep into very subtle issues that could not have been possible with the FGD. In each setting and zone, the knowledgeable were separated from *less knowledgeable* for the FGDs.

d) *Ethical consideration*

Following the Ghanaian custom and research ethics, permission was sought from the chiefs and assembly members of the sampled communities, family heads, as well as the participants before data collection. The purpose and importance of the research outcome were explained to both the community leaders and participants. In addition, each participant was given consent form to sign or thumb-print, and was assured of anonymity and confidentiality before the interview sessions. The participants' consent was also sought for publication of the outcome of the study in both electronic and print format.

e) *Data analysis*

For analysis, expansion of field notes and transcription of recorded data were done within 24 hours. This guided subsequent data collection where the initial analysis suggested change of questioning, and the participants were easily contacted for clarity in cases where important information was missing. To ensure trustworthiness and dependability, the research employed strategies such as combining FGD with IDI, reflexivity during the preparation of the research design and questions, the data collection process and in the various stages of analysis. Also, iterative questioning was employed to ensure that deliberate lies were uncovered during the interactions.

Inductive thematic analysis was the major framework for analysing the data. The data were recorded with a tape recorder during the FGDs and IDIs sessions. Inter-coder analysis was employed, using the results of NVIVO software application and manual coding system. Final coding was done by judging the outcomes of the two techniques. Similar thoughts experienced across the participants were identified, coded and grouped. Out of each group of similar thoughts, a unifying concept or underlying theme was derived. Key points, phrases, and illustrations were also identified to back up the findings, whilst similar emerging themes were grouped together to form major themes.

### III. RESULTS

a) *Awareness of Signs/Symptoms of Obstetric Complications*

i. *The Knowledgeable*

Couples who were aware of maternity and its related complications were able to identify critical periods in pregnancy and complications, such as bleeding, severe waist pain/abdominal pain, persistent severe vomiting (hyperemesis gravidarum) leading to weight loss and dehydration, severe morning sickness and general body weakness, with a few of them mentioning symptoms of pre-eclampsia such as blur vision, frequent headaches and fatigue. In a response to a question on signs of pregnancy, a 28-year old SHS graduate pregnant woman in her first pregnancy in an urban community explained this in IDI:

*"I think nausea, enlargement of breast and the tomy, vomiting and feeling sleeping always are all signs but the seriousness of a particular sign depends on the individual. Mine started with sleeping and I did not realise until I missed my menses before I went to hospital and they told me I was pregnant. I don't vomit and I eat any food without problem but I know some women find it difficult to eat especially, in their first three months in pregnancy."*

The husband had this to say about critical period in IDI:

*"To me, I will say the first 3 months is normally a month women and their spouses need to be very careful. They say the baby is very delicate at this stage. So, any small thing can terminate the pregnancy or affect the baby in the womb. I also realised that women who normally die from pregnancy and childbirth experience this misfortune either at delivery point or immediately after delivery. So, all these periods are very critical for the woman and the foetus' survival."* (34 years, diploma)

These participants explained that the first trimester is critical since the foetus is very delicate, as it undergoes formation within the womb and can be aborted or malformed if not properly taken care of. They mentioned that miscarriages normally occur within this period or within few days after this period. This statement made by a man in IDI was common in the IDIs and FGDs involving the knowledgeable couples:

*"Normally, before miscarriage some blood will come out of the woman's vagina. In the early part of pregnancy or before 4 months in pregnancy the baby is like blood. So, if something happens, it will come out like blood. Therefore, if the woman doesn't go to see doctor she may lose her unborn baby. She needs to go and check the pregnancy so that they give her treatment. They say until the baby is fully grown any blood from vagina can lead to termination"*

of pregnancy. So, they told us to go to see doctor when we feel strong abdominal pains or we see blood." (Husband, 42 years, JSS).

On signs of labour, below were responses from a husband and a wife in separate interviews:

"When I am getting ready for delivery, I see that water comes from my vagina. I can see that some dirty thing like blood also comes from the vagina. I feel like going to urinate or going to toilet. I get back pains in pregnant but when it is getting to time to born the back pain is seriously." (Wife, 40 years, JSS graduate).

"From my experience in my wife's previous pregnancies, normally some water comes out of her organ. Sometimes too she becomes restless with frequent urination or visiting toilet. When I see that then I know she is about to deliver. Another thing is that there is this dirty fluid from the sex organ thicker than water but not clean." (Husband 46 years, SSS graduate).

The conditions associated with labour, such as labour dystocia, prolonged labour and retention of the placenta were also mentioned by these participants. However, the causes of these conditions were received with mixed ideas inclined to women's physiology and spiritualism. They were also well-informed about postpartum haemorrhage, as well as puerperal and neonatal infections. Others who knew about the physiological causes also added that there is a spiritual aspect to retention of baby/placenta during delivery. Whilst agreeing with their colleagues, some participants also held these views expressed by their colleague in FGD:

"I agree with Agi but sometimes it is not a doctor's issue because some bad spirits can attack the women and make them suffer. They may even die. Even the spirits can hold the placenta so that the woman or the baby will die. Some are as a result of curses from fathers, mothers or ancestors of either of the spouses which requires rituals for solution." (36 years, 2 children, primary school, urban community).

They were also well-informed about postpartum haemorrhage and infections associated with delivery, and they also had adequate knowledge about the need for clinic attendance to stop bleeding or screen both the mother and baby early after birth for infections. For instance, in IDI one woman noted:

"Bleeding after childbirth has killed many women. It is not about the birth cert or for weighing alone that we should attend PNC early as some of the women think. You or your baby may be infected but you or the people around you may not detect except clinical test. I was told by the midwife that after delivery it is important to be screened for and treated

of any infections that might affect the mother or baby." (25 years, social worker, diploma).

Below was the response from her husband in IDI:

"After delivery women have to see the doctor as soon as possible to avoid bleeding because it can kill. It has killed many women in this village. They go for traditional treatment and they die. Doctors say women should go to hospital early after delivery for care to check bleeding and other things." (33 years, farmer, SSS)

#### ii. The Less Knowledgeable

Most of the less knowledgeable were aware of common signs such as nausea/vomiting, cessation of menses and weight gain. The statement below made by a woman in FGD was supported by her colleagues, and similar statements were repeated in other FGDs and IDIs involving the same category of participants.

"To me I only realise that I am pregnant when I see my breast enlarging and my stomach becoming bigger and bigger. I don't agree that stoppage of menses is a sign of pregnancy because sometimes the menses can delay and if you eat food that does not contain enough pepper and ginger you can vomit and that does not mean you are pregnant. I have been vomiting and even my husband vomits as well." (40 years, mother of 4 children, SHS graduate).

This was her husband's reaction in the FGD:

"It is serious. For me to be sure that a woman is pregnant the stomach must be very big. It is women who normally know those things. I think my wife knows better." (47 years, A'Level).

They had limited or no idea about the physiological causes of labour dystocia, prolonged labour and retention of placenta, as well as post-partum haemorrhage (PPH). Below are the response of a 45-year old woman and mother of 3 children and her husband in FGD on obstructed labour and retention of the placenta. These statements were supported by most members of the group.

"Women who suffer before they born may have very big babies or they had sex with different men. Some people too have been cursed by their family gods if they sinned against them. It is the same way for those who experience delays in expulsion of the placenta. If the baby is big what about the placenta? It is because of sin. We have to be faithful in our marriages so that God will have mercy on us." (Wife).

"I know many slim women who delivered very big babies but they did not suffer in giving birth. They didn't also struggle before the placenta came out. I think most women who suffer prolonged labour and delay of placenta expulsion only suffer the consequences of their sins and nothing else." (Husband, 47 years).

Though some of them were informed about early postnatal care for preventing PPH, they were not aware of infections that women and babies contract during and after delivery.

"Yea, when my wife born I take her go to doctor after 2 days so that they can do my baby birth certificate and do weighing for him. Some women in this town die when they give birth some time. They are bleeding and go hospital late and they die." (Husband, 44 years, MSLC)

"Yes, I go for check up when I born. My first born I go after three days and my second born I go after naming ceremony for doctor to do my daughter papers and to do weighing. Oh!! I don't go early because I was not weak and my baby is strong. If you born and you are weak you for go and see doctor so that they give you medicine. If the baby too is weak you take it to the doctor to check and give drugs. Women that are weak after delivery go for doctor early or when the baby is sick when they born. They go for doctor to check the baby. Yes, that is what we do normally." (Wife, 40 years, MSLC).

#### b) Intra-Household Decision-Making

##### i. The Knowledgeable

Generally, amongst most knowledgeable couples, decision-making was the responsibility of both spouses. The men did not exercise their traditional gender powers, but rather gave the women some degree of liberty to take autonomous decisions where necessary without seeking approval from husbands. They considered women as the best people in the right position to make decisions towards maternity care since it is women who directly experience the conditions related to pregnancy and childbirth. These couples also mentioned that, even though some men may equally have knowledge about maternity, women usually attend maternal healthcare promotion programmes and are therefore more exposed to issues of pregnancy and childbirth as compared with men. Decisions taken by the knowledgeable were mostly joint decisions or non-tentative individual decisions taken by the women. There were no evidences of delays in decision-making by the knowledgeable. Most of them chose clinical care for treatment of complications and general check-ups during maternity. Below are responses of a couple in separate IDI sessions:

"I don't have the eyes to see what is in the woman's womb. I can only see the seriousness when my wife tells me. I cannot decide for my pregnant wife regarding where and when she should seek care. It is even better for women to decide for themselves." (Husband, 34 years old, SSS graduate).

"Sometimes I decide with my husband but in most cases I decide when, where and the type of care to seek. My husband always say that I should be better

than him in determining the seriousness of what I experience. One thing is that I go to seek care whenever I need care without delays." (Wife, 28 years, SSS graduate).

##### ii. The less knowledgeable

The case of the less knowledgeable was almost a direct opposite of the knowledgeable. Though there were few traces of joint decision-making between spouses, the husbands were the final decision-makers. Unlike the knowledgeable, most husbands amongst these couples were conservative in exercising traditional gender powers. Women did not have the liberty to make autonomous decisions towards care-seeking. Whatever they decided were tentative and subject to approval by husbands. The statements below were made by a couple in FGD session, supported by the discussants and repeated in other FGD and IDI sessions.

"You see, certain things are done by men and some by women. Would you allow your son or daughter to decide for you? We are supposed to act as fathers to our wives and we must be accountable to any decision or action taken about their welfare including health issues. Decision-making is a sole responsibility of the family head. Yes, I listen to my wife but I make my decisions and implement them with her." (Husband, 42 years old, A' Level Graduate).

"As wives, we normally look up to our husbands to tell us what to do. We can't do anything without their approval. They are mandated to play the role of fathers so we are supposed to wait for them or listen to them. I always tell my husband what I experience then he decides whether I should go to hospital or for alternative care from the traditional practitioners." (Wife, 38 years, JSS graduate).

##### c) Interplay of Awareness, Culture and Gender

The decisions made by the knowledgeable were devoid of cultural and gender influences. In the light of this, most of the wives had the freedom to make autonomous decisions which resulted in quicker decision-making and positive clinic attendance. Amongst the knowledgeable, joint decision making was to grant women the listening ears and help the couple to make informed decisions. However, the less knowledgeable reached decisions based on the judgments of the men. Their delayed decisions were male-dominated, culturally determined and gender driven. Most of the less knowledgeable supported this statement, which run through various interviews.

"It is difficult to allow women to make decisions especially in pregnancy because some conditions will provoke the woman to move to the health facility where she would end her life. Certain pregnancy related conditions can best be treated with spiritual eyes or with herbs. Traditionally, women are not

*supposed to make spiritual consultations and therefore, cannot be part of decision-making that is based on spiritual consultation outcomes."* (Husband, 39 years old).

d) *Background of Couples and Intra-Household Decision-Making*

Apart from level of education and media exposure, which were the major sources of knowledge to the couples, other factors, such as religion, location of participants, age, occupation and experiences of maternity were important in determining the involvement of wives in decision-making towards care-seeking. Most Muslim husbands were more conservative and did not involve their spouses in decision-making, although some of them had adequate knowledge and much experience of maternity from their wives. Most Muslim couples supported this statement which was common across the datasets of IDIs and FGDs:

*"I am the man and head of the family. I have to take family decisions. Women only decide on what to cook for the family. ... She only told me she felt like giving birth. So, I took it upon myself to make a good decision about where to go and the time she should seek care."* (52 years old, Muslim, A'level)

Members of traditional religion were also conservative, delayed decision-making, and denied their wives access to modern maternity care in some cases, as the primary choice of care amongst them was traditional practitioners. However, they were more liberal as compared with the Muslims because a few of them involved their wives in decision-making. Most Christians were more liberal and granted their wives autonomy in making and implementing decisions as compared with the members of traditional religion and Muslims. Irrespective of the level of awareness, urban couples were more liberal and the women were more autonomous. Generally, couples who worked together made joint and quicker decisions as compared with those who worked in different environments or workplaces. Although most of them worked together, farmers were more conservative and their decision-making process and outcome were influenced by gender norms and cultural factors.

Husbands whose wives had several pregnancy experiences were more liberal and their wives had opportunities in making decisions about care-seeking as compared with their counterparts without experiences. The men noted that they were comfortable with such women because they had both knowledge and experience to make informed decisions. Also, husbands with much experience of pregnancy were more democratic, as they sought and used the views of their wives in intra-household decision-making as compared with their colleagues with less experience. Joint decision making was more common amongst the younger couples as compared with older ones. Couples with

younger wives and older husbands exhibited a different tendency. The older husbands treated their wives like children by marginalising them in decision-making.

e) *Patriarchy, Matriarchy and Intra-Household Decision Making*

The study observed that patriarchal and matriarchal communities exhibited differences in decision-making regarding the involvement of women and the type of decisions taken by the couples. Generally, couples from patriarchal society were less gender responsive, less democratic and more conservative, which resulted in male-dominated and culture-driven decisions. Data from the Muslim dominated Yendi Municipality and the Chereponi District of the patriarchal Northern Region of Ghana showed that women had limited opportunities to make autonomous decisions as compared with the women from matriarchal Awutu community. Men from Awutu communities gave women more power and autonomy in taking decisions. The women had the chances of finalising decisions which were not likely to be contested by husbands.

f) *Implication for Maternal and Neonatal Healthcare*

The knowledgeable took prompt and informed decisions for care-seeking, which resulted in early antenatal care attendance, mostly in the first month of the first trimester; timely attendance for institution-supervised delivery and postnatal care within the critical period as well as increased number of ANC and PNC visits. The data indicates that most of the women met the recommended ANC attendance of four visits with at least one visit in each trimester for women without serious risk factors. Most of them made about six visits before delivery. In an FGD a couple noted:

*"My wife made several ANC visits. The visits were timely as well including PNC visits. Because she knew I would not worry her she was able to go to the clinic as early as possible when she felt like giving birth in my absence. I only received a phone call from the clinic and joined her there till she delivered."* (Husband, 48 years, No formal education).

*My husband does not interfere in decisions. He allows me to do what I think is good for me and my unborn baby. I don't need to seek permission before going to the hospital for treatment."* (Wife, 42 years, JSS graduate)

However, apart from delayed decision-making, the less knowledgeable couples took uniformed decisions, which had implications for the health of women and their babies. The poor decisions translated into poor clinic attendance in the form of low ANC visits, late ANC, delivery care and PNC. In separate IDIs, a couple shared their experiences:

*"... from the herbalist then we decided to carry her to the clinic but when we got there the midwife said it*

*was too late so we lost the baby and my wife sustained some injuries. Later, they had to clean her womb." (Husband, 45 years old, MSLC).*

*"When I felt like going to toilet several times, I told my husband to take me to the clinic but he went out and when he returned he gave me some herbs. After some time he took me an herbalist but the baby delayed in coming out. Later, they carried me to the clinic but my baby died in the womb and I sustained injuries." (Wife, 39 years, JSS graduate).*

#### IV. DISCUSSION

The observation of this study concurs with the observations of previous studies that pregnancy is usually associated with complications that require proper care from the onset to the post-partum period (Graham, 1998; Stevens, 2000; GSS, 2015). To a large extent, appropriate care-seeking depends on individuals' awareness of signs of obstetric complications for making informed decisions (Pembe et al., 2009; Kabakyenga, Östergren, Turyakira & Pettersson, 2011). This study discovered that awareness of pregnancy and its related complications forms an essential component of intra-household decision-making. Couples' awareness of maternity was instrumental in spousal decision-making towards care-seeking, as the well-informed couples were able to make quicker and informed decisions. The couples' awareness did not only enable the husbands to actively involve their spouses in decision-making with equal opportunity, but also granted women the autonomy to make and implement decisions about their own health. This resulted in timely and regular clinic attendance amongst the knowledgeable which is essential for positive maternal and neonatal health outcomes (Ministry of Health, Uganda cited in Kabakyenga et al., 2011, pg.2; Comb Thorsen et al., 2012; Waiswa et al., 2010). The findings sustain the observation that gender role affects women's autonomy to make decisions about healthcare at the household level which, in turn, play out at the agency level (ICRW, 2008, 2010; WHO Commission on Social Determinants Health, 2008, 2010; Bougangue, 2017).

The findings also confirm previous studies that awareness of danger signs of obstetric complications enables women or the care-givers to take timely and appropriate actions towards healthcare (Pembe et al., 2009; Kabakyenga et al., 2011). The timely and regular clinic visits noticed amongst the knowledgeable positioned them for better chances of maternal screening and early treatment of complications. Most knowledgeable couples were able to stand dominant against gender norms and cultural beliefs to make informed decisions which is a pre-condition for better maternal outcome. The argument that awareness of pregnancy and its related complications translates into

informed decisions and positive attitudes towards care-seeking is maintained in this study (ICRW, 2008, 2010, 2014). As shown in earlier studies, the awareness empowered the couples and guided them to take appropriate decisions during the maternity period (Bhutta, Darmstadt, Hasan & Haws, 2005; Hagman, 2013; Winta, 2013; UNDP/ICRW, 2014). However, women from the less knowledgeable families had higher risks of developing maternal complications or aggravating existing conditions because most of them missed early screening and identification of risks factors for timely intervention due to wrong decisions that led to late clinic attendance (WHO, 2013; GSS, 2008, 2009, 2015).

The observed entrenched gender norms and cultural adherence amongst the less knowledgeable and some of the knowledgeable had a serious repercussion on decision type as well as maternal clinic attendance. This observation supports the findings in previous studies that have shown evidences of gender inequalities in decision-making towards maternal healthcare (Tsikata, 2007; ICRW, 2008, 2010, 2014). The views of women about their own health were central in determining the timing and choice of care type, which guarantees the need for gender equality and women's empowerment in healthcare decision-making at the household level. The marginalisation of women in decision-making and the normalisation of cultural and traditional gender norms as observed in this study had a serious implication for maternal health, particularly amongst the less knowledgeable who entrusted the decision-making power to the sole hands of men. Some husbands did not count women's experience of pregnancy-related symptoms as important for decision-making, and this resulted in some women developing serious complications and injuries during delivery with others losing their lives in the process of giving life.

Self-medication and the use of herbs during pregnancy as this study observed, are associated with health risks in the form of pregnancy termination such as premature birth, spontaneous abortion and stillbirth (Cnatingius et al., 2000; Kirsten, Ulrik, Bodil, Morten, & Tine, 2003; Abasiubong et al., 2012; Liao et al., 2015). The herbs may contain substances such as caffeine, which has adverse effects on the growth and development of the foetus (Bakker et al., 2010; Creanga et al., 2012). There was no guarantee of purity and safety of the herbs used by the women, because they were not subjected to laboratory investigations to ascertain the efficacy, composition, expiry dates as well as the side effects (Lapi et al., 2010). Besides, traditional practitioners lack the skills and equipment to save women's lives in the event of obstructed labour or delay in placenta expulsion, which were evident amongst the less knowledgeable. This exposes the women and their unborn babies to risks of further complications which may result in injuries, lifetime disabilities and deaths



(WHO, 2001, 2013; GSS, 2008; Neilson, Lavender, Quenby & Wray, 2003). As highlighted in the findings, some of the maternal complications, injuries and deaths observed in this study were mainly due to uninformed decision-making that led the women to non-professionals who could not give the needed treatment.

Another finding of this study is that the women's decision-making power about reproduction and sexuality was extremely limited, particularly amongst the less knowledgeable couples. The exclusion of women from decision-making concerning their own health is a violation of women's reproductive rights (Republic of Ghana, 1992; African Union, 2006). Women are the direct objects of complications arising from pregnancy and childbirth, and, therefore, must be given the autonomy to take non-tentative care-seeking decisions. The fundamental human rights guarantee women the right to make decisions and to access healthcare services of their choice (Republic of Ghana, 1992). Studies show that if women are given the power, they make meaningful decisions that can positively affect their lives (Babalola & Fatusi, 2009; Hagman, 2013). However, differences in status between women and men lead to differences in opportunities to claim, benefit from, and enjoy human rights, including the right to decision-making and health (WHO, 2013, 2014).

The interplay of couples' awareness of signs and complications of maternity with gender norms and cultural beliefs influenced the involvement of female spouses in intra-household decisions and the type of decisions they took. Whilst the educational level of couples remains important and necessary for informed decision-making, the study also noticed that couples' exposure to the media as well as their inclination to cultural and gender norms impacted their decisions. Irrespective of awareness, some couples considered decision-making as a sole responsibility of men, and the women relied fully on the decisions made by their spouses for care-seeking. However, despite the cultural, religious and gender influences, the study observed a seeming changing pattern of men's behaviour in intra-household decision-making, with a gradual shift from the hitherto male-centred decision-making to joint and female autonomous decision-making even amongst some illiterate couples. This behavioural pattern was evident, particularly amongst the Christian husbands and the believers of traditional religion. Whilst this is positive and essential for improvement of maternal healthcare (ICRW, 2010, 2014), the norms and principles governing marriage amongst the Muslims, particularly regarding gender ideologies, were observed to be very strong and resilient.

Media exposure had a far-reaching impact, particularly on the illiterates and rural couples. Some of the knowledgeable gained awareness from local radio programmes which were accessible to them even in their farms. The illiterate couples in the urban

communities had more knowledge than their rural counterparts due to exposure to multiple sources of the media and information (different TV and Radio stations/programmes) and the opportunity to interact with literates and experienced people from whom they tapped knowledge.

Spousal age difference emerged as a crucial factor in intra-household decision-making. Men who were married to very young women as compared to their ages considered their wives as incapable of taking appropriate decision. Most of the participants aged between 25 and 35 years who had at least secondary education were in support of joint decision-making and female autonomous decisions. However, regardless of age and experience, female educational attainment was central to women's involvement in decision-making. Female spouses who had at least secondary education were more involved in decision-making as compared with their illiterate counterparts, as well as those who had basic education. This observation maintains the importance of formal education, which is often used as a proxy for knowledge, informed decisions and empowerment, as well as positive behaviour towards care-seeking though there were evidences of exceptional cases (Preston, 1989; McAlister & Baskett, 2006; GSS, 2008, 2009, 2015; Hagman, 2013; Perumal et al., 2013; UNFPA/ICRW, 2013, 2014; Bougangue, 2017; Bougangue & Ling, 2017).

The less knowledgeable women, particularly those from rural areas with much experience of pregnancy and antenatal counselling, were not well-informed about maternity complications, and their implications were observed to be attributable to poor messaging during ANC counselling. This maintains previous discoveries that poor messaging during ANC counselling contributes to low knowledge about pregnancy and childbirth - a recipe for poor maternal health (Perumal et al., 2013). This reinforces the view that dialectical differences and poor communication create barriers to women during antenatal counselling, and these deny women the desired knowledge, thereby leading to lack of perceived need for clinic attendance (Andersen & Newman, 2005; Bougangue & Kumi-Kyereme, 2015).

Most of the couples in the Muslim-dominated patriarchal societies of the Northern Region entrusted decision-making to the hands of men. They were more influenced by gender and cultural norms as compared with their counterparts from the matriarchal Awutu community in the south most of whom gave autonomy to female spouses to take and implement decisions towards care-seeking. In effect, the Awutu female spouses put up better attitudes towards care-seeking as compared with their northern counterparts. This caused late clinic attendance and denial of professional care in some instances, which partly explains the persistent high maternal and child morbidities and mortalities

observed in the Northern Region (GSS, 2008, 2009, 2015).

## V. CONCLUSIONS

The study observed that awareness of signs of obstetric complications as well as their implications is an essential ingredient in reaching well-informed decisions towards care-seeking. However, cultural, religious and gender norms may collide with couples' awareness to adversely influence the involvement of female spouses in decision-making, which has far-reaching implications for health outcome. Adherence to cultural and gender norms, and the associated male-dominant decision-making were acts of disempowerment and marginalisation of women with the tendency of making uninformed decisions and the resultant negative health outcomes.

Healthcare promotion programmes should be directed towards the dangers of cultural, religious and gender norms around maternity care and the need to empower women to make decisions about their own health and that of their babies. Also, there should be public education on pregnancy and childbirth and the associated benefits and dangers to equip couples, especially men, to recognise pregnancy-related complications for informed decision-making.

This study established that couples' awareness of signs of obstetric complications is essential for female spousal involvement in decision-making as well as reaching well-informed decisions for positive maternal outcome.

## REFERENCES RÉFÉRENCES REFERENCIAS

- Abasiubong, F., Bassey, E. A., Udobang, J. A., Akinbami, O. S., Udoh, S. B & Idung A. U (2012). Self-Medication: potential risks and hazards among pregnant women in Uyo, Nigeria. *The Pan African Medical Journal*. Volume 13: 15.
- African Union Commission (2006). *The state of African population report, population dynamics: implications for achieving the MDGs*. Department of Social Affairs: African Union Commission.
- Ampim, A.G (2013). *Men's Involvement in Maternal Healthcare in Accra, Ghana. From Household to Delivery Room*. M. Phil Thesis, Gender and Development: Universitas Bergensis; 2013. Google Scholar
- Andersen, R., & Newman, J. F. (2005). Societal and individual determinants of medical care utilization in the united states. *The Milbank Quarterly*, 83(4), 1-28.
- Awutu-Senya West District Assembly (2015). District profile, Awutu-Breku, CR, Ghana.
- Babalola, S. & Fatusi, A. (2009). Determinants of use of maternal health services in Nigeria Looking beyond individual and household factor. *Pregnancy and Childbirth*, 15(9): 43.
- Bakker, R., Steegers, E. A., Obradov, A., Raat, H., Hofman, A & Jaddoe, V.W.V (2010). Maternal caffeine intake from coffee and tea, fetal growth, and the risk of adverse birth outcomes: the generation R study. *Am J Clin Nutr*, 91 (6) pp. 1691–1698.
- Bhutta, Z. A., Darmstadt, G. L., Hasan, B. S., & Haws, R. A. (2005). Community-based interventions for improving perinatal and neonatal health outcomes in developing countries: a review of the evidence. *Pediatrics*, 115(2 Suppl), 519-617.
- Bougangue, B. & Kumi-Kyereme, A. (2015). Maternal health clinic attendance in Awutu-Senya District, Ghana. *International Journal of Economics & Social Science Volume 5 (2015)*.
- Bougangue, B. (2017). *Socio-cultural dimensions of maternal healthcare in Ghana*. PhD Thesis, Faculty of Social Sciences, Universiti Malaysia Sarawak.
- Chereponi District Assembly (2015). *District profile*, District Planning Office, Chereponi, Northern Region, Ghana.
- Cnatingius, S., Signorello, L. B., Anneren, G., Clausson, B., Ekbohm, A., Ljunger, E et al. (2000). Caffeine intake and the risk of first-trimester spontaneous abortion *N Engl J Med*, 343 (25) pp. 1839–1845.
- Combs Thorsen, V., Sundby, J. & Malata, L. (2012). Piecing together the maternal death puzzle through narratives: The three delays model revisited. *PLOS ONE* 7(12): e52090.doi.io.1371/journal-pone.0052090.
- Creanga, A.A., Sabel, J.C., Ko, J.Y., Wasserman, C.R., Shapiro-Mendoza, C.K., Taylor, P., et al. (2012). Maternal drug use and its effect on neonates: a population-based study in Washington State. *Obstet Gynecol*, 119 (5) pp. 924–933.
- Ghana Statistical Service, Ghana Health Service & Macro International Inc. (2008). *Ghana maternal health survey 2007*. Calverton, Maryland, USA: Ghana Statistical Service, Ghana Health Service, and Macro International Inc.
- Ghana Statistical Service, Ghana Health Service & ICF Macro (2009). *Ghana demographic and health survey 2008*. Calverton, Maryland, USA: Ghana Statistical Service.
- Ghana Statistical Service, Ghana Health Service & ICF International (2015). *Ghana demographic and health survey 2014*. Rockville, Maryland, USA: GSS, GHS, and ICF International.
- Graham, W: (1998). Every pregnancy faces risks. *Plan Parent Chall*. 1998 (1): 13-14.
- Hagman, M. (2013). Maternal mortality: Gender and access to health services. The case of Ghana. *Journal of Politics and International Studies*, Vol.9. ISSN 2047-7651.
- Kabakyenga, J. K., Östergren, P. O, Turyakira, E & Pettersson, K.O (2011). Knowledge of obstetric

- danger signs and birth preparedness practices among women in rural Uganda. *Journal of Reproductive Health*. doi:10.1186/1742-4755-8-33.
21. Kirsten, W., Ulrik, K., Bodil, H. B., Morten, H & Tine, B. H (2003). Maternal consumption of coffee during pregnancy and still birth and infant death in first year of life: prospective study, *BMJ*, 326, pp. 7386–7420.
  22. Lapi, F., Vannacci, A., Moschini, M., Cipollini, F., Morsuillo, M., Gallo, E., et al.(2010). Use, Attitudes and Knowledge of Complementary and Alternative Drugs (CADs) among Pregnant Women: a Preliminary Survey in Tuscany. *PubMed*, eCAM.; 7(4): 477.
  23. Liao, S., Luo, B., Feng, X., Yin, Y., Yang, Y & Jing, W (2015). Substance use and self-medication during pregnancy and associations with socio-demographic data: A cross-sectional survey. *International Journal of Nursing Sciences*. Volume 2, Issue 1, Pages 28–33.
  24. McAlister, C. & Baskett, F. T. (2006). *Female education and maternal Mortality: A worldwide survey*. Department of Obstetrics and Gynaecology, Halifax: Dalhousie University.
  25. Milena, Z. R., Dainora, G., Alin, S. (2008). Qualitative research methods: a comparison between focus group and in-depth interview. *Annals of Faculty of Economics, University of Oradea*, 4 (1)1279-1283.
  26. Neilson, J. P., Lavender, T., Quenby, S & Wray, S. (2003). Obstructed labour: Reducing maternal death and disability during pregnancy. *British Medical Bulletin*, 67, 191-204.
  27. Nwokocho, E. E. (2007). Maternal crises and the role of African men. The case of a Nigerian community. *African Population Studies*, 22(1), 39–62.
  28. Pembe AB, Urassa DP, Carlstedt A, Lindmark G, Nystrom L, Darj E: Rural Tanzanian women's awareness of danger signs of obstetric complications. *BMC Pregnancy Childbirth*. 2009, 9: 12-10.1186/1471-2393-9-12.
  29. Perumal, N., Cole, D. C., Ouédraogo, H. Z., Sindi, K., Loechl, C., Low, J., Levin, C., Kiria, C., Kurji, J & Oyunga, M (2013). Health and nutrition knowledge, attitudes and practices of pregnant women attending and not-attending ANC clinics in Western Kenya: a cross-sectional analysis. *BMC Pregnancy and Childbirth*, 2013 13:146, DOI: 10.1186/1471-2393-13-146.
  30. Preston, S. H. (1989). *Resources, knowledge and child mortality: Selected readings in the cultural and behavioural determinants of health*. A comparison of the US in the late nineteenth century and developing countries today. In: Caldwell, J. C. & Santow, W. G. (Eds.). Canberra: Health Transition Centre.
  31. Republic of Ghana (1992). *The constitution. Abridged*, Accra: NCD.
  32. Senah, K. (2003). Maternal mortality in Ghana: The other side. Research review. *Socio-cultural Dimensions of Reproductive Health and Human Development*. Institute of African Studies, University of Ghana, Legon: New Series, 19(1), 47–56.
  33. Stevens, R.D (2000). Safe motherhood: an insight into maternal mortality in the developing world. *Health Millions*. 2000, 26 (3): 34-37.PubMedGoogle Scholar
  34. Tsikata, D. (2007). Women in Ghana at 50 still struggling to achieve full citizenship? *Ghana Studies*, 10: 163-206.
  35. United Nations Population Fund/International Center for Research on Women (2014). Bridging the gap. The 5th International Conference on Women's Issues in Transportation, Paris.
  36. Waiswa, P., Kallander, K., Peterson, S., Tomson, G. & Pariyo, G. W (2010). Using the three delays model to understand why newborn babies die in eastern Uganda. *Tropical Medicine and International Health*, Volume 15 no. 8 pp 964–972. Division of Global Health, Department of Public Health Sciences, Karolinska Institutet, Stockholm, Sweden.
  37. Winta, N. A (2013). *Knowledge and Practice of Reproductive Health Among Mothers and Their Impact on Fetal Birth Outcomes: A Case of Eritrea* A PhD dissertation, College of Nursing, University of South Florida, USA.
  38. World Health Organisation, United Nations Children's Fund & World Bank (2001). *Making motherhood safer: Overcoming obstacles on the pathway to care*. In Elizabeth I. Ransom & Nancy V. Yinger (Eds). Geneva: World Health Organisation.
  39. World Health Organisation (2002). *Promoting Effective Perinatal Care; Essential Antenatal, Perinatal and Postpartum Care Training modules*, WHO Regional Office for Europe.
  40. World Health Organisation (2003). Skilled attendants for pregnancy, childbirth and postnatal care. In: L de Bernis, D. R Sherratt, C. AbouZahr & W.V. Lerberghen (Eds), *British Medical Bulletin* 2003; 67: 39–57. World Health Organisation, Geneva, Switzerland.
  41. World Health Organisation Commission on Social Determinants of Health (2008). *Closing the gap in a generation: Health equity through action on the social determinants of health*. The Final Report of the WHO Commission on Social Determinants of Health, 28 August, 2008. WHO, Geneva.
  42. World Health Organisation Commission on Social Determinants of Health (2010). *A conceptual framework for action on the social determinants of health*. Social Determinants of Health Discussion Paper 2 (Policy & Practice). In Solar O, Irwin A. (Ed). WHO, Geneva.

43. World Health Organisation (2013). World health report 2013: Research for universal coverage. WHO Press, Geneva.
44. World Health Organisation (2014). *Gender, women and health*: WHO Press, Geneva. Accessed at [www.who.int/genderandhealth/en/](http://www.who.int/genderandhealth/en/) on 6<sup>th</sup> October, 2014.
45. Yendi municipal assembly (2015). Municipal profile. Yendi, Northern Region.



This page is intentionally left blank



GLOBAL JOURNAL OF MEDICAL RESEARCH: E  
GYNECOLOGY AND OBSTETRICS  
Volume 17 Issue 2 Version 1.0 Year 2017  
Type: Double Blind Peer Reviewed International Research Journal  
Publisher: Global Journals Inc. (USA)  
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

## Massive Ascites Complicating Severe Preeclampsia : A Case Report

By Bharathi K.R, Vijayalakshmi S & Mahendra G

*Introduction-* Pre-eclampsia earlier known as Toxemia in pregnancy/pregnancy induced hypertension is primarily a endothelial disorder which is thought to be triggered by the placenta.<sup>1</sup> It affects about 4-8% of all pregnancies. Hypertensive disorders in pregnancy remains one of major component of the lethal triad in causing maternal mortality even in the era of modern obstetrics with the advent of newer management options and intensive care facilities being available.<sup>2</sup> The complications associated with pre-eclampsia are estimated to be around 20%. Complications like intracranial hemorrhage, cerebral edema, pulmonary edema, renal failure, global haemolytic tendency, HELLP syndrome and maternal ascites are all found in pre-eclampsia. The finding of presence of fluid in various body cavities is due to increased endothelial permeability and microvascular damage which is not unusual.<sup>3,4</sup> Some amount of fluid in peritoneal cavity is seen frequently seen in pre-eclamptic women but presence of massive ascites is a rare complication and needs cautious management.

*GJMR-E Classification: NLMC Code: WJ 190*



*Strictly as per the compliance and regulations of:*



# Massive Ascites Complicating Severe Preeclampsia : A Case Report

Bharathi K.R <sup>α</sup>, Vijayalakshmi S <sup>σ</sup> & Mahendra G <sup>ρ</sup>

## I. INTRODUCTION

**P**re-eclampsia earlier known as Toxemia in pregnancy/pregnancy induced hypertension is primarily a endothelial disorder which is thought to be triggered by the placenta.<sup>1</sup> It affects about 4-8% of all pregnancies. Hypertensive disorders in pregnancy remains one of major component of the lethal triad in causing maternal mortality even in the era of modern obstetrics with the advent of newer management options and intensive care facilities being available.<sup>2</sup> The complications associated with pre-eclampsia are estimated to be around 20%. Complications like intracranial hemorrhage, cerebral edema, pulmonary edema, renal failure, global haemolytic tendency, HELLP syndrome and maternal ascites are all found in pre-eclampsia. The finding of presence of fluid in various body cavities is due to increased endothelial permeability and microvascular damage which is not unusual.<sup>3,4</sup> Some amount of fluid in peritoneal cavity is seen frequently seen in pre-eclamptic women but presence of massive ascites is a rare complication and needs cautious management.

## II. CASE REPORT

A 19year old primigravida with h/o 9 months of amenorrhea with good perception of fetal movements and regular antenatal checkups was referred from a private hospital to our antenatal OPD at AIMS, BG Nagara in view of high blood pressure readings. She gave h/o pedal edema and high blood pressure since 2 months for which she was on treatment with tab. Nicardipine 5mg thrice daily which has been increased to 10 mg thrice daily since 3 days. No h/o imminent symptoms. On examination, there was no pallor, icterus. Bilateral pitting type of pedal edema present. Pulse rate 94/min, BP was 160/120mmhg, respiratory rate 22/min, regular. Cardiovascular & respiratory system examination were within normal limits. Abdominal examination showed abdominal wall edema, gross maternal ascites with fluid thrill, 28weeks size uterus, FHS couldnot be localised clinically but fetal cardiac activity and rate noted with the help of bedside scan. Vulval edema present. Vaginal examination revealed

uneffaced and closed cervical os and diagnosis of 36weeks gestation with severe preeclampsia with severe IUGR with oligohydramnios with gross maternal ascites was made. Urine albumin was 4+. Serum uric acid 6mg/dl. Complete blood count, blood urea, serum creatinine, serum bilirubin, ALT, AST, LDH were within normal limits. Fundoscopy revealed grade I hypertensive retinopathy changes. She was given 10mg Nicardipine stat dose and 12mg betamethasone injection was given. Decision for caesarean section was taken in view of severe IUGR and oligohydramnios. Under spinal anesthesia, caesarean section was done and delivered a live preterm male baby weighing 1.5kg. 2litres of ascitic fluid drained intraoperatively, BP 140/100mmhg. Immediate postoperative period was uneventful. She was started on clear liquids orally after 12 hours, BP was 130/100mmhg, pulse rate 88/min.

On 2<sup>nd</sup> postoperative day patient developed abdominal distension with breathing difficulty. On examination pulse was 116/min, BP 130/100mmhg, with RR 24/min. Abdomen soft, nontender, ascites +. Ultrasonography revealed gross ascites and bilateral mild pleural effusion. Echocardiogram was normal. Serum albumin was 2.6g/dl, all other investigations were within normal limits. Patient was shifted to intensive care unit. Ascitic fluid tapping done. Strict input/output charting maintained. Tab Nicardipine 10mg thrice daily continued. On 3<sup>rd</sup> and 4<sup>th</sup> post op day she continued to have tachycardia and abdominal distension. High protein diet started. From 5<sup>th</sup> postop day the tachycardia started settling down and also the blood pressure with BP 140/90mmhg, abdominal distension also started reducing which was monitored by measuring abdominal girth. Complete hemogram, Blood urea, creatinine, AST, ALT, LDH, serum bilirubin were within normal limits. By 10<sup>th</sup> day distension reduced completely. Review ultrasound revealed no pleural effusion but mild ascetic fluid. Nicardipine was stopped on 15<sup>th</sup> day and BP monitoring was done which was within normal limits. She was discharged on 20<sup>th</sup> day in stable condition with instructions for weekly follow up. Review follow up was uneventful with normal BP readings.

## III. DISCUSSION

Pre-eclampsia is a multisystem disorder characterised by the involvement of global microvascular system associated with endothelial dysfunction. Out of all the complications known to occur

*Author α ρ: Asst professor, AIMS, BG Nagara.*

*e-mail: bharathikr84@gmail.com*

*Author σ: Prof and head, dept of OBG, AIMS, BG Nagara.*

*Author: Postgraduate AIMS, BG Nagar.*

in pre-eclampsia, ascites is a rare complication. Fluid in the peritoneal cavity is a frequent finding in pre-eclamptic women but massive ascites is very rare and only few cases have been reported in the literature. Incidence of ascites in pre-eclampsia is 1 in 1000 patients. Fluid in multiple body cavities in pre-eclampsia is thought to be due to increased capillary permeability (capillary leak syndrome), portal hypertension, hypoproteinemia due to proteinuria and altered albumin/globulin ratio.<sup>5</sup>

Cong and Wang conducted a study in 23 patients with pre-eclampsia and ascites where they found albumin/globulin ratio  $<1.5$  in all patients.<sup>6</sup> The finding of altered albumin/globulin ratio was thought to reduce the intravascular oncotic pressure leading to transudation of serum to various body cavities predominantly peritoneal cavity.<sup>7</sup>

A case of massive ascites in pre-eclampsia with bilateral hydrothorax reported by Ashmore et al reveals the use of intraabdominal drain to monitor the ascetic fluid production.<sup>8</sup>

A case series conducted by Woo et al in pregnancy with severe-eclampsia and ascites reported rapid deterioration in maternal condition with rise in the blood pressure associated with increasing proteinuria requiring early termination of pregnancy.<sup>9</sup>

#### IV. CONCLUSION

Ascites in preeclampsia is a rare complication which need cautious management to avoid complications of massive ascites and hydrothorax like respiratory distress and cardiac compromise. Patients need to be monitored well in intensive care set ups and decision regarding termination of pregnancy should be taken based on the clinical scenario rather than making a decision for early termination of pregnancy.

#### REFERENCES RÉFÉRENCES REFERENCIAS

1. Calvin S, Silva M, Weinstein L et al. Characterization of ascites present at Caesarean Section. *Amer J Perinatal* 1991; 8: 99-102.
2. Vaijyanath AM, Nayar B, Malhotra N, et al. Massive ascites in severe pre-eclampsia : a rare complication. *J Obstet Gynecol Res* 2002; 28: 199-202.
3. Woods JB, Blake PG, Perry KG et al. Ascites : a portent of cardiopulmonary complications in the preeclamptic patient with the syndrome of hemolysis, elevated liver enzymes and low platelets. *Obstet Gynecol* 1992; 80: 87-91.
4. Foreman CS. Massive ascites as a complication of severe pre-eclampsia – a case report. *J Reprod Med* 1989; 34: 307-10.
5. Jimmy PW, Damar P. The unique presentation of massive ascites complicating severe preeclampsia

and HELLP syndrome. *Indonesia J Obstet and Gynecol* 2011; 35(2): 67-9.

6. Cong KJ, Wang TT. Complications of ascites in pregnancy induced hypertension. *Zhonghua Fu Chan Ke Za Zhi* 1994; 29(1): 7-9.
7. Lee Ko M, Huang LW, Chang JZ. Massive ascites complicating pre-eclampsia. *Taiwanese J Obstet Gynecol* 2005; 44(3): 267-9.
8. Ashmore j, Attapattu JA. Massive ascites and bilateral hydrothorax complicating severe preeclampsia. *Ceylon Med J* 1998; 43(4): 235-7.
9. Woo JS, Liang ST, Wong VT. Ultrasonic detection of maternal ascites in patients with severe pre-eclampsia. *Aust NZ J Obstet Gynecol* 1982; 22: 237-9.





GLOBAL JOURNAL OF MEDICAL RESEARCH: E  
GYNECOLOGY AND OBSTETRICS  
Volume 17 Issue 2 Version 1.0 Year 2017  
Type: Double Blind Peer Reviewed International Research Journal  
Publisher: Global Journals Inc. (USA)  
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

## Incidence, Clinical Characteristics and Outcome of Unexpected Uterine Sarcoma after Hysterectomy for Uterine Mass: A Retrospective Study of 774 Cases

By Monika Jindal & Rekha Ratnani

*Chandulal Chandrakar Memorial Medical College*

**Abstract- Background:** Women often require a hysterectomy via laparotomy or laparoscopy. Morcellation is often necessary to perform a laparoscopic surgery. The objective of this study is to determine the incidence of unexpected uterine sarcomas (UUSs) after hysterectomy and to reduce the occurrence and avoid the morcellation of UUSs by analyzing their characteristics.

**Methods:** Women who had a hysterectomy for various reasons in Chandulal Chandrakar Memorial Medical College, Durg, India between January 2014 and September 2016 were selected for the study, and their clinical characteristics were analyzed.

**Results:** During the period, 5 UUSs were found in 774 cases, and the overall incidence was 0.65%. Most of the UUSs were stage I (80%), which occurred more commonly (80%) in women aged 40–49. Abnormal uterine bleeding (60%) was the main clinical manifestation. The margins of most UUSs (80%) were regular, which may cause UUSs to be misdiagnosed as uterine fibroids.

**Keywords:** *laparoscopy, morcellation, uterine sarcoma, uterine fibroid, hysterectomy.*

**GJMR-E Classification:** *NLMC Code: WQ 440*



*Strictly as per the compliance and regulations of:*



# Incidence, Clinical Characteristics and Outcome of Unexpected Uterine Sarcoma after Hysterectomy for Uterine Mass: A Retrospective Study of 774 Cases

Monika Jindal <sup>α</sup> & Rekha Ratnani <sup>ο</sup>

**Abstract- Background:** Women often require a hysterectomy via laparotomy or laparoscopy. Morcellation is often necessary to perform a laparoscopic surgery. The objective of this study is to determine the incidence of unexpected uterine sarcomas (UUSs) after hysterectomy and to reduce the occurrence and avoid the morcellation of UUSs by analyzing their characteristics.

**Methods:** Women who had a hysterectomy for various reasons in Chandulal Chandrakar Memorial Medical College, Durg, India between January 2014 and September 2016 were selected for the study, and their clinical characteristics were analyzed.

**Results:** During the period, 5 UUSs were found in 774 cases, and the overall incidence was 0.65%. Most of the UUSs were stage I (80%), which occurred more commonly (80%) in women aged 40–49. Abnormal uterine bleeding (60%) was the main clinical manifestation. The margins of most UUSs (80%) were regular, which may cause UUSs to be misdiagnosed as uterine fibroids.

**Conclusion:** The incidence of UUSs after hysterectomy for uterine fibroids was low, and their clinical characteristics are atypical. It is necessary and very critical to make a complete and cautious preoperative evaluation to reduce the occurrence and avoid the morcellation of UUSs.

**Keywords:** laparoscopy, morcellation, uterine sarcoma, uterine fibroid, hysterectomy.

## I. INTRODUCTION

Uterine fibroids are one of the most common types of pelvic tumors in women. Various treatments are available for the management of uterine fibroids, such as having a hysterectomy or a myomectomy. The surgical route is shifting from an abdominal to a laparoscopic approach, which confers a more rapid recovery and fewer perioperative complications.<sup>1</sup> Morcellation of the specimen is often necessary to perform a laparoscopy.

Uterine sarcoma is a rare, malignant tumor affecting the female genital system that accounts for only 1%–3% of uterine malignancies. The 5-year survival rate is approximately 30%.<sup>2,3</sup> The degree of malignancy

is high and the prognosis is poor. Inadvertent morcellation of unexpected uterine sarcomas (UUSs) is a surgical risk of laparoscopic hysterectomies and myomectomies. Evidence suggests that morcellation of UUSs potentially upstages the disease, which portends a poor prognosis.<sup>4,5</sup> The lack of specific symptoms, signs, or diagnostic techniques for preoperative differentiation from uterine fibroids results in most patients being diagnosed after surgery. If a uterine sarcoma is mistakenly diagnosed as a uterine fibroid and is morcellated via laparoscopy, serious consequences may arise.

The objective of this study is to determine the incidence of UUSs after hysterectomy for uterine fibroids. Until now, few studies have described the characteristics of UUSs in detail. This study is also designed to help reduce the occurrence and avoid the morcellation of UUSs in future by analyzing their characteristics.

## II. MATERIALS AND METHODS

### a) Case selection

This study was approved by the Institutional Review Board at CCM Medical College, Durg, India and all subjects provided informed consent. Women who had a hysterectomy in CCM Medical College, Durg, India between January 2014 and September 2016 were selected for the study. The average patient age was  $48.20 \pm 7.64$  years. Among them, patients who were diagnosed with uterine sarcomas based on postoperative pathology were also selected. Their median age was 41 years. Data analysis was performed to determine the incidence of UUSs. Finally, we reviewed the patient's clinical manifestations, laboratory tests, imaging studies, intraoperative findings, surgical pathologic stage, postoperative pathology, and prognosis.

### b) Statistical analysis

Statistical analysis was performed with SPSS 19.0 software (IBM Corporation, Armonk, NY, USA). All results are expressed as the mean  $\pm$  standard deviation. The Pearson's chi-square test was used for

Author <sup>α</sup>: Chandulal Chandrakar Memorial Medical College, India.  
e-mail: drmonickajindal@gmail.com

comparison of percentages between the groups. Statistical significance was assumed at a  $P \leq 0.05$ .

### III. RESULTS

#### a) Incidence of unexpected uterine sarcomas

During the period, 5 UUSs were found in 774 cases, and the overall incidence was 0.65%. Among the 5 patients, open surgery was performed on 4 patients

(80%), accounting for approximately 0.01% of 625 laparotomies. There was one laparoscopy (20%), accounting for approximately 0.01% of 149 laparoscopies. There was no statistical difference ( $P=1.05$ ) regarding the incidence of UUSs between laparotomies and laparoscopies.

The details of the UUSs are shown in Table 1.

| S.no | Age (years) | Clinical Presentation   | Associated Co-morbidities         | Type and Stage                | Treatment Received  | Outcome                                 |
|------|-------------|-------------------------|-----------------------------------|-------------------------------|---|---|
| 1    | 49          | Lump Abdomen (24 weeks) | Hypertension, Sickle Cell disease | Leiomyosarcoma 3              | Abdominal Hysterectomy with Bilateral Salpingo-oophorectomy With regional Lymphadenectomy | Lost to follow-up                       |
| 2    | 41          | AUB                     | Nil                               | Leiomyosarcoma 1              | Total Laparoscopic Hysterectomy   | Died                                    |
| 3    | 23          | AUB Dysmenorrhoea       | Anemia                            | Endometrial Stromal Sarcoma 1 | First Surgery – Myomectomy<br>Second Surgery – Wertheim's Hysterectomy                    | Chemotherapy                            |
| 4    | 40          | Acute Abdomen           | Nil                               | Carcinosarcoma 1              | Radical Hysterectomy with Bilateral retroperitoneal pelvic lymphadenectomy                | Chemotherapy and Radiotherapy completed |
| 5    | 45          | AUB                     | Nil                               | Leiomyosarcoma 1              | Total Abdominal Hysterectomy with Bilateral Salpingo-oophorectomy                         | Chemoradiotherapy                       |

The results showed that 4 patients were between 40 and 49 years, comprising the highest proportion (80%) of the 5 UUSs. There was only one patient aged 23 years, accounting for 20%. Thus, it can be seen that UUSs occurred more commonly in women aged 40–49, but patients aged 21–39 and aged 50–59 cannot be ignored.

#### b) Clinical manifestations

The clinical manifestations of uterine sarcomas are nonspecific. Classically, uterine sarcomas are always present as rapidly growing pelvic masses, which may be accompanied by abnormal uterine bleeding and abdominal or pelvic pain.<sup>6,7</sup> In our study, 3 UUSs (60%) presented with abnormal uterine bleeding (menorrhagia, menostaxis, and irregular uterine bleeding), 1 UUSs (20%) presented with acute abdomen, and one UUS (20%) presented with rapidly growing pelvic masses. Abnormal uterine bleeding was the main clinical manifestation. A rapidly growing pelvic mass may be an indication of uterine sarcoma.

#### c) Laboratory tests and imaging studies involving unexpected uterine sarcomas

Of the 5 UUSs, laboratory tests for tumor markers, especially CA-125, were performed on 2 (40%). Only one case had high CA-125 values i.e. 51.2 mIU/mL, and in other it was found to be normal i.e. 23.0 mIU/mL. The sensitivity of CA-125 for uterine sarcomas is only 50% and its specificity is poor. We concluded, therefore, that CA-125 contributes minimally to the early diagnosis of uterine sarcoma.

The 5 UUS cases all underwent ultrasonic examinations, and the preoperative ultrasound

diagnosis was a uterine fibroid in 4(80%) and ovarian mass in 1 (20%). The diameters of the pelvic mass ranged from 6 to 15 cm and the median value was 7 cm. The diameter of the mass in 3 cases (60%) was less than 8 cm, in one case (20%) from 8 to 10 cm, and only one case (20%) was above 10 cm. Among the 5 UUSs, only one case (20%) had irregular margins. The margins of typical uterine sarcomas are mostly nodular, irregular, or ill defined, and the uterus is significantly enlarged, but the size and margins of most UUSs prompted by ultrasonography in this study were atypical, which may cause UUSs to be misdiagnosed as uterine fibroids. Color Doppler flow imaging of one of the 5 (20%) UUSs demonstrated rich blood flow signals, and degeneration of uterine fibroids was found in one of 5 (20%) cases. A rich blood flow signal around a pelvic mass and the degeneration of a uterine fibroid may suggest that more methods are needed to rule out the possibility of a uterine sarcoma.

In our study, 2 out of 5 UUSs (40%) were examined by computed tomography (CT). The preoperative CT diagnoses was uterine fibroid in one and ovarian mass in one. CT cannot be used to help distinguish uterine sarcomas from uterine fibroids. This study showed that it is more difficult to distinguish between benign degenerating uterine fibroids and malignant uterine sarcomas. When the pathology reports were available, the CT images were reviewed again by the pathologist, but a preoperative CT diagnosis of a uterine sarcoma still could not be made so easily. In such patients, alternative treatment options should be carefully considered.

d) *Surgical stage, postoperative pathology, and the prognosis of UUSs*

According to the postoperative pathology report after the first operation, there were three leiomyosarcomas (60%), two low-grade endometrial stromal sarcomas (ESSs, 40%). According to the FIGO (International Federation of Gynecology and Obstetrics) 2009 staging for uterine sarcomas, the number of uterine sarcomas for stages I, II, and III after the first operation was four, zero, and one, respectively, thus accounting for approximately 80%, 0%, and 20% of the five UUSs. When postoperative pathology was confirmed, one case (20%) underwent a second operation. Stage I was the first operative stage. This case underwent laparoscopy that was converted to laparotomy during the first surgery. The time interval between the two operations was 15 days.

The follow-up percentage was 60%. The follow-up time was 4–11 months, and the median follow-up time was 7 months. All these patients are of Stage I sarcoma. One case died after 9 months of diagnosis and was found after a telephonic call as she was lost to follow-up. She took 2 cycles of chemotherapy. One more case lost to follow-up and is not traceable.

#### IV. DISCUSSION

A number of studies<sup>10,14,15</sup> have reported a poor prognosis for uterine sarcomas closely related to the clinical stage and the method used in the primary operation. In this study, the staging of most UUSs (80%) is stage I, which may indicate a good prognosis, but how to make an accurate diagnosis and choose an appropriate treatment is critical. To reduce the occurrence of UUSs and the risk of spreading UUSs by laparoscopic morcellation, the method used for the preoperative diagnosis of a uterine sarcoma is very important. Brohl et al<sup>16</sup> found the risk of UUSs varied significantly across age groups, and the risk of uterine sarcoma ranged from a peak of 10.1 cases per 1,000 for patients aged 75–79 years to, 1 case per 500 for patients aged, 30 years. However, our study found that UUSs occurred more commonly in women aged 40–49. Abnormal uterine bleeding was the main clinical manifestation, and cases with no symptoms cannot be ignored. A rapidly growing pelvic mass may be indicative of a uterine sarcoma. The size and margins of most UUSs as suggested by ultrasonography were atypical, and UUSs can be easily misdiagnosed as uterine fibroids. Attention should be paid to a rich blood flow signal around a pelvic mass and the degeneration of uterine fibroids, and more methods are needed to rule out the possibility of uterine sarcomas. CT may have a role in the evaluation of pelvic masses. In our study, CT was of no use as it was unable to differentiate between fibroids and sarcomas.

When the pathology reports were available, the pathologist reviewed the CT images again, but it was still difficult to make a preoperative CT diagnosis of a uterine sarcoma. This told us that it is more difficult to distinguish between benign degenerating uterine fibroids and malignant uterine sarcomas. When the malignant potential of a uterine mass is uncertain, alternative treatment options should be carefully considered.

A number of studies<sup>20–25</sup> have shown that the incidence of UUSs ranges from 0.09% to 0.49% among women undergoing benign hysterectomy or myomectomy. In this, the incidence of UUSs during hysterectomy performed for various reasons was 0.65%, and there was no statistical difference ( $P=1.04$ ) in the incidence between laparotomy and laparoscopy (0.01% vs 0.01%).

#### V. CONCLUSION

In conclusion, the incidence of UUSs after hysterectomy and myomectomy was low and their clinical characteristics are atypical. It is critical to make a complete and cautious preoperative evaluation to reduce the occurrence and avoid the morcellation of UUSs. Due to the limitations of sample size, further research should be carried out to assess the impact of morcellation on the prognosis of UUSs and summarize the clinical characteristics of UUSs to avoid the risk in future.

#### REFERENCES RÉFÉRENCES REFERENCIAS

1. Nieboer TE, Johnson N, Lethaby A, et al. Surgical approach to hysterectomy for benign gynaecological disease. *Cochrane Database Syst Rev*. 2009; (3): CD003677.
2. Sehnal B, Driak D, Kmonickova E, et al. Current classification of malignant tumours in gynecological oncology – part I. *Ceska Gynekol*. 2011; 76(4): 279–284. Czech.
3. Sehnal B, Driak D, Kmonickova E, et al. Current classification of malignant tumours in gynecological oncology – part II. *Ceska Gynekol*. 2011; 76(5): 360–366. Czech.
4. Einstein MH, Barakat RR, Chi DS, et al. Management of uterine malignancy found incidentally after supracervical hysterectomy or uterine morcellation for presumed benign disease. *Int J Gynecol Cancer*. 2008; 18(5):1065–1070.
5. Oduyebo T, Rauh-Hain AJ, Meserve EE, et al. The value of re-exploration in patients with inadvertently morcellated uterine sarcoma. *Gynecol Oncol*. 2014; 132(2): 360–365.
6. Seddon BM, Davda R. Uterine sarcomas – recent progress and future challenges. *Eur J Radiol*. 2011; 78(1): 30–40.



7. Wu TI, Yen TC, Lai CH. Clinical presentation and diagnosis of uterine sarcoma, including imaging. *Best Pract Res Clin Obstet Gynaecol.* 2011; 25(6): 681–689.
8. FDA. Laparoscopic Uterine Power Morcellation in Hysterectomy and Myomectomy: FDA Safety Communication. Available from: <http://www.fda.gov/MedicalDevices/Safety/AlertsandNotices/ucm393576.htm>. Accessed April 17, 2014.
9. George S, Barysaukas C, Serrano C, et al. Retrospective cohort study evaluating the impact of intraperitoneal morcellation on outcomes of localized uterine leiomyosarcoma. *Cancer.* 2014; 120(20): 3154–3158.
10. Morice P, Rodriguez A, Rey A, et al. Prognostic value of initial surgical procedure for patients with uterine sarcoma: analysis of 123 patients. *Eur J Gynaecol Oncol.* 2003; 24(3–4): 237–240.
11. Cusido M, Fargas F, Baulies S, et al. Impact of surgery on the evolution of uterine sarcomas. *J Minim Invasive Gynecol.* 2015; 22(6): 1068–1074.
12. Park JY, Kim DY, Kim JH, Kim YM, Kim YT, Nam JH. The impact of tumor morcellation during surgery on the outcomes of patients with apparently early low-grade endometrial stromal sarcoma of the uterus. *Ann Surg Oncol.* 2011; 18(12): 3453–3461.
13. Park JY, Park SK, Kim DY, et al. The impact of tumor morcellation during surgery on the prognosis of patients with apparently early uterine leiomyosarcoma. *Gynecol Oncol.* 2011; 122(2): 255–259.
14. Gadducci A, Cosio S, Romanini A, Genazzani AR. The management of patients with uterine sarcoma: a debated clinical challenge. *Crit Rev Oncol Hematol.* 2008; 65(2): 129–142.
15. Perri T, Korach J, Sadetzki S, Oberman B, Fridman E, Ben-Baruch G. Uterine leiomyosarcoma: does the primary surgical procedure matter? *Int J Gynecol Cancer.* 2009; 19(2): 257–260.
16. Brohl AS, Li L, Andikyan V, et al. Age-stratified risk of unexpected uterine sarcoma following surgery for presumed benign leiomyoma. *Oncologist.* 2015; 20(4): 433–439.
17. Santos P, Cunha TM. Uterine sarcomas: clinical presentation and MRI features. *Diagn Interv Radiol.* 2015; 21(1):4–9.
18. Takeuchi M, Matsuzaki K, Harada M. Preliminary observations and clinical value of lipid peak in high-grade uterine sarcomas using in vivo proton MR spectroscopy. *Eur Radiol.* 2013; 23(9): 2358–2363.
19. Thomassin-Naggara I, Dechoux S, Bonneau C, et al. How to differentiate benign from malignant myometrial tumours using MR imaging. *Eur Radiol.* 2013; 23(8): 2306–2314.
20. Wright JD, Tergas AI, Cui R, et al. Use of electric power morcellation and prevalence of underlying cancer in women who undergo myomectomy. *JAMA Oncol.* 2015; 1(1): 69–77.
21. Parker WH, Fu YS, Berek JS. Uterine sarcoma in patients operated on for presumed leiomyoma and rapidly growing leiomyoma. *Obstet Gynecol.* 1994; 83(3): 414–418.
22. Takamizawa S, Minakami H, Usui R, et al. Risk of complications and uterine malignancies in women undergoing hysterectomy for presumed benign leiomyomas. *Gynecol Obstet Invest.* 1999; 48(3): 193–196.
23. Theben JU, Schellong AR, Altgassen C, Kelling K, Schneider S, Grosse-Drieling D. Unexpected malignancies after laparoscopic-assisted supracervical hysterectomies (LASH): an analysis of 1,584 LASH cases. *Arch Gynecol Obstet.* 2013; 287(3): 455–462.
24. Zhang J, Zhang J, Dai Y, Zhu L, Lang J, Leng J. Clinical characteristics and management experience of unexpected uterine sarcoma after myomectomy. *Int J Gynaecol Obstet.* 2015; 130(2): 195–199.
25. Graebe K, Garcia-Soto A, Aziz M, et al. Incidental power morcellation of malignancy: a retrospective cohort study. *Gynecol Oncol.* 2015; 136(2):274–277.
26. McKenna JB, Kanade T, Choi S, et al. The Sydney contained in bag morcellation technique. *J Minim Invasive Gynecol.* 2014; 21(6): 984–985.
27. Cholkeri-Singh A, Miller CE. Power morcellation in a specimen bag. *J Minim Invasive Gynecol.* 2015; 22(2): 160.



GLOBAL JOURNAL OF MEDICAL RESEARCH: E  
GYNECOLOGY AND OBSTETRICS  
Volume 17 Issue 2 Version 1.0 Year 2017  
Type: Double Blind Peer Reviewed International Research Journal  
Publisher: Global Journals Inc. (USA)  
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

## Effect of Pregnancy and Mode of Delivery on HPV Infection and on Cervical Cytological Changes

By Lizhou Sun, Preetam Kona Herkanaidu & Purnima Mohur  
*Nanjing Medical University*

**Abstract- Background:** To know if the hormonal changes during pregnancy and the mode of delivery affect HPV (human papillomavirus) infection and on cervical cytological changes concerning the rates of persistence, progression and regression in the postpartum period following hormonal normalization.

**Methods:** A prospective study was carried out on 57 pregnant women who attended the Nanjing Maternity and Child Health Care Hospital between 2015 and 2017 who were positive for cervical cytological changes and HPV during their antenatal visits by TCT (PAP smear) and hybrid capture test2 respectively. Initial and postpartum results of these tests were compared where the rates of persistence, progression and regression of HPV and the cervical cytological changes were analyzed. For analysis of the results, a control group of 57 nonpregnant patients from the gynecological department was included and followed in this study.

**Results:** The postpartum evaluation of the pregnant cohort revealed a significantly higher tendency to spontaneous regression as compared to the non-pregnant control group (59.9% versus 31.6 %,  $p = 0.005$ ) and the mode of delivery did not affect cervical cytological changes and HPV infection ( $p=0.140$ ).

**Conclusion:** The high regression rate suggests that in the absence of invasive disease, definite management of abnormal cervical cytology can be carried out in the postpartum period.



*Strictly as per the compliance and regulations of:*



# Effect of Pregnancy and Mode of Delivery on HPV Infection and on Cervical Cytological Changes

Lizhou Sun <sup>α</sup>, Preetam Kona Herkanaidu <sup>σ</sup> & Purnima Mohur <sup>ρ</sup>

**Abstract- Background:** To know if the hormonal changes during pregnancy and the mode of delivery affect HPV (human papillomavirus) infection and on cervical cytological changes concerning the rates of persistence, progression and regression in the postpartum period following hormonal normalization.

**Methods:** A prospective study was carried out on 57 pregnant women who attended the Nanjing Maternity and Child Health Care Hospital between 2015 and 2017 who were positive for cervical cytological changes and HPV during their antenatal visits by TCT (PAP smear) and hybrid capture test2 respectively. Initial and postpartum results of these tests were compared where the rates of persistence, progression and regression of HPV and the cervical cytological changes were analyzed. For analysis of the results, a control group of 57 nonpregnant patients from the gynecological department was included and followed in this study.

**Results:** The postpartum evaluation of the pregnant cohort revealed a significantly higher tendency to spontaneous regression as compared to the non-pregnant control group (59.9% versus 31.6 %,  $p = 0.005$ ) and the mode of delivery did not affect cervical cytological changes and HPV infection ( $p=0.140$ )

**Conclusion:** The high regression rate suggests that in the absence of invasive disease, definite management of abnormal cervical cytology can be carried out in the postpartum period.

## I. BACKGROUND

HPV infection is highly prevalent in sexually active women worldwide, mostly in the non-vaccinated females, and even among those presenting normal cytology [1]. It is one of the most common sexually transmitted infections in the world and most sexually active women will be infected with HPV at least once during their lifetime, with the majority of infections cleared within two years [2,3]. Based on their association with cervical cancer (CC), HPV genotypes are classified as low-risk or high-risk [4]. High-risk (HPV) is a small DNA tumor virus that infects the mucosal squamous epithelium and causes various malignant

*Author α:* Department of Obstetrics, Nanjing Maternity and Child Health Care Hospital, Affiliated with Nanjing Medical University, Jiangsu Province, Nanjing, China. e-mails: lizhou\_sun121@sina.com, pkherkanaidu@yahoo.com

*Author σ:* Department of Obstetrics, Nanjing Maternity and Child Health Care Hospital, Affiliated with Nanjing Medical University, China.

*Author ρ:* Department of Obstetrics and Gynecology, JN Hospital, Mauritius.

diseases in humans, including cancers of the cervix [5]. It is well established that persistent infection with high-risk HPV genotypes is the necessary although not a sufficient cause of CC [6]. The involvement of other factors, in addition to HPV, is needed to induce cervical carcinogenesis and adequate immune response is crucial for HPV clearance while immune deficiency favors viral persistence and cervical cancer [7, 8]. Only 10-15% of women develop a persistent infection, which is an important risk factor for cervical carcinogenesis [9]. HPV persistence, even for a short time, has been associated with higher risk for cervical intra-epithelial neoplasia, compared to women without a history of HPV infection [10, 11]. After initial infection, HPV clearance is very frequent in the first six months, with rates of 50-70% per follow-up year [12, 13].

Physiological changes in immunity and other biological parameters (for example, changes in the levels of different hormones) during pregnancy and postpartum may change the natural history of HPV infection and most authors have found a reduction in HPV positivity during the postpartum period [14]. The aim of this study is to find out how pregnancy affects the cervical cytological changes and the natural history of HPV infection.

## II. MATERIALS AND METHODS

A prospective study was carried out in Nanjing Maternity and Child Health Care Hospital from January 2015 to December 2017.

The inclusion criteria were as follows;

- Adults aged 18 to 30 years with given consent
- Pregnancy in the first 24 weeks
- Females:
  - i. with positive TCT (Thin prep cytology test) and HPV positive
  - ii. having only one partner
  - iii. who do not smoke and do not indulge in alcohol or substance abuse.

The exclusion criteria were as follows;

- All immunosuppressed patients
- Patients who are or were known cases of cervical carcinoma
- Patients who had previous surgeries of the cervix

- Females who are known to be HPV positive and positive cervical cytology before pregnancy

All women up to 24 weeks of gestation attending the antenatal clinic for the first time in their pregnancy had TCT done. Here, after insertion of a vaginal speculum, we used plastic broom to collect material from the endocervix and this was then immediately distributed in a buffer solution by shaking the broom vigorously. The samples were then transported to the Molecular Biology. Those with positive cytological changes, that is positive PAPS smear, were asked in their next review to do the hybrid capture test 2 for HPV testing. These patients were seen as per the normal ANC review and no treatment was given for HPV positivity. However, after delivery, all those patients who were positive (ASCUS+HPV positive) were asked to repeat the TCT and the hybrid capture test 2 to know their status postpartum. The mode of delivery included both normal vaginal delivery and Caesarian section (Cs). We repeated the tests after a minimum of six weeks, that is the puerperium, when all the hormone levels and physiological changes during pregnancy return back to normal. During the same period of study, we followed a control group of 57 patients from the gynecological department with initial positive PAP smear (ASCUS-atypical squamous cell of undetermined significance) and positive hybrid capture test 2 (HPV positive). The mean period for the repeat tests for both pregnant and nonpregnant patients was 40 weeks. In this study, regression was defined as those cases where the first cervical cytology and HPV test was positive but became negative after the second test. Persistence were those cases in which both the first and the second tests were positive. Progression were those cases which worsened on the second test based on the cytological findings.

We used SPSS 19.0 statistical software for statistical analyses. p-values of  $<0.05$  were considered statistically significant. The impact of pregnancy and the mode of delivery on regression persistence and progression were assessed using the Chi-squared test. Binary logistic regression analysis was performed to calculate odds ratios and 95% confidence interval.

### III. RESULTS

Within this study period, a total of 114 patients with proven cytological changes and HPV infections were included in the analysis. Among them, 57 were pregnant and 57 who were nonpregnant served as the control group. As far as the effect of the mode of delivery is concerned, as shown in table 1, where the vaginal delivery rate was 75.4%, the regression rate was 65.1%. For the Cs group, the regression rate was 42.9%. Based on the p-value which was 0.140, that is  $>0.05$ , we did not find any significant difference between the effect of Cs and vaginal delivery on regression of HPV

and its cytological changes. On the other hand, as shown in table 2, in the group of pregnant women, the overall regression rate was 59.6%, which was higher than the regression rate in the control group that is 31.6%. Based on the chi-squared test, the p-value was 0.005 which is statistically significant and supports the fact that pregnancy affects the regression (that is a higher regression rate postpartum) of cervical HPV infection and its associated cytological changes.

*Table 1:* Comparison of effect of mode of delivery on HPV infection

| Mode of delivery | Persistence | Regression | X <sup>2</sup> | P     |
|------------------|-------------|------------|----------------|-------|
| Cs               | 8           | 6          | 2.174          | 0.140 |
| Vaginal delivery | 15          | 28         |                |       |

*Table 2:* Comparison of effect of pregnancy on HPV infection

|              | Persistence | Progression | Regression | X <sup>2</sup> | P     |
|--------------|-------------|-------------|------------|----------------|-------|
| Pregnant     | 23          | 0           | 34         | 10.78          | 0.005 |
| Non-pregnant | 36          | 3           | 18         |                |       |

*Table 3:* Characteristics associated with all the 114 pregnant and non-pregnant patients (multivariable analysis)

| Variables   | OR    | OR 95% CI |       |
|-------------|-------|-----------|-------|
|             |       | Low       | Up    |
| Regression  | 1     |           |       |
| Persistence | 0.395 | 0.186     | 0.839 |

### IV. DISCUSSION

The present study revealed high spontaneous regression in pregnant women. No cases of progression to higher grades cytological changes or invasive disease during the postpartum follow up period was found. These data support the opinion that once invasive disease can be excluded by colposcopy and CGB, definitive therapy in pregnant women can be deferred until after delivery. The initial obstetrical consultation provides an excellent opportunity to detect patients with abnormal PAP smears [15, 16, 17]. Historically, women with high-grade CIN were treated by cone biopsy during pregnancy [18, 19, 20]. Several studies reported that cone biopsy in pregnancy is associated with an impaired pregnancy outcome [21, 22]. Other reports showed that loop electrosurgical excision procedures are safe during pregnancy with a miscarriage rate  $<1\%$  [23]. Due to the low rates of progression during pregnancy, it is nowadays accepted that most patients may safely undergo expectant management if an invasive disease has been ruled out [24, 25]. Within the last decades, several authors studied the natural history of HPV diagnosed during



pregnancy with different outcomes, shown in table 4. The present study reports a significantly high postpartum regression rate of HPV in the group of pregnant patients that is 59.6%. The high regression rate is in accordance with recent previous studies, which report regression rates between 37 and 74 % for pregnant women at the time of postpartum follow up [26, 27, 28, 29, 30, 31, 32].

**Table 4:** Studies with reported outcomes

| Author             | Regression (%) |
|--------------------|----------------|
| Lurain [32]        | 77.4           |
| Yost [26]          | 69.3           |
| Vlahos [27]        | 61.6           |
| Paraskevaides [28] | 37.5           |
| Serati [31]        | 47.3           |

Many theories to explain these high regression rates can be found in the recent literature and among them most commonly the ones discussed below. Possible biological mechanisms for this could be that the raised levels of estrogen and progesterone during pregnancy which bring the following changes; the vaginal flora specifically presents an imbalance that, together with the dampness particular to that area, favors the development of infectious agents, including HPV. Also, during pregnancy, there occur anatomical modifications of the genital tract such as hypertrophy and congestion of the cervix, which increase, and is followed by metaplasia. The squamocolumnar junction undergoes alterations and maintains the transformation zone (TZ) on the exo-cervix (ectopy) for many years as a result of which this area of immature squamous metaplasia becomes more susceptible to the development of HPV infections and pre-neoplastic lesions [33]. These hormones also alter the local immune microenvironment of the cervix and sensitize the TZ to cervical cancer formation. The squamous epithelium of the cervix is composed of keratinocytes (primary target of HPV) and a type of immature dendritic cell (DC), the Langerhans cells (LC), which are important for the immuno-surveillance of the squamous epithelium [34].

Estradiol and progesterone influence the APC (antigen-presenting cell) functions of DC, with estradiol suppressing APC function, which may be due to decreased recruitment of DC or due to hormone-induced TGF- production that maintains DC in an immature state [35]. Moreover, in the transformation zone, estradiol has a high rate of conversion to 16 alpha hydroxyestrone [36, 37] which covalently bind and activate ER $\alpha$  (Estrogen receptor alpha). ER $\alpha$  is necessary for the genesis and continued growth of cancer and its expression in stromal cells is required for disease progression [38- 40]. The activated ER $\alpha$  is assumed to bind to responsive elements within the LCR (Long control region) and further induce E6 and E7

transcription to maintain HPV gene activity [37, 41]. Thus, it is hypothesized that both HPV and estradiol enhance the effects of each other, either directly through functional EREs (Estrogen responsive elements) in the viral genome or indirectly encourage uncontrolled cellular proliferation, thus enhancing malignant proliferation [42]. This synergistic combination of estrogen and HPV is the strongest factor in such carcinogenic transformations [38, 43-46]. Also, estrogen has a mitogenic activity which can be amplified by viral oncogenes [47]. It has been shown to stimulate the proliferation of human keratinocytes by promoting the expression of cyclin D2 and inducing G1 to S phase progression in the cell cycle [48]. Moreover, estrogen inhibits the oxidative stress-induced apoptosis in keratinocytes by promoting expression of the anti-apoptotic protein Bcl-2 [49]. It can also induce direct DNA damage via its catechol metabolites [50] and HPV infection has shown to considerably increase the formation of these potentially carcinogenic estrogen metabolites [36]. Also, there is increasing evidence that estrogen has the property to influence the immune system by acting on the cytokine production [51]. Estradiol has been shown to inhibit the expression of GM-CSF in the U2OS cell line through its interaction with ER $\alpha$  and to decrease this production via contact with ER $\beta$  [52].

Progesterone has also been shown to act on cytokine production to affect the immune system [51]. It increases the production of I $\kappa$ B, an inhibitor of NF- $\kappa$ B [53] and has an inhibitory effect on GMCSF secretion [54]. The immunosuppression caused mainly by the increased progesterone, which is necessary for maintaining a fetus during pregnancy, further predisposes a woman to the acquisition and development of lesions induced by HPV [33, 55, 56]. Researchers in molecular biology have also found an interaction between progesterone and HPV. It is known now that HPV 16 encodes a protein located in the region of E 6 and E 7 reading frames that cooperates with activated ras oncogene to transform primary cells [57]. The LCR of HPV 16 is reported to contain a deoxyribonucleic acid sequence that enhances response to both progesterone and glucocorticoids thereby increasing E 6 and E 7 transcription [58]. E6 and E7 oncoproteins bind to p53 and Rb, respectively, and surpass the host defense system [59-61]. In pregnancy, the elevated progesterone level increases HPV gene expression, giving rise to larger numbers of viral copy and multiplication of virus-transformed cells [62]. During the first trimester, there is a low immune response to HPV, thus accounting for the higher frequency of persistence of the virus. This deficient response, however, undergoes an intense recovery at the beginning of the third trimester, with reinforcement during the postpartum period to eventually lead to regression of the infection [63].

Furthermore, in cervical carcinogenesis, HPV DNA is often integrated into the host genome, leading to the loss of E2 gene. HPV E2 is a modulator of HPV gene expression and an inducer of apoptosis [64]. E2 and E7 proteins can induce apoptosis in transformed cells [65] and progesterone and estrogen increase the levels of E2 and E7-induced apoptosis [66]. However, with the loss of E2 in HPV infection, cell proliferation might increase [64]. Also, in the absence of E2, these hormones have been found to be a possible risk factor for cervical carcinogenesis by altering HPV gene expression [67]. Other authors speculated that the performance of multiple cervical biopsies in the antepartum evaluation could give the appearance of spontaneous regression [30]. Alternatively, other studies suggested a correlation between CIN course (cervical intraepithelial neoplasia) and mode of delivery and found a higher rate of regression of cervical dysplasia in association with vaginal delivery compared with cesarean section (67 versus 13 %) [30, 68]. A possible mechanism for this finding may be the loss of the dysplastic cervical epithelium during cervical ripening and the passage of the fetus through the birth canal [28, 69]. Other studies have not reported any differences in regression rates among patients who delivered vaginally or by cesarean section [26, 70], which is also the case in the present study.

Limitations of this study include

1. The limited number of cases.
2. The limited data on specific HPV subtypes have to be recognized as a limitation of the study.
3. Also, in the group of pregnant women we did not categorize the patients according to age. This can be a limitation as the females aged less than 24 years are known to be more often infected with HPV while at the same time they are known to clear the infection more spontaneously.
4. Inter-observer variability of TCT and HPV test results.

## V. CONCLUSION

This study supports the fact that a conservative management of HPV infection in pregnancy is safe since it reports high regression rates and no progression after delivery. However, this mode of management of HPV infection during pregnancy is valid as long as an invasive disease is ruled out.

## ACKNOWLEDGEMENT

Thank you to all those who supported me in this endeavor.

## REFERENCES RÉFÉRENCES REFERENCIAS

1. Bruni L, Diaz M, Castellsagué M, Ferrer E, Bosch FX, de Sanjosé S. Cervical Human Papillomavirus Prevalence in 5 continents: meta-analysis of 1 million women with normal cytological findings. *J Infect Dis.* 2010; 202: 1789–1799.
2. Baseman JG and Koutsky LA. The epidemiology of human papillomavirus infections. *J Clin Virol.* 2005; 32 Suppl 1: S16–S24.
3. Sawaya GF, Kulasingam S, Denberg TD, Qaseem A, Clinical Guidelines Committee of American College of Physicians. Cervical Cancer Screening in Average-Risk Women: Best Practice Advice From the Clinical Guidelines Committee of the American College of Physicians. *Ann Intern Med.* 2015; 162 (12): 851-9.
4. Cogliano V, Baan R, Straif K, Grosse Y, Secretan B, El Ghissassi F, WHO International Agency for Research on Cancer. Carcinogenicity of human papillomaviruses. *Lancet Oncol.* 2005; 6: 2042.
5. Leemans C.R., Braakhuis B.J., Brakenhoff R.H. The molecular biology of head and neck cancer. *Nat Rev Cancer.* 2011; 11: 9–22.
6. Munoz N, Castellsague X, de Gonzalez AB, Gissmann L. Chapter 1: HPV in the etiology of human cancer. *Vaccine.* 2006; 24 Suppl 3: S3-1-S310.
7. Song SH, Lee JK, Lee NW, Saw HS, Kang JS, Lee KW. Interferon- $\gamma$  (IFN- $\gamma$ ): a possible prognostic marker for clearance of high-risk human papillomavirus (HPV). *Gynecol Oncol.* 2008; 108: 543–548.
8. Einstein KH, Schiller JT, Viscidi RP, Strickler HD, Coursaget P, Tan T et al. Clinician's guide to human papillomavirus immunology: knowns and unknowns. *Lancet Infect Dis.* 2009; 9: 347–356.
9. Koshiol J, Lindsay L, Pimenta JM, Poole C, Jenkins D, Smith JS. Persistent human papillomavirus infection and cervical neoplasia: a systematic review and meta-analysis. *Am J Epidemiol.* 2008; 168: 123–137.
10. Naucler P, Ryd W, Tomberg S, Strand A, Wadell G, Elfgrén K et al. Human papillomavirus and Papanicolaou tests to screen for cervical cancer. *N Engl J Med.* 2007; 357: 1589–1597.
11. Castle PE, Rodriguez AC, Burk RD, Herrero R, Wacholder S, Alfaro M et al. Proyecto Epidemiológico Guanacaste (PEG) Group. Short term persistence of human papillomavirus and risk of cervical precancer and cancer: population based cohort study. *BMJ.* 2009; 339: b2569.
12. Winer RL, Hughes JP, Feng Q, Xi LF, Chernesky S, O'Reilly S et al. Early natural history of incident type-specific human papillomavirus in newly sexually active young women. *Cancer Epidemiol Biomarkers Prev.* 2011; 20: 699–707.
13. Oh JK, Ju YH, Franceschi S, Quint W, Shin HR. Acquisition of new infection and clearance of type-specific human papillomavirus infections in female

- students in Busan, South Korea: a follow-up study. *BMC Infect Dis.* 2008; 8: 13–18.
14. Ziegler A, Kastner C, Chang-Claude J. Analysis of pregnancy and other factors on detection of human papillomavirus (HPV) infection using eight estimating equations for follow-up data. *Statist Med.* 2003; 22: 2217–2233.
  15. Frega A, Scirpa P, Corosu R, Verrico M, Scarciglia ML, Primieri MR, et al. Clinical management and follow up of squamous intraepithelial cervical lesions during pregnancy and postpartum. *Anticancer Res.* 2007; 27: 2743–6.
  16. Xavier Júnior JC, Duflloth RM, do Vale DB, Tavares TA, Zeferino LC. Highgrade squamous intraepithelial lesions in pregnant and nonpregnant women. *Eur J Obstet Gynecol Reprod Biol.* 2014; 175: 103–6.
  17. Stonehocker J. Cervical cancer screening in pregnancy. *Obstet Gynecol Clin North Am.* 2013; 40: 269–82.
  18. Morice P, Uzan C, Gouy S, Verschraegen C, HaieMeder C. Gynaecological cancers in pregnancy. *Lancet.* 2012; 379: 558–69.
  19. Economos K, Perez Veridiano N, Delke I, Collado MI, Tancer MI. Abnormal cervical cytology in pregnancy: A 17year experience. *Obstet Gynecol.* 1993; 81: 915–8.
  20. Benedet JL, Selke RA, Nickerson KG. Colposcopic evaluation of abnormal Papanicolaou smears in pregnancy. *Am J Obstet Gynecol.* 1987; 157: 932–7.
  21. Averette HE, Nasser N, Yankow SL, Little WA. Cervical conization in pregnancy. Analysis of 180 operations. *Am J Obstet Gynecol.* 1970; 106: 543–9.
  22. Hannigan EV, Whitehouse 3rd HH, Atkinson WD, Becker SN. Cone biopsy during pregnancy. *Obstet Gynecol.* 1982; 60: 450–5.
  23. Kärrberg C, Brännström M, Strander B, Ladfors L, Rådberg T. Colposcopically directed cervical biopsy during pregnancy; minor surgical and obstetrical complications and high rates of persistence and regression. *Acta Obstet Gynecol Scand.* 2013; 92: 692–9.
  24. Jain AG, Higgins RV, Boyle MJ. Management of lowgrade squamous intraepithelial lesions during pregnancy. *Am J Obstet Gynecol.* 1997; 177: 298–302.
  25. Holowaty P, Miller AB, Rohan T, To T. Natural history of dysplasia of the uterine cervix. *J Natl Cancer Inst.* 1999; 91: 252–8.
  26. Yost NP, Santoso JT, McIntire DD, Iliya FA. Postpartum regression rates of antepartum cervical intraepithelial neoplasia II and III lesions. *Obstet Gynecol.* 1999; 93: 359–62.
  27. Vlahos G, Rodolakis A, Diakomanolis E, Stefanidis K, Haidopoulos D, Abela K, et al. Conservative Management of cervical intraepithelial neoplasia (CIN 2–3) in pregnant women. *Gynecol Obstet Invest.* 2002; 54: 78–81.
  28. Paraskevaidis E, Koliopoulos G, Kalantaridou S, Pappa L, Navrozoglou I, Zikopoulos K, et al. Management and evolution of cervical intraepithelial neoplasia during pregnancy and postpartum. *Eur J Obstet Gynecol and Reprod Biol.* 2002; 104: 67–9.
  29. Fader AN, Alward EK, Niederhauser A, Chirico C, Lesnock JL, Zwiesler DJ, et al. Cervical 7dysplasia in pregnancy: a multiinstitutional evaluation. *Am J Obstet Gynecol.* 2010 ; 203(2): 113. e16.
  30. Ahdoot D, Van Nostrand KM, Nguyen NJ, Tewari DS, Kurasaki T, DiSaia PJ, et al. The effect of route of delivery on regression of abnormal cervical cytologic findings in the postpartum period. *Am J Obstet Gynecol.* 1998; 178: 1116–20.
  31. Serati M, Uccella S, Laterza RM, Salvatore S, Beretta P, Riva C, et al. Natural history of cervical intraepithelial neoplasia during pregnancy. *Acta Obstet Gynecol.* 2008; 87: 1296–300.
  32. Lurain JR, Gallup DG. Management of abnormal PAP smears in pregnancy. *Obstet Gynecol.* 1979; 53: 484–8.
  33. Correia HS, Cornetta MCM, Gonçalves AKS. Infecção genital pelo papilomavírus humano (HPV) em mulheres grávidas. *Rev Brasil de Genit.* 2000; 1: 14-19.
  34. Romani N, Holzmann S, Tripp CH, Koch, F, Stoitzner P. Langerhans cells—dendritic cells of the epidermis. *APMIS.* 2003; 111: 725–740.
  35. Kenneth W. Beagley, Christine M. Gockel. Regulation of innate and adaptive immunity by the female sex hormones oestradiol and progesterone. *FEMS Immunology and Medical Microbiology.* 2003; 38: 13-22.
  36. Auburn KJ, Woodworth C, DiPaolo JA, Bradlow HL. The interaction between HPV infection and estrogen metabolism in cervical carcinogenesis. *Int J Cancer.* 1991; 49: 867-869.
  37. Matos A, Castelão C, Pereira da Silva A, Alho I, Bicho M, Medeiros R et al. Epistatic interaction of CYP1A1 and COMT polymorphisms in cervical cancer. *Oxidative Medicine and Cellular Longevity.* 2016; article 2769804.
  38. Chung SH, Wiedmeyer K, Shai A, Korach KS, Lambert PF. Requirement for estrogen receptor alpha in a mouse model for human papillomavirus-associated cervical cancer. *Cancer Research.* 2008; 68: 9928–9934.
  39. Chung SH, Lambert PF. Prevention and treatment of cervical cancer in mice using estrogen receptor antagonists. *Proc Natl Acad Sci USA.* 2009; 106: 19467–19472.
  40. Chung SH, Shin MK, Korach KS, Lambert PF. Requirement for stromal estrogen receptor alpha in cervical neoplasia. *Horm Cancer.* 2013; 4: 50–59.

41. De Villiers EM. Relationship between steroid hormone contraceptives and HPV cervical intraepithelial neoplasia and cervical carcinoma. *International Journal of Cancer*. 2003; 103: 705–708.
42. Ramachandran B. Functional association of oestrogen receptors with HPV infection in cervical carcinogenesis. *Endocrine-Related Cancer*. 2017; 24: R99–R108.
43. Brake T, Lambert PF. Estrogen contributes to the onset persistence and malignant progression of cervical cancer in a human papillomavirus-transgenic mouse model. *PNAS*. 2005; 102: 2490–2495.
44. Marks M, Gravitt PE, Gupta SB, Liaw KL, Kim E, Tadesse A et al. The association of hormonal contraceptive use and HPV prevalence. *International Journal of Cancer*. 2011; 128: 2962–2970.
45. Marks MA, Gupta S, Liaw KL, Tadesse A, Kim E, Phongnarisorn C et al. Prevalence and correlates of HPV among women attending family-planning clinics in Thailand. *BMC Infectious Diseases*. 2015; 15: 159.
46. Liao SF, Lee WC, Chen HC, Chuang LC, Pan MH, Chen CJ. Baseline human papillomavirus infection high vaginal parity and their interaction on cervical cancer risks after a follow-up of more than 10 years. *Cancer Causes and Control*. 2012; 23: 703–708.
47. Liehr JG. Is estradiol a genotoxic mutagenic carcinogen? *Endocr Rev*. 2000; 21: 40–54.
48. Kanda N, Watanabe S. 17beta-estradiol stimulates the growth of human keratinocytes by inducing cyclin D2 expression. *J Invest Dermatol*. 2004; 123: 319–328.
49. Kanda N, Watanabe S. 17beta-estradiol inhibits oxidative stress induced apoptosis in keratinocytes by promoting Bcl-2 expression. *J Invest Dermatol*. 2003; 121: 1500–1509.
50. Newfield L, Bradlow HL, Sepkovic DW, Auburn K. Estrogen metabolism and the malignant potential of human papillomavirus immortalized keratinocytes. *Proc Soc Expt Biol Med*. 1998; 217: 322–326.
51. McMurray RW, Ndebele K, Hardy KJ, Jenkins JK. 17-betaestradiol suppresses IL-2 and IL-2 receptor. *Cytokine*. 2001; 14: 324–333.
52. Brady H, Doubleday M, Gayo-Fung LM, Hickman M, Khammungkhune S, Kois A et al. Differential response of estrogen receptors alpha and beta to SP500263, a novel potent selective estrogen receptor modulator. *Mol Pharmacol*. 2002; 61: 562–568.
53. Miller L, Hunt JS. Regulation of TNF-alpha production in activated mouse macrophages by progesterone. *J Immunol*. 1998; 160: 5098–5104.
54. Robertson SA, Mayrhofer G, Seamark RF. Ovarian steroid hormones regulate granulocytemacrophage colony-stimulating factor synthesis by uterine epithelial cells in the mouse. *Biol Reprod*. 1996; 54: 183–196.
55. Rando RF, Lindheim S, Hasty L, Sedlacek TV, Woodland M, Eder C. Increased frequency of detection of human papillomavirus deoxyribonucleic acid in exfoliated cervical cells during pregnancy. *Am J Obstet Gynecol*. 1989; 161: 50–55.
56. Armbruster-Moraes E, Toshimoto LM, Leão E, Zugaib M. Prevalence of "high risk" human papillomavirus in the lower genital tract of Brazilian gravidas. *Int J Gynaecol Obstet*. 2000; 69: 223–227.
57. Matlaschewski G, Schneider J, Banks L, Jones N, Murray A, Crawford L. Human papillomavirus type 16 DNA cooperates with activated ras oncogene in transforming primary cells. *EMBO J*. 1987; 6: 1741.
58. Crook T, Storey A, Almond N, Osborn K, Crawford L. Human papillomavirus type 16 cooperates with activated ras and fos oncogenes in hormone dependent transformation of primary mouse cells. *Proc Natl Acad Sci USA*. 1988; 85: 8820.
59. Moodley M, Moodley J, Chetty R, Herrington CS. The role of steroid contraceptive hormones in the pathogenesis of invasive cervical cancer: a review. *International Journal of Gynecological Cancer*. 2003; 13: 103–110.
60. Sima N, Wang W, Kong D, Deng D, Xu Q, Zhou J, Xu G, Meng L, Lu Y, Wang S, et al. RNA interference against HPV16 E7 oncogene leads to viral E6 and E7 suppression in cervical cancer cells and apoptosis via upregulation of Rb and p53. *Apoptosis*. 2008; 2: 273–281.
61. Chung SH, Franceschi S, Lambert PF. Estrogen and ERalpha: culprits in cervical cancer? *Trends in Endocrinology and Metabolism*. 2010; 21: 504–511.
62. Chan WK, Klock G, Bernard HU. Progesterone and glucocorticoid response elements occur in the long control regions of several Human papillomaviruses involved in oncogenital neoplasia. *J Virol*. 1989; 63:4417.
63. Puig-Tintoré LM, Menendez AA, Bordoy XC, Bosch FX, Bladé AT, Castellsague X. La infección por papilomavirus, Documentos de consenso, SEGO, Barcelona 2002, p 56.
64. Webster K, Parish J, Pandya M, Stern PL, Clarke AR, Gaston K. The human papillomavirus (HPV) 16 E2 protein induces apoptosis in the absence of other HPV proteins and via a p53- dependent pathway. *J Biol Chem*. 2000; 275: 87–94.
65. Demeret C, Garcia-Carranca A, Thierry F. Transcription-independent triggering of the extrinsic pathway of apoptosis by human papillomavirus 18 E2 protein. *Oncogene*. 2003; 22: 168–175.
66. Webster K, Taylor A, Gaston K. Oestrogen and progesterone increase the levels of apoptosis induced by the human papillomavirus type 16 E2 and E7 proteins. *J Gen Virol*. 2001; 82: 201–213.

67. Delvenne P , Herman L, Kholod N, Caberg JH, Herfs M, Boniver J et al. Role of hormone cofactors in the human papillomavirus-induced carcinogenesis of the uterine cervix. *Molecular and Cellular Endocrinology*. 2007; 264: 1-5.
68. Siristatidis C, Vitoratos N, Michailidis E, Syciotis C, Panagiotopoulos N, Kassanos D, et al. The role of the mode of delivery in the alteration of intrapartum pathological cervical cytologic findings during the postpartum period. *Eur J Gynecol Oncol*. 2002; 23: 358–60.
69. Strinic T, Bukovic D, Karelavic D, Bojic L, Stipic I. The effect of delivery on regression of abnormal cervical cytologic findings. *Coll Antropol*. 2002; 26: 577–82.
70. Coppola A, Sorosky J, Casper R, Anderson B, Buller RE. The clinical course of cervical carcinoma in situ diagnosed during pregnancy. *Gynecol Oncol*. 1997; 67: 162–5.



GLOBAL JOURNALS INC. (US) GUIDELINES HANDBOOK 2017

---

[WWW.GLOBALJOURNALS.ORG](http://WWW.GLOBALJOURNALS.ORG)

# FELLOWS

## FELLOW OF ASSOCIATION OF RESEARCH SOCIETY IN MEDICAL (FARSM)

Global Journals Incorporate (USA) is accredited by Open Association of Research Society (OARS), U.S.A and in turn, awards “FARSM” title to individuals. The 'FARSM' title is accorded to a selected professional after the approval of the Editor-in-Chief/Editorial Board Members/Dean.



- The “FARSM” is a dignified title which is accorded to a person’s name viz. Dr. John E. Hall, Ph.D., FARSS or William Walldroff, M.S., FARSM.

FARSM accrediting is an honor. It authenticates your research activities. After recognition as FARSM, you can add 'FARSM' title with your name as you use this recognition as additional suffix to your status. This will definitely enhance and add more value and repute to your name. You may use it on your professional Counseling Materials such as CV, Resume, and Visiting Card etc.

*The following benefits can be availed by you only for next three years from the date of certification:*



FARSM designated members are entitled to avail a 40% discount while publishing their research papers (of a single author) with Global Journals Incorporation (USA), if the same is accepted by Editorial Board/Peer Reviewers. If you are a main author or co-author in case of multiple authors, you will be entitled to avail discount of 10%.

Once FARSM title is accorded, the Fellow is authorized to organize a symposium/seminar/conference on behalf of Global Journal Incorporation (USA). The Fellow can also participate in conference/seminar/symposium organized by another institution as representative of Global Journal. In both the cases, it is mandatory for him to discuss with us and obtain our consent.



You may join as member of the Editorial Board of Global Journals Incorporation (USA) after successful completion of three years as Fellow and as Peer Reviewer. In addition, it is also desirable that you should organize seminar/symposium/conference at least once.


We shall provide you intimation regarding launching of e-version of journal of your stream time to time. This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.





The FARSM can go through standards of OARS. You can also play vital role if you have any suggestions so that proper amendment can take place to improve the same for the benefit of entire research community.

As FARSM, you will be given a renowned, secure and free professional email address with 100 GB of space e.g. [johnhall@globaljournals.org](mailto:johnhall@globaljournals.org). This will include Webmail, Spam Assassin, Email Forwarders, Auto-Responders, Email Delivery Route tracing, etc.



The FARSM will be eligible for a free application of standardization of their researches. Standardization of research will be subject to acceptability within stipulated norms as the next step after publishing in a journal. We shall depute a team of specialized research professionals who will render their services for elevating your researches to next higher level, which is worldwide open standardization.

The FARSM member can apply for grading and certification of standards of their educational and Institutional Degrees to Open Association of Research, Society U.S.A. Once you are designated as FARSM, you may send us a scanned copy of all of your credentials. OARS will verify, grade and certify them. This will be based on your academic records, quality of research papers published by you, and some more criteria. After certification of all your credentials by OARS, they will be published on your Fellow Profile link on website <https://associationofresearch.org> which will be helpful to upgrade the dignity.



The FARSM members can avail the benefits of free research podcasting in Global Research Radio with their research documents. After publishing the work, (including published elsewhere worldwide with proper authorization) you can upload your research paper with your recorded voice or you can utilize chargeable services of our professional RJs to record your paper in their voice on request.



The FARSM member also entitled to get the benefits of free research podcasting of their research documents through video clips. We can also streamline your conference videos and display your slides/ online slides and online research video clips at reasonable charges, on request.







The FARSM is eligible to earn from sales proceeds of his/her researches/reference/review Books or literature, while publishing with Global Journals. The FARSS can decide whether he/she would like to publish his/her research in a closed manner. In this case, whenever readers purchase that individual research paper for reading, maximum 60% of its profit earned as royalty by Global Journals, will be credited to his/her bank account. The entire entitled amount will be credited to his/her bank account exceeding limit of minimum fixed balance. There is no minimum time limit for collection. The FARSM member can decide its price and we can help in making the right decision.

The FARSM member is eligible to join as a paid peer reviewer at Global Journals Incorporation (USA) and can get remuneration of 15% of author fees, taken from the author of a respective paper. After reviewing 5 or more papers you can request to transfer the amount to your bank account.



## MEMBER OF ASSOCIATION OF RESEARCH SOCIETY IN MEDICAL (MARSM)

The ' MARSM ' title is accorded to a selected professional after the approval of the Editor-in-Chief / Editorial Board Members/Dean.

The “MARSM” is a dignified ornament which is accorded to a person’s name viz. Dr. John E. Hall, Ph.D., MARSM or William Walldroff, M.S., MARSM.



MARSM accrediting is an honor. It authenticates your research activities. After becoming MARSM, you can add 'MARSM' title with your name as you use this recognition as additional suffix to your status. This will definitely enhance and add more value and repute to your name. You may use it on your professional Counseling Materials such as CV, Resume, Visiting Card and Name Plate etc.

*The following benefits can be availed by you only for next three years from the date of certification.*



MARSM designated members are entitled to avail a 25% discount while publishing their research papers (of a single author) in Global Journals Inc., if the same is accepted by our Editorial Board and Peer Reviewers. If you are a main author or co-author of a group of authors, you will get discount of 10%.

As MARSM, you will be given a renowned, secure and free professional email address with 30 GB of space e.g. [johnhall@globaljournals.org](mailto:johnhall@globaljournals.org). This will include Webmail, Spam Assassin, Email Forwarders, Auto-Responders, Email Delivery Route tracing, etc.





We shall provide you intimation regarding launching of e-version of journal of your stream time to time. This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.

The MARSM member can apply for approval, grading and certification of standards of their educational and Institutional Degrees to Open Association of Research, Society U.S.A.



Once you are designated as MARSM, you may send us a scanned copy of all of your credentials. OARS will verify, grade and certify them. This will be based on your academic records, quality of research papers published by you, and some more criteria.

It is mandatory to read all terms and conditions carefully.



## AUXILIARY MEMBERSHIPS

### Institutional Fellow of Open Association of Research Society (USA) - OARS (USA)

Global Journals Incorporation (USA) is accredited by Open Association of Research Society, U.S.A (OARS) and in turn, affiliates research institutions as “Institutional Fellow of Open Association of Research Society” (IFOARS).

The “FARSC” is a dignified title which is accorded to a person’s name viz. Dr. John E. Hall, Ph.D., FARSC or William Walldroff, M.S., FARSC.



The IFOARS institution is entitled to form a Board comprised of one Chairperson and three to five board members preferably from different streams. The Board will be recognized as “Institutional Board of Open Association of Research Society”-(IBOARS).

*The Institute will be entitled to following benefits:*



The IBOARS can initially review research papers of their institute and recommend them to publish with respective journal of Global Journals. It can also review the papers of other institutions after obtaining our consent. The second review will be done by peer reviewer of Global Journals Incorporation (USA) The Board is at liberty to appoint a peer reviewer with the approval of chairperson after consulting us.

The author fees of such paper may be waived off up to 40%.

The Global Journals Incorporation (USA) at its discretion can also refer double blind peer reviewed paper at their end to the board for the verification and to get recommendation for final stage of acceptance of publication.



The IBOARS can organize symposium/seminar/conference in their country on behalf of Global Journals Incorporation (USA)-OARS (USA). The terms and conditions can be discussed separately.

The Board can also play vital role by exploring and giving valuable suggestions regarding the Standards of “Open Association of Research Society, U.S.A (OARS)” so that proper amendment can take place for the benefit of entire research community. We shall provide details of particular standard only on receipt of request from the Board.



Journals Research  
inducing researches

The board members can also join us as Individual Fellow with 40% discount on total fees applicable to Individual Fellow. They will be entitled to avail all the benefits as declared. Please visit Individual Fellow-sub menu of GlobalJournals.org to have more relevant details.



We shall provide you intimation regarding launching of e-version of journal of your stream time to time. This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.



After nomination of your institution as “Institutional Fellow” and constantly functioning successfully for one year, we can consider giving recognition to your institute to function as Regional/Zonal office on our behalf. The board can also take up the additional allied activities for betterment after our consultation.

**The following entitlements are applicable to individual Fellows:**

Open Association of Research Society, U.S.A (OARS) By-laws states that an individual Fellow may use the designations as applicable, or the corresponding initials. The Credentials of individual Fellow and Associate designations signify that the individual has gained knowledge of the fundamental concepts. One is magnanimous and proficient in an expertise course covering the professional code of conduct, and follows recognized standards of practice.



Open Association of Research Society (US)/ Global Journals Incorporation (USA), as described in Corporate Statements, are educational, research publishing and professional membership organizations. Achieving our individual Fellow or Associate status is based mainly on meeting stated educational research requirements.

Disbursement of 40% Royalty earned through Global Journals : Researcher = 50%, Peer Reviewer = 37.50%, Institution = 12.50% E.g. Out of 40%, the 20% benefit should be passed on to researcher, 15 % benefit towards remuneration should be given to a reviewer and remaining 5% is to be retained by the institution.



We shall provide print version of 12 issues of any three journals [as per your requirement] out of our 38 journals worth \$ 2376 USD.

**Other:**

**The individual Fellow and Associate designations accredited by Open Association of Research Society (US) credentials signify guarantees following achievements:**

- The professional accredited with Fellow honor, is entitled to various benefits viz. name, fame, honor, regular flow of income, secured bright future, social status etc.



- In addition to above, if one is single author, then entitled to 40% discount on publishing research paper and can get 10% discount if one is co-author or main author among group of authors.
- The Fellow can organize symposium/seminar/conference on behalf of Global Journals Incorporation (USA) and he/she can also attend the same organized by other institutes on behalf of Global Journals.
- The Fellow can become member of Editorial Board Member after completing 3yrs.
- The Fellow can earn 60% of sales proceeds from the sale of reference/review books/literature/publishing of research paper.
- Fellow can also join as paid peer reviewer and earn 15% remuneration of author charges and can also get an opportunity to join as member of the Editorial Board of Global Journals Incorporation (USA)
- • This individual has learned the basic methods of applying those concepts and techniques to common challenging situations. This individual has further demonstrated an in-depth understanding of the application of suitable techniques to a particular area of research practice.

### Note :

//

- In future, if the board feels the necessity to change any board member, the same can be done with the consent of the chairperson along with anyone board member without our approval.
- In case, the chairperson needs to be replaced then consent of 2/3rd board members are required and they are also required to jointly pass the resolution copy of which should be sent to us. In such case, it will be compulsory to obtain our approval before replacement.
- In case of “Difference of Opinion [if any]” among the Board members, our decision will be final and binding to everyone.

//



## PROCESS OF SUBMISSION OF RESEARCH PAPER

---

The Area or field of specialization may or may not be of any category as mentioned in 'Scope of Journal' menu of the GlobalJournals.org website. There are 37 Research Journal categorized with Six parental Journals GJCST, GJMR, GJRE, GJMBR, GJSFR, GJHSS. For Authors should prefer the mentioned categories. There are three widely used systems UDC, DDC and LCC. The details are available as 'Knowledge Abstract' at Home page. The major advantage of this coding is that, the research work will be exposed to and shared with all over the world as we are being abstracted and indexed worldwide.

The paper should be in proper format. The format can be downloaded from first page of 'Author Guideline' Menu. The Author is expected to follow the general rules as mentioned in this menu. The paper should be written in MS-Word Format (\*.DOC, \*.DOCX).

The Author can submit the paper either online or offline. The authors should prefer online submission. Online Submission: There are three ways to submit your paper:

**(A) (I) First, register yourself using top right corner of Home page then Login. If you are already registered, then login using your username and password.**

**(II) Choose corresponding Journal.**

**(III) Click 'Submit Manuscript'. Fill required information and Upload the paper.**

**(B) If you are using Internet Explorer, then Direct Submission through Homepage is also available.**

**(C) If these two are not convenient, and then email the paper directly to dean@globaljournals.org.**

Offline Submission: Author can send the typed form of paper by Post. However, online submission should be preferred.



# PREFERRED AUTHOR GUIDELINES

## MANUSCRIPT STYLE INSTRUCTION (Must be strictly followed)

Page Size: 8.27" X 11"

- Left Margin: 0.65
- Right Margin: 0.65
- Top Margin: 0.75
- Bottom Margin: 0.75
- Font type of all text should be Swis 721 Lt BT.
- Paper Title should be of Font Size 24 with one Column section.
- Author Name in Font Size of 11 with one column as of Title.
- Abstract Font size of 9 Bold, "Abstract" word in Italic Bold.
- Main Text: Font size 10 with justified two columns section
- Two Column with Equal Column with of 3.38 and Gaping of .2
- First Character must be three lines Drop capped.
- Paragraph before Spacing of 1 pt and After of 0 pt.
- Line Spacing of 1 pt
- Large Images must be in One Column
- Numbering of First Main Headings (Heading 1) must be in Roman Letters, Capital Letter, and Font Size of 10.
- Numbering of Second Main Headings (Heading 2) must be in Alphabets, Italic, and Font Size of 10.

**You can use your own standard format also.**

### Author Guidelines:

1. General,
2. Ethical Guidelines,
3. Submission of Manuscripts,
4. Manuscript's Category,
5. Structure and Format of Manuscript,
6. After Acceptance.

### 1. GENERAL

Before submitting your research paper, one is advised to go through the details as mentioned in following heads. It will be beneficial, while peer reviewer justify your paper for publication.

### Scope

The Global Journals Inc. (US) welcome the submission of original paper, review paper, survey article relevant to the all the streams of Philosophy and knowledge. The Global Journals Inc. (US) is parental platform for Global Journal of Computer Science and Technology, Researches in Engineering, Medical Research, Science Frontier Research, Human Social Science, Management, and Business organization. The choice of specific field can be done otherwise as following in Abstracting and Indexing Page on this Website. As the all Global

Journals Inc. (US) are being abstracted and indexed (in process) by most of the reputed organizations. Topics of only narrow interest will not be accepted unless they have wider potential or consequences.

## 2. ETHICAL GUIDELINES

Authors should follow the ethical guidelines as mentioned below for publication of research paper and research activities.

Papers are accepted on strict understanding that the material in whole or in part has not been, nor is being, considered for publication elsewhere. If the paper once accepted by Global Journals Inc. (US) and Editorial Board, will become the copyright of the Global Journals Inc. (US).

**Authorship: The authors and coauthors should have active contribution to conception design, analysis and interpretation of findings. They should critically review the contents and drafting of the paper. All should approve the final version of the paper before submission**

The Global Journals Inc. (US) follows the definition of authorship set up by the Global Academy of Research and Development. According to the Global Academy of R&D authorship, criteria must be based on:

- 1) Substantial contributions to conception and acquisition of data, analysis and interpretation of the findings.
- 2) Drafting the paper and revising it critically regarding important academic content.
- 3) Final approval of the version of the paper to be published.

All authors should have been credited according to their appropriate contribution in research activity and preparing paper. Contributors who do not match the criteria as authors may be mentioned under Acknowledgement.

Acknowledgements: Contributors to the research other than authors credited should be mentioned under acknowledgement. The specifications of the source of funding for the research if appropriate can be included. Suppliers of resources may be mentioned along with address.

**Appeal of Decision: The Editorial Board's decision on publication of the paper is final and cannot be appealed elsewhere.**

**Permissions: It is the author's responsibility to have prior permission if all or parts of earlier published illustrations are used in this paper.**

Please mention proper reference and appropriate acknowledgements wherever expected.

If all or parts of previously published illustrations are used, permission must be taken from the copyright holder concerned. It is the author's responsibility to take these in writing.

Approval for reproduction/modification of any information (including figures and tables) published elsewhere must be obtained by the authors/copyright holders before submission of the manuscript. Contributors (Authors) are responsible for any copyright fee involved.

## 3. SUBMISSION OF MANUSCRIPTS

Manuscripts should be uploaded via this online submission page. The online submission is most efficient method for submission of papers, as it enables rapid distribution of manuscripts and consequently speeds up the review procedure. It also enables authors to know the status of their own manuscripts by emailing us. Complete instructions for submitting a paper is available below.

Manuscript submission is a systematic procedure and little preparation is required beyond having all parts of your manuscript in a given format and a computer with an Internet connection and a Web browser. Full help and instructions are provided on-screen. As an author, you will be prompted for login and manuscript details as Field of Paper and then to upload your manuscript file(s) according to the instructions.





To avoid postal delays, all transaction is preferred by e-mail. A finished manuscript submission is confirmed by e-mail immediately and your paper enters the editorial process with no postal delays. When a conclusion is made about the publication of your paper by our Editorial Board, revisions can be submitted online with the same procedure, with an occasion to view and respond to all comments.

Complete support for both authors and co-author is provided.

#### 4. MANUSCRIPT'S CATEGORY

Based on potential and nature, the manuscript can be categorized under the following heads:

Original research paper: Such papers are reports of high-level significant original research work.

Review papers: These are concise, significant but helpful and decisive topics for young researchers.

Research articles: These are handled with small investigation and applications

Research letters: The letters are small and concise comments on previously published matters.

#### 5. STRUCTURE AND FORMAT OF MANUSCRIPT

The recommended size of original research paper is less than seven thousand words, review papers fewer than seven thousands words also. Preparation of research paper or how to write research paper, are major hurdle, while writing manuscript. The research articles and research letters should be fewer than three thousand words, the structure original research paper; sometime review paper should be as follows:

**Papers:** These are reports of significant research (typically less than 7000 words equivalent, including tables, figures, references), and comprise:

(a) Title should be relevant and commensurate with the theme of the paper.

(b) A brief Summary, "Abstract" (less than 150 words) containing the major results and conclusions.

(c) Up to ten keywords, that precisely identifies the paper's subject, purpose, and focus.

(d) An Introduction, giving necessary background excluding subheadings; objectives must be clearly declared.

(e) Resources and techniques with sufficient complete experimental details (wherever possible by reference) to permit repetition; sources of information must be given and numerical methods must be specified by reference, unless non-standard.

(f) Results should be presented concisely, by well-designed tables and/or figures; the same data may not be used in both; suitable statistical data should be given. All data must be obtained with attention to numerical detail in the planning stage. As reproduced design has been recognized to be important to experiments for a considerable time, the Editor has decided that any paper that appears not to have adequate numerical treatments of the data will be returned un-refereed;

(g) Discussion should cover the implications and consequences, not just recapitulating the results; conclusions should be summarizing.

(h) Brief Acknowledgements.

(i) References in the proper form.

Authors should very cautiously consider the preparation of papers to ensure that they communicate efficiently. Papers are much more likely to be accepted, if they are cautiously designed and laid out, contain few or no errors, are summarizing, and be conventional to the approach and instructions. They will in addition, be published with much less delays than those that require much technical and editorial correction.



The Editorial Board reserves the right to make literary corrections and to make suggestions to improve brevity.

It is vital, that authors take care in submitting a manuscript that is written in simple language and adheres to published guidelines.

## Format

*Language: The language of publication is UK English. Authors, for whom English is a second language, must have their manuscript efficiently edited by an English-speaking person before submission to make sure that, the English is of high excellence. It is preferable, that manuscripts should be professionally edited.*

Standard Usage, Abbreviations, and Units: Spelling and hyphenation should be conventional to The Concise Oxford English Dictionary. Statistics and measurements should at all times be given in figures, e.g. 16 min, except for when the number begins a sentence. When the number does not refer to a unit of measurement it should be spelt in full unless, it is 160 or greater.

Abbreviations supposed to be used carefully. The abbreviated name or expression is supposed to be cited in full at first usage, followed by the conventional abbreviation in parentheses.

Metric SI units are supposed to generally be used excluding where they conflict with current practice or are confusing. For illustration, 1.4 l rather than  $1.4 \times 10^{-3} \text{ m}^3$ , or 4 mm somewhat than  $4 \times 10^{-3} \text{ m}$ . Chemical formula and solutions must identify the form used, e.g. anhydrous or hydrated, and the concentration must be in clearly defined units. Common species names should be followed by underlines at the first mention. For following use the generic name should be constricted to a single letter, if it is clear.

## Structure

All manuscripts submitted to Global Journals Inc. (US), ought to include:

Title: The title page must carry an instructive title that reflects the content, a running title (less than 45 characters together with spaces), names of the authors and co-authors, and the place(s) wherever the work was carried out. The full postal address in addition with the e-mail address of related author must be given. Up to eleven keywords or very brief phrases have to be given to help data retrieval, mining and indexing.

*Abstract, used in Original Papers and Reviews:*

### Optimizing Abstract for Search Engines

Many researchers searching for information online will use search engines such as Google, Yahoo or similar. By optimizing your paper for search engines, you will amplify the chance of someone finding it. This in turn will make it more likely to be viewed and/or cited in a further work. Global Journals Inc. (US) have compiled these guidelines to facilitate you to maximize the web-friendliness of the most public part of your paper.

### Key Words

A major linchpin in research work for the writing research paper is the keyword search, which one will employ to find both library and Internet resources.

One must be persistent and creative in using keywords. An effective keyword search requires a strategy and planning a list of possible keywords and phrases to try.

Search engines for most searches, use Boolean searching, which is somewhat different from Internet searches. The Boolean search uses "operators," words (and, or, not, and near) that enable you to expand or narrow your affords. Tips for research paper while preparing research paper are very helpful guideline of research paper.

Choice of key words is first tool of tips to write research paper. Research paper writing is an art. A few tips for deciding as strategically as possible about keyword search:



- One should start brainstorming lists of possible keywords before even begin searching. Think about the most important concepts related to research work. Ask, "What words would a source have to include to be truly valuable in research paper?" Then consider synonyms for the important words.
- It may take the discovery of only one relevant paper to let steer in the right keyword direction because in most databases, the keywords under which a research paper is abstracted are listed with the paper.
- One should avoid outdated words.

Keywords are the key that opens a door to research work sources. Keyword searching is an art in which researcher's skills are bound to improve with experience and time.

Numerical Methods: Numerical methods used should be clear and, where appropriate, supported by references.

*Acknowledgements: Please make these as concise as possible.*

#### References

References follow the Harvard scheme of referencing. References in the text should cite the authors' names followed by the time of their publication, unless there are three or more authors when simply the first author's name is quoted followed by et al. unpublished work has to only be cited where necessary, and only in the text. Copies of references in press in other journals have to be supplied with submitted typescripts. It is necessary that all citations and references be carefully checked before submission, as mistakes or omissions will cause delays.

References to information on the World Wide Web can be given, but only if the information is available without charge to readers on an official site. Wikipedia and Similar websites are not allowed where anyone can change the information. Authors will be asked to make available electronic copies of the cited information for inclusion on the Global Journals Inc. (US) homepage at the judgment of the Editorial Board.

The Editorial Board and Global Journals Inc. (US) recommend that, citation of online-published papers and other material should be done via a DOI (digital object identifier). If an author cites anything, which does not have a DOI, they run the risk of the cited material not being noticeable.

The Editorial Board and Global Journals Inc. (US) recommend the use of a tool such as Reference Manager for reference management and formatting.

#### Tables, Figures and Figure Legends

*Tables: Tables should be few in number, cautiously designed, uncrowned, and include only essential data. Each must have an Arabic number, e.g. Table 4, a self-explanatory caption and be on a separate sheet. Vertical lines should not be used.*

*Figures: Figures are supposed to be submitted as separate files. Always take in a citation in the text for each figure using Arabic numbers, e.g. Fig. 4. Artwork must be submitted online in electronic form by e-mailing them.*

#### Preparation of Electronic Figures for Publication

Even though low quality images are sufficient for review purposes, print publication requires high quality images to prevent the final product being blurred or fuzzy. Submit (or e-mail) EPS (line art) or TIFF (halftone/photographs) files only. MS PowerPoint and Word Graphics are unsuitable for printed pictures. Do not use pixel-oriented software. Scans (TIFF only) should have a resolution of at least 350 dpi (halftone) or 700 to 1100 dpi (line drawings) in relation to the imitation size. Please give the data for figures in black and white or submit a Color Work Agreement Form. EPS files must be saved with fonts embedded (and with a TIFF preview, if possible).

For scanned images, the scanning resolution (at final image size) ought to be as follows to ensure good reproduction: line art: >650 dpi; halftones (including gel photographs) : >350 dpi; figures containing both halftone and line images: >650 dpi.



Color Charges: It is the rule of the Global Journals Inc. (US) for authors to pay the full cost for the reproduction of their color artwork. Hence, please note that, if there is color artwork in your manuscript when it is accepted for publication, we would require you to complete and return a color work agreement form before your paper can be published.

*Figure Legends: Self-explanatory legends of all figures should be incorporated separately under the heading 'Legends to Figures'. In the full-text online edition of the journal, figure legends may possibly be truncated in abbreviated links to the full screen version. Therefore, the first 100 characters of any legend should notify the reader, about the key aspects of the figure.*

## **6. AFTER ACCEPTANCE**

Upon approval of a paper for publication, the manuscript will be forwarded to the dean, who is responsible for the publication of the Global Journals Inc. (US).

### **6.1 Proof Corrections**

The corresponding author will receive an e-mail alert containing a link to a website or will be attached. A working e-mail address must therefore be provided for the related author.

Acrobat Reader will be required in order to read this file. This software can be downloaded

(Free of charge) from the following website:

[www.adobe.com/products/acrobat/readstep2.html](http://www.adobe.com/products/acrobat/readstep2.html). This will facilitate the file to be opened, read on screen, and printed out in order for any corrections to be added. Further instructions will be sent with the proof.

Proofs must be returned to the dean at [dean@globaljournals.org](mailto:dean@globaljournals.org) within three days of receipt.

As changes to proofs are costly, we inquire that you only correct typesetting errors. All illustrations are retained by the publisher. Please note that the authors are responsible for all statements made in their work, including changes made by the copy editor.

### **6.2 Early View of Global Journals Inc. (US) (Publication Prior to Print)**

The Global Journals Inc. (US) are enclosed by our publishing's Early View service. Early View articles are complete full-text articles sent in advance of their publication. Early View articles are absolute and final. They have been completely reviewed, revised and edited for publication, and the authors' final corrections have been incorporated. Because they are in final form, no changes can be made after sending them. The nature of Early View articles means that they do not yet have volume, issue or page numbers, so Early View articles cannot be cited in the conventional way.

### **6.3 Author Services**

Online production tracking is available for your article through Author Services. Author Services enables authors to track their article - once it has been accepted - through the production process to publication online and in print. Authors can check the status of their articles online and choose to receive automated e-mails at key stages of production. The authors will receive an e-mail with a unique link that enables them to register and have their article automatically added to the system. Please ensure that a complete e-mail address is provided when submitting the manuscript.

### **6.4 Author Material Archive Policy**

Please note that if not specifically requested, publisher will dispose off hardcopy & electronic information submitted, after the two months of publication. If you require the return of any information submitted, please inform the Editorial Board or dean as soon as possible.

### **6.5 Offprint and Extra Copies**

A PDF offprint of the online-published article will be provided free of charge to the related author, and may be distributed according to the Publisher's terms and conditions. Additional paper offprint may be ordered by emailing us at: [editor@globaljournals.org](mailto:editor@globaljournals.org) .



Before start writing a good quality Computer Science Research Paper, let us first understand what is Computer Science Research Paper? So, Computer Science Research Paper is the paper which is written by professionals or scientists who are associated to Computer Science and Information Technology, or doing research study in these areas. If you are novel to this field then you can consult about this field from your supervisor or guide.

#### TECHNIQUES FOR WRITING A GOOD QUALITY RESEARCH PAPER:

**1. Choosing the topic:** In most cases, the topic is searched by the interest of author but it can be also suggested by the guides. You can have several topics and then you can judge that in which topic or subject you are finding yourself most comfortable. This can be done by asking several questions to yourself, like Will I be able to carry our search in this area? Will I find all necessary recourses to accomplish the search? Will I be able to find all information in this field area? If the answer of these types of questions will be "Yes" then you can choose that topic. In most of the cases, you may have to conduct the surveys and have to visit several places because this field is related to Computer Science and Information Technology. Also, you may have to do a lot of work to find all rise and falls regarding the various data of that subject. Sometimes, detailed information plays a vital role, instead of short information.

**2. Evaluators are human:** First thing to remember that evaluators are also human being. They are not only meant for rejecting a paper. They are here to evaluate your paper. So, present your Best.

**3. Think Like Evaluators:** If you are in a confusion or getting demotivated that your paper will be accepted by evaluators or not, then think and try to evaluate your paper like an Evaluator. Try to understand that what an evaluator wants in your research paper and automatically you will have your answer.

**4. Make blueprints of paper:** The outline is the plan or framework that will help you to arrange your thoughts. It will make your paper logical. But remember that all points of your outline must be related to the topic you have chosen.

**5. Ask your Guides:** If you are having any difficulty in your research, then do not hesitate to share your difficulty to your guide (if you have any). They will surely help you out and resolve your doubts. If you can't clarify what exactly you require for your work then ask the supervisor to help you with the alternative. He might also provide you the list of essential readings.

**6. Use of computer is recommended:** As you are doing research in the field of Computer Science, then this point is quite obvious.

**7. Use right software:** Always use good quality software packages. If you are not capable to judge good software then you can lose quality of your paper unknowingly. There are various software programs available to help you, which you can get through Internet.

**8. Use the Internet for help:** An excellent start for your paper can be by using the Google. It is an excellent search engine, where you can have your doubts resolved. You may also read some answers for the frequent question how to write my research paper or find model research paper. From the internet library you can download books. If you have all required books make important reading selecting and analyzing the specified information. Then put together research paper sketch out.

**9. Use and get big pictures:** Always use encyclopedias, Wikipedia to get pictures so that you can go into the depth.

**10. Bookmarks are useful:** When you read any book or magazine, you generally use bookmarks, right! It is a good habit, which helps to not to lose your continuity. You should always use bookmarks while searching on Internet also, which will make your search easier.

**11. Revise what you wrote:** When you write anything, always read it, summarize it and then finalize it.



**12. Make all efforts:** Make all efforts to mention what you are going to write in your paper. That means always have a good start. Try to mention everything in introduction, that what is the need of a particular research paper. Polish your work by good skill of writing and always give an evaluator, what he wants.

**13. Have backups:** When you are going to do any important thing like making research paper, you should always have backup copies of it either in your computer or in paper. This will help you to not to lose any of your important.

**14. Produce good diagrams of your own:** Always try to include good charts or diagrams in your paper to improve quality. Using several and unnecessary diagrams will degrade the quality of your paper by creating "hotchpotch." So always, try to make and include those diagrams, which are made by your own to improve readability and understandability of your paper.

**15. Use of direct quotes:** When you do research relevant to literature, history or current affairs then use of quotes become essential but if study is relevant to science then use of quotes is not preferable.

**16. Use proper verb tense:** Use proper verb tenses in your paper. Use past tense, to present those events that happened. Use present tense to indicate events that are going on. Use future tense to indicate future happening events. Use of improper and wrong tenses will confuse the evaluator. Avoid the sentences that are incomplete.

**17. Never use online paper:** If you are getting any paper on Internet, then never use it as your research paper because it might be possible that evaluator has already seen it or maybe it is outdated version.

**18. Pick a good study spot:** To do your research studies always try to pick a spot, which is quiet. Every spot is not for studies. Spot that suits you choose it and proceed further.

**19. Know what you know:** Always try to know, what you know by making objectives. Else, you will be confused and cannot achieve your target.

**20. Use good quality grammar:** Always use a good quality grammar and use words that will throw positive impact on evaluator. Use of good quality grammar does not mean to use tough words, that for each word the evaluator has to go through dictionary. Do not start sentence with a conjunction. Do not fragment sentences. Eliminate one-word sentences. Ignore passive voice. Do not ever use a big word when a diminutive one would suffice. Verbs have to be in agreement with their subjects. Prepositions are not expressions to finish sentences with. It is incorrect to ever divide an infinitive. Avoid clichés like the disease. Also, always shun irritating alliteration. Use language that is simple and straight forward. put together a neat summary.

**21. Arrangement of information:** Each section of the main body should start with an opening sentence and there should be a changeover at the end of the section. Give only valid and powerful arguments to your topic. You may also maintain your arguments with records.

**22. Never start in last minute:** Always start at right time and give enough time to research work. Leaving everything to the last minute will degrade your paper and spoil your work.

**23. Multitasking in research is not good:** Doing several things at the same time proves bad habit in case of research activity. Research is an area, where everything has a particular time slot. Divide your research work in parts and do particular part in particular time slot.

**24. Never copy others' work:** Never copy others' work and give it your name because if evaluator has seen it anywhere you will be in trouble.

**25. Take proper rest and food:** No matter how many hours you spend for your research activity, if you are not taking care of your health then all your efforts will be in vain. For a quality research, study is must, and this can be done by taking proper rest and food.

**26. Go for seminars:** Attend seminars if the topic is relevant to your research area. Utilize all your resources.



**27. Refresh your mind after intervals:** Try to give rest to your mind by listening to soft music or by sleeping in intervals. This will also improve your memory.

**28. Make colleagues:** Always try to make colleagues. No matter how sharper or intelligent you are, if you make colleagues you can have several ideas, which will be helpful for your research.

**29. Think technically:** Always think technically. If anything happens, then search its reasons, its benefits, and demerits.

**30. Think and then print:** When you will go to print your paper, notice that tables are not be split, headings are not detached from their descriptions, and page sequence is maintained.

**31. Adding unnecessary information:** Do not add unnecessary information, like, I have used MS Excel to draw graph. Do not add irrelevant and inappropriate material. These all will create superfluous. Foreign terminology and phrases are not apropos. One should NEVER take a broad view. Analogy in script is like feathers on a snake. Not at all use a large word when a very small one would be sufficient. Use words properly, regardless of how others use them. Remove quotations. Puns are for kids, not grunt readers. Amplification is a billion times of inferior quality than sarcasm.

**32. Never oversimplify everything:** To add material in your research paper, never go for oversimplification. This will definitely irritate the evaluator. Be more or less specific. Also too, by no means, ever use rhythmic redundancies. Contractions aren't essential and shouldn't be there used. Comparisons are as terrible as clichés. Give up ampersands and abbreviations, and so on. Remove commas, that are, not necessary. Parenthetical words however should be together with this in commas. Understatement is all the time the complete best way to put onward earth-shaking thoughts. Give a detailed literary review.

**33. Report concluded results:** Use concluded results. From raw data, filter the results and then conclude your studies based on measurements and observations taken. Significant figures and appropriate number of decimal places should be used. Parenthetical remarks are prohibitive. Proofread carefully at final stage. In the end give outline to your arguments. Spot out perspectives of further study of this subject. Justify your conclusion by at the bottom of them with sufficient justifications and examples.

**34. After conclusion:** Once you have concluded your research, the next most important step is to present your findings. Presentation is extremely important as it is the definite medium through which your research is going to be in print to the rest of the crowd. Care should be taken to categorize your thoughts well and present them in a logical and neat manner. A good quality research paper format is essential because it serves to highlight your research paper and bring to light all necessary aspects in your research.

## INFORMAL GUIDELINES OF RESEARCH PAPER WRITING

### Key points to remember:

- Submit all work in its final form.
- Write your paper in the form, which is presented in the guidelines using the template.
- Please note the criterion for grading the final paper by peer-reviewers.

### Final Points:

A purpose of organizing a research paper is to let people to interpret your effort selectively. The journal requires the following sections, submitted in the order listed, each section to start on a new page.

The introduction will be compiled from reference matter and will reflect the design processes or outline of basis that direct you to make study. As you will carry out the process of study, the method and process section will be constructed as like that. The result segment will show related statistics in nearly sequential order and will direct the reviewers next to the similar intellectual paths throughout the data that you took to carry out your study. The discussion section will provide understanding of the data and projections as to the implication of the results. The use of good quality references all through the paper will give the effort trustworthiness by representing an alertness of prior workings.



Writing a research paper is not an easy job no matter how trouble-free the actual research or concept. Practice, excellent preparation, and controlled record keeping are the only means to make straightforward the progression.

**General style:**

Specific editorial column necessities for compliance of a manuscript will always take over from directions in these general guidelines.

To make a paper clear

- Adhere to recommended page limits

Mistakes to evade

- Insertion a title at the foot of a page with the subsequent text on the next page
- Separating a table/chart or figure - impound each figure/table to a single page
- Submitting a manuscript with pages out of sequence

In every sections of your document

- Use standard writing style including articles ("a", "the," etc.)
- Keep on paying attention on the research topic of the paper
- Use paragraphs to split each significant point (excluding for the abstract)
- Align the primary line of each section
- Present your points in sound order
- Use present tense to report well accepted
- Use past tense to describe specific results
- Shun familiar wording, don't address the reviewer directly, and don't use slang, slang language, or superlatives
- Shun use of extra pictures - include only those figures essential to presenting results

**Title Page:**

Choose a revealing title. It should be short. It should not have non-standard acronyms or abbreviations. It should not exceed two printed lines. It should include the name(s) and address (es) of all authors.





## Abstract:

The summary should be two hundred words or less. It should briefly and clearly explain the key findings reported in the manuscript-- must have precise statistics. It should not have abnormal acronyms or abbreviations. It should be logical in itself. Shun citing references at this point.

An abstract is a brief distinct paragraph summary of finished work or work in development. In a minute or less a reviewer can be taught the foundation behind the study, common approach to the problem, relevant results, and significant conclusions or new questions.

Write your summary when your paper is completed because how can you write the summary of anything which is not yet written? Wealth of terminology is very essential in abstract. Yet, use comprehensive sentences and do not let go readability for brevity. You can maintain it succinct by phrasing sentences so that they provide more than lone rationale. The author can at this moment go straight to shortening the outcome. Sum up the study, with the subsequent elements in any summary. Try to maintain the initial two items to no more than one ruling each.

- Reason of the study - theory, overall issue, purpose
- Fundamental goal
- To the point depiction of the research
- Consequences, including definite statistics - if the consequences are quantitative in nature, account quantitative data; results of any numerical analysis should be reported
- Significant conclusions or questions that track from the research(es)

## Approach:

- Single section, and succinct
- As an outline of job done, it is always written in past tense
- A conceptual should situate on its own, and not submit to any other part of the paper such as a form or table
- Center on shortening results - bound background information to a verdict or two, if completely necessary
- What you account in an abstract must be regular with what you reported in the manuscript
- Exact spelling, clearness of sentences and phrases, and appropriate reporting of quantities (proper units, important statistics) are just as significant in an abstract as they are anywhere else

## Introduction:

The **Introduction** should "introduce" the manuscript. The reviewer should be presented with sufficient background information to be capable to comprehend and calculate the purpose of your study without having to submit to other works. The basis for the study should be offered. Give most important references but shun difficult to make a comprehensive appraisal of the topic. In the introduction, describe the problem visibly. If the problem is not acknowledged in a logical, reasonable way, the reviewer will have no attention in your result. Speak in common terms about techniques used to explain the problem, if needed, but do not present any particulars about the protocols here. Following approach can create a valuable beginning:

- Explain the value (significance) of the study
- Shield the model - why did you employ this particular system or method? What is its compensation? You strength remark on its appropriateness from a abstract point of vision as well as point out sensible reasons for using it.
- Present a justification. Status your particular theory (es) or aim(s), and describe the logic that led you to choose them.
- Very for a short time explain the tentative propose and how it skilled the declared objectives.

## Approach:

- Use past tense except for when referring to recognized facts. After all, the manuscript will be submitted after the entire job is done.
- Sort out your thoughts; manufacture one key point with every section. If you make the four points listed above, you will need a least of four paragraphs.



- Present surroundings information only as desirable in order hold up a situation. The reviewer does not desire to read the whole thing you know about a topic.
- Shape the theory/purpose specifically - do not take a broad view.
- As always, give awareness to spelling, simplicity and correctness of sentences and phrases.

#### **Procedures (Methods and Materials):**

This part is supposed to be the easiest to carve if you have good skills. A sound written Procedures segment allows a capable scientist to replacement your results. Present precise information about your supplies. The suppliers and clarity of reagents can be helpful bits of information. Present methods in sequential order but linked methodologies can be grouped as a segment. Be concise when relating the protocols. Attempt for the least amount of information that would permit another capable scientist to spare your outcome but be cautious that vital information is integrated. The use of subheadings is suggested and ought to be synchronized with the results section. When a technique is used that has been well described in another object, mention the specific item describing a way but draw the basic principle while stating the situation. The purpose is to text all particular resources and broad procedures, so that another person may use some or all of the methods in one more study or referee the scientific value of your work. It is not to be a step by step report of the whole thing you did, nor is a methods section a set of orders.

#### **Materials:**

- Explain materials individually only if the study is so complex that it saves liberty this way.
- Embrace particular materials, and any tools or provisions that are not frequently found in laboratories.
- Do not take in frequently found.
- If use of a definite type of tools.
- Materials may be reported in a part section or else they may be recognized along with your measures.

#### **Methods:**

- Report the method (not particulars of each process that engaged the same methodology)
- Describe the method entirely
- To be succinct, present methods under headings dedicated to specific dealings or groups of measures
- Simplify - details how procedures were completed not how they were exclusively performed on a particular day.
- If well known procedures were used, account the procedure by name, possibly with reference, and that's all.

#### **Approach:**

- It is embarrassed or not possible to use vigorous voice when documenting methods with no using first person, which would focus the reviewer's interest on the researcher rather than the job. As a result when script up the methods most authors use third person passive voice.
- Use standard style in this and in every other part of the paper - avoid familiar lists, and use full sentences.

#### **What to keep away from**

- Resources and methods are not a set of information.
- Skip all descriptive information and surroundings - save it for the argument.
- Leave out information that is immaterial to a third party.

#### **Results:**

The principle of a results segment is to present and demonstrate your conclusion. Create this part a entirely objective details of the outcome, and save all understanding for the discussion.

The page length of this segment is set by the sum and types of data to be reported. Carry on to be to the point, by means of statistics and tables, if suitable, to present consequences most efficiently. You must obviously differentiate material that would usually be incorporated in a study editorial from any unprocessed data or additional appendix matter that would not be available. In fact, such matter should not be submitted at all except requested by the instructor.



## Content

- Sum up your conclusion in text and demonstrate them, if suitable, with figures and tables.
- In manuscript, explain each of your consequences, point the reader to remarks that are most appropriate.
- Present a background, such as by describing the question that was addressed by creation an exacting study.
- Explain results of control experiments and comprise remarks that are not accessible in a prescribed figure or table, if appropriate.
- Examine your data, then prepare the analyzed (transformed) data in the form of a figure (graph), table, or in manuscript form.

### What to stay away from

- Do not discuss or infer your outcome, report surroundings information, or try to explain anything.
- Not at all, take in raw data or intermediate calculations in a research manuscript.
- Do not present the similar data more than once.
- Manuscript should complement any figures or tables, not duplicate the identical information.
- Never confuse figures with tables - there is a difference.

### Approach

- As forever, use past tense when you submit to your results, and put the whole thing in a reasonable order.
- Put figures and tables, appropriately numbered, in order at the end of the report
- If you desire, you may place your figures and tables properly within the text of your results part.

### Figures and tables

- If you put figures and tables at the end of the details, make certain that they are visibly distinguished from any attach appendix materials, such as raw facts
- Despite of position, each figure must be numbered one after the other and complete with subtitle
- In spite of position, each table must be titled, numbered one after the other and complete with heading
- All figure and table must be adequately complete that it could situate on its own, divide from text

### Discussion:

The Discussion is expected the trickiest segment to write and describe. A lot of papers submitted for journal are discarded based on problems with the Discussion. There is no head of state for how long a argument should be. Position your understanding of the outcome visibly to lead the reviewer through your conclusions, and then finish the paper with a summing up of the implication of the study. The purpose here is to offer an understanding of your results and hold up for all of your conclusions, using facts from your research and generally accepted information, if suitable. The implication of result should be visibly described. Infer your data in the conversation in suitable depth. This means that when you clarify an observable fact you must explain mechanisms that may account for the observation. If your results vary from your prospect, make clear why that may have happened. If your results agree, then explain the theory that the proof supported. It is never suitable to just state that the data approved with prospect, and let it drop at that.

- Make a decision if each premise is supported, discarded, or if you cannot make a conclusion with assurance. Do not just dismiss a study or part of a study as "uncertain."
- Research papers are not acknowledged if the work is imperfect. Draw what conclusions you can based upon the results that you have, and take care of the study as a finished work
- You may propose future guidelines, such as how the experiment might be personalized to accomplish a new idea.
- Give details all of your remarks as much as possible, focus on mechanisms.
- Make a decision if the tentative design sufficiently addressed the theory, and whether or not it was correctly restricted.
- Try to present substitute explanations if sensible alternatives be present.
- One research will not counter an overall question, so maintain the large picture in mind, where do you go next? The best studies unlock new avenues of study. What questions remain?
- Recommendations for detailed papers will offer supplementary suggestions.

### Approach:

- When you refer to information, differentiate data generated by your own studies from available information
- Submit to work done by specific persons (including you) in past tense.
- Submit to generally acknowledged facts and main beliefs in present tense.



## THE ADMINISTRATION RULES

Please carefully note down following rules and regulation before submitting your Research Paper to Global Journals Inc. (US):

**Segment Draft and Final Research Paper:** You have to strictly follow the template of research paper. If it is not done your paper may get rejected.

- The **major constraint** is that you must independently make all content, tables, graphs, and facts that are offered in the paper. You must write each part of the paper wholly on your own. The Peer-reviewers need to identify your own perceptives of the concepts in your own terms. NEVER extract straight from any foundation, and never rephrase someone else's analysis.
- Do not give permission to anyone else to "PROOFREAD" your manuscript.
- **Methods to avoid Plagiarism is applied by us on every paper, if found guilty, you will be blacklisted by all of our collaborated research groups, your institution will be informed for this and strict legal actions will be taken immediately.)**
- To guard yourself and others from possible illegal use please do not permit anyone right to use to your paper and files.



CRITERION FOR GRADING A RESEARCH PAPER (COMPILATION)  
BY GLOBAL JOURNALS INC. (US)

Please note that following table is only a Grading of "Paper Compilation" and not on "Performed/Stated Research" whose grading solely depends on Individual Assigned Peer Reviewer and Editorial Board Member. These can be available only on request and after decision of Paper. This report will be the property of Global Journals Inc. (US).

| Topics                        | Grades   |   |  |
|-------------------------------|--|---|--|
|                               | A-B  | C-D   | E-F  |
| <i>Abstract</i>               | Clear and concise with appropriate content, Correct format. 200 words or below   | Unclear summary and no specific data, Incorrect form<br><br>Above 200 words                         | No specific data with ambiguous information<br><br>Above 250 words |
| <i>Introduction</i>           | Containing all background details with clear goal and appropriate details, flow specification, no grammar and spelling mistake, well organized sentence and paragraph, reference cited | Unclear and confusing data, appropriate format, grammar and spelling errors with unorganized matter | Out of place depth and content, hazy format                        |
| <i>Methods and Procedures</i> | Clear and to the point with well arranged paragraph, precision and accuracy of facts and figures, well organized subheads  | Difficult to comprehend with embarrassed text, too much explanation but completed                   | Incorrect and unorganized structure with hazy meaning              |
| <i>Result</i>                 | Well organized, Clear and specific, Correct units with precision, correct data, well structuring of paragraph, no grammar and spelling mistake   | Complete and embarrassed text, difficult to comprehend  | Irregular format with wrong facts and figures                      |
| <i>Discussion</i>             | Well organized, meaningful specification, sound conclusion, logical and concise explanation, highly structured paragraph reference cited   | Wordy, unclear conclusion, spurious   | Conclusion is not cited, unorganized, difficult to comprehend      |
| <i>References</i>             | Complete and correct format, well organized  | Beside the point, Incomplete  | Wrong format and structuring                                       |



# INDEX

---

---

## **A**

Angiotensin · 12  
Antepartum · 4

---

## **B**

Bilirubin · 14, 29  
Bougangue · 17, 18, 23, 24, 25

---

## **C**

Counselled · 7  
Cystadenoma · 8  
Cystectomy · 1, 7, 9

---

## **D**

Darmstadt · 23, 25  
Dystocia · 20

---

## **E**

Eclampsia · 1, 2, 4, 6, 11  
Ekpoudom · 13  
Erythematosis · 12  
Etiology · 11

---

## **H**

Haemolytic · 29  
Hypertensive · 1, 4, 11, 13, 14, 15, 29

---

## **I**

Intrauterine · 4, 7, 15

---

## **L**

Laparotomies · 32  
Leiomyosarcomas · 33

---

## **M**

Marginalising · 22  
Meticulous · 4

---

## **N**

Nicardipine · 29  
Normotensive · 14  
Nwokocha · 18, 26

---

## **O**

Oliguria · 11  
Oophoropexy · 9

---

## **P**

Preeclampsia · 1, 29, 30  
Primigravida · 13  
Proteinuria · 11, 30

---

## **S**

Salphingoophorectomy · 8  
Sarcoma · 1, 31, 33

---

## **T**

Tertiary · 1, 2, 4, 13  
Thrombocytopenia · 11



save our planet




# Global Journal of Medical Research

---

Visit us on the Web at [www.GlobalJournals.org](http://www.GlobalJournals.org) | [www.MedicalResearchJournal.org](http://www.MedicalResearchJournal.org)  
or email us at [helpdesk@globaljournals.org](mailto:helpdesk@globaljournals.org)

ISSN 9755896



© Global Journals