



GLOBAL JOURNAL OF MEDICAL RESEARCH: I  
SURGERIES AND CARDIOVASCULAR SYSTEM  
Volume 18 Issue 2 Version 1.0 Year 2018  
Type: Double Blind Peer Reviewed International Research Journal  
Publisher: Global Journals  
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

# The Role of 2D Speckle Tracking Echocardiography in Early Detection of Left Ventricular Dysfunction in Type II Diabetic Patients

By Mahmoud Shawky Abd El Moneum

*Benha University*

**Abstract- Aim:** The Early detection of diabetic heart disease is important for the timely interventions resulting in the prevention for the future development of heart failure. Left ventricular (LV) systolic dysfunction may be identified by a reduction in longitudinal function, which can be assessed using 2D speckle tracking echocardiography (STE).

**Methods and Results:** To determine longitudinal, radial, and circumferential function, three LV short-axis and three LV apical views were acquired in 100 asymptomatic diabetic patients with normal LV ejection fraction (EF) and 25 age-matched healthy volunteers. Using 2D strain software, end-systolic longitudinal strain (LS), radial strain (RS), and circumferential strain (CS) were measured in 18 LV segments. No significant differences in LVEF were noted between two groups. Diabetic patients had more advanced diastolic dysfunction and increased LV mass compared with normal subjects. Basal, middle, and apical LSs were significantly lower in diabetic patients compared with control subjects, with 43% (43/100) of the diabetic patients showing abnormal global LS values (cut-off value: 217.2, mean  $\pm$  2 SD in control subjects).

**Keywords:** *left ventricle; speckle tracking; longitudinal function; strain; diabetes mellitus.*

**GJMR-I Classification:** FOR Code: WG 141.5.E2



*Strictly as per the compliance and regulations of:*



# The Role of 2D Speckle Tracking Echocardiography in Early Detection of Left Ventricular Dysfunction in Type II Diabetic Patients

Mahmoud Shawky Abd El Moneum

**Abstract- Aim:** The Early detection of diabetic heart disease is important for the timely interventions resulting in the prevention for the future development of heart failure. Left ventricular (LV) systolic dysfunction may be identified by a reduction in longitudinal function, which can be assessed using 2D speckle tracking echocardiography (STE).

**Methods and Results:** To determine longitudinal, radial, and circumferential function, three LV short-axis and three LV apical views were acquired in 100 asymptomatic diabetic patients with normal LV ejection fraction (EF) and 25 age-matched healthy volunteers. Using 2D strain software, end-systolic longitudinal strain (LS), radial strain (RS), and circumferential strain (CS) were measured in 18 LV segments. No significant differences in LVEF were noted between two groups. Diabetic patients had more advanced diastolic dysfunction and increased LV mass compared with normal subjects. Basal, middle, and apical LSs were significantly lower in diabetic patients compared with control subjects, with 43% (43/100) of the diabetic patients showing abnormal global LS values (cut-off value: 217.2, mean  $\pm$  2 SD in control subjects). Basal RS and apical CS were also significantly lower in diabetic patients. Multivariate linear regression analysis showed that diabetic duration was the only independent confounder for the reduction of LS ( $t$   $\frac{1}{4}$  2.22,  $P$   $\frac{1}{4}$  0.0313).

**Conclusion:** In addition to diastolic dysfunction, LV longitudinal dysfunction is preferentially and frequently observed in asymptomatic diabetes patients with normal LVEF. The decrease in LS correlated with duration of diabetes. 2D speckle tracking echocardiography (STE) has the potential for detecting subclinical LV systolic dysfunction and might provide useful information of the risk stratification in an asymptomatic diabetic population.

**Keywords:** left ventricle; speckle tracking; longitudinal function; strain; diabetes mellitus.

## I. INTRODUCTION

DM is often associated with coronary risk factors, resulting in significant cardiac morbidity and mortality.[1,2] The Early detection of diabetic heart disease is of paramount importance, because timely life-style modifications and medical interventions could prevent or delay the subsequent development of heart failure which is considered one of major burdens for health insurance costs.[3,4] Diabetic patients with normal left ventricular ejection fraction (LVEF) are

frequently associated with diastolic dysfunction.[5,6] However, LVEF is known not to be a sensitive marker for the detection of subclinical LV systolic dysfunction.[7] The Early manifestation of diabetic LV systolic dysfunction can be appeared longitudinally, because sub-endocardial fibres, which are prone to vulnerable to myocardial ischaemia, have a longitudinal trajectory.[8,10] Presence of impaired longitudinal function in diabetic patients has been reported when using tissue Doppler imaging.[11] But tissue Doppler imaging has its own limitations including angle dependency and the 1D nature of its measurement. The recent development of 2D speckle tracking echocardiography (STE) overcomes some of these limitations, and its accuracy [12,13] and clinical usefulness[14,15] have been reported. Assessment of longitudinal strain (LS) profile curve also provides information on post-systolic shortening (PSS), which is considered as a marker of myocardial dysfunction. [16,17] Thus, the aim of this study was mainly to measure LS, radial strain (RS), and circumferential strain (CS) in asymptomatic diabetic patients using 2DSTE, to determine which LV strain is preferentially impaired, and finally to elucidate the characteristics of PSS in a diabetic population.

## II. PATIENTS AND METHODS

### a) Study population

Our study included 100 patients with diabetes mellitus (54 males and 46 females; mean age 63 $\pm$ 12 years). All patients had normal LVEF with no regional wall motion abnormalities on 2D echocardiography. Exclusion criteria included a history of coronary artery disease, the presence of moderate-to-severe valvular heart disease, and/or significant rhythm disturbances. We also enrolled 25 healthy control subjects (15 males and 10 females; mean age 62 $\pm$ 11 years) from our database for normal subjects. Healthy subjects were predominantly hospital employee or their relatives and/or friends. Because ageing affects diastolic function, we selected control subjects in order to adjust the same range of age. The Ethics Committee of the hospital approved the protocol and informed consent was obtained in every subject.

**Author:** MD cardiology, Faculty of Medicine, Benha University, Egypt.  
e-mail: husseine86@gmail.com

b) *Echocardiography*

The Echocardiography was performed using a commercially available ultrasound equipment (M3S probe, Vivid S7.). All 2D grey-scale echocardiographic images were obtained using second harmonic imaging. LV volumes and EF was measured using the modified Simpson method from the apical four- and two-chamber views. For the assessment of LV RS and CS, three LV short-axis planes were acquired at the basal, middle, and apical levels of the LV at high frame rates (range: 67–92 frame/s; mean 81+5 frame/s). Care was taken to ensure that the basal short-axis views were obtained at the level of the mitral valve, the middle planes at the level of the papillary muscles, and apical planes distal to the papillary muscles. For LS assessment, three LV apical views, apical four-chamber, two chamber, and long-axis views were acquired at high frame rates (range: 59–82 frame/s; mean 72+6 frame/s). In each plane, three consecutive cardiac cycles were acquired during a breath hold and digitally stored in a hard disk for off-line analysis. In order to measure the timing of cardiac events, LV inflow and outflow velocities were recorded using pulsed-wave Doppler echocardiography. Mitral annular velocity at the septal corner of the mitral annulus was also recorded to determine peak systolic, early diastolic, and late diastolic annular velocities.

c) *Two-dimensional speckle tracking analysis*

By using commercially available 2D strain software (Echopac PC, version 6.0), the endocardial border in the end-systolic frame was manually traced. A region of interest was then drawn to include the entire myocardium. The software algorithm automatically segmented the LV into six equidistant segments and selected suitable speckles in the myocardium for tracking. The software algorithm then tracked the speckle patterns on a frame-by-frame basis using the sum of absolute difference algorithm. Finally, the software automatically generated time-domain LV strain profiles for each of the six segments of each view, from which end-systolic strain was measured. The average value of strain at each level (basal, middle, and apical) and global strain obtained from averaging the strain values of 18 LV segments was calculated. We also evaluated longitudinal PSS. From time-domain LS waveforms throughout the cardiac cycle, we measured strain at end-systole (LSes) as well as post-systolic peak LS (LSpss). The post-systolic index (PSI) was calculated as  $((LSpss - LSes) / LSes) \times 100$  (%) in each segment, and these values displayed in a parametric bull's eye map (Figure 1). Whenever PSS was not present, PSI value of 0 was given.




Figure 1: Measurements of strain and post-systolic index. Upper three panels show longitudinal regional strain curve of six segments from the apical four-chamber, two-chamber, and long-axis views in a diabetic patient. The vertical line denotes aortic valve closure. In addition to longitudinal strain values at end-systole, post-systolic peak longitudinal strain was measured, if the regional strain curve reached its peak after the aortic valve closure. Lower panels show parametric image of end-systolic stain and post-systolic index. Note that post-systolic index was observed in the basal part of the myocardium.

d) *Statistical analysis*

Our study was designed with 90% power to detect a significant difference in global LS between diabetic patients and control subjects with a  $\frac{1}{4}$  0.05. A difference in global LS between two groups of 3.0% was defined as clinically important, with an estimated SD of 3.0%. Due to reliability of multivariate analysis in diabetic patients, we enrolled 100 diabetic patients. Data were expressed as mean values +SD or median values (inter-quartile range). Frequencies were expressed as percentages. All statistical analysis was carried out using commercially available statistical software (JMP, version 7.0 SAS). Differences in continuous variables between both groups were evaluated using paired or unpaired t-tests. Categorical variables were compared using Fisher's exact test or  $\chi^2$  test whenever appropriate. Linear regression analysis was used to investigate the relation between two parameters. Univariate and multivariate analyses were performed to

determine independent predictors between LS and clinical and echocardiographic parameters. A P-value of 0.05 was considered significant.

III. RESULTS

a) *Clinical and standard echocardiographic characteristics*

**Table (1)** shows the clinical characteristics of both groups. The mean diabetic duration was 8.7 years.

**Table (2)** shows standard echocardiographic parameters. Although LVEF was not different between groups, LV mass index, relative wall thickness, and left atrial volume index were significantly higher in diabetic patients. Peak systolic and early diastolic annular velocity (E0) was significantly lower in the diabetic group, resulting in a higher E/E0 compared with control subjects.

*Table 1:* Clinical characteristics of diabetic patients and control subjects

	Diabetic patients (n = 100)	Control subjects (n = 25)	P-value
Age (years)	63 ± 12	62 ± 11	NS
Sex, male/female (N)	54/46	15/10	NS
BSA (m <sup>2</sup> )	1.60 ± 0.19	1.61 ± 0.15	NS
HR (bpm)	73 ± 13	67 ± 9	<0.05
HTN (N)	47	0 (0)	<0.001
Hyperlipidemia (%)	47(47)	8 (32)	NS
Smoker (%)	49 (49)	6 (24)	<0.05
Diabetic treatment			
Insulin (%)	51 (51)	N/A	
SU (%)	49 (49)	N/A	

BSA, body surface area; DM, diabetes mellitus; HbA1c, haemoglobin A1c; HL, hypertension; HR, heart rate; HTN, hypertension; SU, sulfonil urea.

*Table 2:* Standard echocardiographic data

	Diabetic patients (n = 100)	Control subjects (n = 25)	P-value
IVS (mm)	11.2 ± 1.6	9.4 ± 0.9	<0.001
PW (mm)	10.8 ± 1.5	9.5 ± 1.2	<0.001
LVDd (mm)	46 ± 6	44 ± 4	NS
LVDs (mm)	29 ± 5	28 ± 3	NS
LVEDV (mL)	71.7 ± 20.7	81.2 ± 17.7	<0.05
LVESV (mL)	26.2 ± 11.3	29.1 ± 7.6	NS
LVEF (%)	64.4 ± 7.2	64.2 ± 5.7	NS
LVMI (M-mode) (g/m <sup>2</sup> )	115 ± 30	80 ± 16	<0.001
RWT	0.49 ± 0.10	0.43 ± 0.05	<0.005
LAVI (mL/m <sup>2</sup> )	38.1 ± 11.4	26.7 ± 4.6	<0.005
E (cm/s)	64 ± 17	66 ± 14	NS
A (cm/s)	82 ± 20	67 ± 13	<0.001
DcT (ms)	243 ± 68	215 ± 64	NS
E / A	0.8 ± 0.3	1.0 ± 0.3	<0.05
IVCT (ms)	46.3 ± 27.7	36.3 ± 21.6	NS
IVRT (ms)	104.7 ± 23.7	88.3 ± 14.7	<0.005
S' velocity (cm/s)	6.9 ± 1.8	8.0 ± 1.5	<0.05
E' velocity (cm/s)	5.6 ± 2.0	7.5 ± 1.9	<0.001
E / E'	12.9 ± 5.1	9.1 ± 2.4	<0.005




A, mitral late diastolic peak velocity; DcT, deceleration time of the E-wave velocity; E, mitral early diastole velocity; E', peak mitral annular velocity during early diastole; IVCT, isovolumic contraction time; IVRT, isovolumic relaxation time; IVS, interventricular septum; LAVI, left atrial volume index; LVDD, left ventricular end-diastolic diameter; LVDs, left ventricular end-systolic diameter; LVEDV, left ventricular end-diastolic volume; LVEF, left ventricular ejection fraction; LVESV, left ventricular end-systolic volume; LVMI, left ventricular mass index; NS, not significant; PW, posterior wall; RWT, relative wall thickness; S', peak mitral annular velocity during systole.

b) End-systolic strain values

The global and regional three-principal strain values are shown in **Figure 2**. Global and regional LSs at the base-, mid-, and apical-LV levels were significantly lower in diabetic patients compared with control subjects. Global LS in control subject was  $220.8 \pm 1.8$ .

These data were used to establish abnormal cut-off value of global LS. This was calculated as the value of the mean  $2 \text{ 2SD}$ . Using the cut-off value of 217.2, 43% (26/60) of the diabetic patients showed abnormal global LS value. CS at the base- and the mid-LV levels did not differ between groups. However, CS at the apical level was significantly lower in diabetic patients, resulting in a significant reduction in global CS. Diabetic patients had also significantly lower regional RS at the basal level and global RS compared with control subjects. Similar results were obtained when diabetic patients with LV hypertrophy on 2D echocardiography were excluded from the analysis. No significant correlation was noted between LVEF and global LS ( $r = 0.205$ ,  $P = \text{NS}$ ) or RS ( $r = 0.24$ ,  $P = \text{NS}$ ). A weak albeit significant negative correlation was noted between LVEF and global CS ( $r = -0.2038$ ,  $P = 0.005$ ).



**Figure 2:** Showing global and regional strain values of diabetic patients and control subjects. Numerical values are represented as mean  $\pm$  SD. In each box plot, upper and lower bars represent 90th and 10th percentiles. Top of the box represents 75th percentile, line in the box median value and bottom of the box means 25th percentile.

Univariate analysis revealed that the reduction of global LS was independently associated with E0 (P, 0.0001), relative wall thickness (P, 0.0001), duration of diabetic disease (P ¼ 0.0006), albuminuria (P ¼ 0.0037), and E-wave velocity (P ¼ 0.0257). No correlation was noted between the reduction of global LS and fasting blood glucose (P ¼ 0.7489) or glycosylated haemoglobin (P ¼ 0.7524). Multivariate linear regression analysis demonstrated that diabetic duration was the only independent predictor for LS reduction (t ¼ 2.22, P ¼ 0.0313). When dividing diabetic patients into two groups according to the duration of disease (.5 and .5 years), global LS was significantly lower in the diabetic group with longer disease duration (216.7+3.0) compared with the short diabetic duration group (218.2+1.9, P , 0.05). Although no significant differences in global RS (46.0+11.7 vs. 44.9+11.7) and CS (223.1+3.9 vs. 222.2+3.5) were noted, RS at the

apical level (38.4+18.7 vs. 27.5+16.0, P , 0.05) and CS at the middle level (223.5+4.2 vs. 221.2+3.6, P , 0.05) were significantly higher in the diabetic group with prolonged disease duration compared with the short duration group.

c) *Post-systolic shortening*

**Figure 3** shows the PSS indices in both groups. The PSI value was significantly larger in diabetic patients compared with control subjects. PSI was significantly larger at the basal level compared with the middle or apical LV levels in both groups. Parametric PSS maps revealed that the distribution did not correlate with the perfusion territory of any coronary artery. PSI values significantly correlated with endsystolic LS in all subjects (n ¼ 85, r ¼ 0.69, P, 0.001) as well as diabetic patients (n ¼ 60, r ¼ 0.64, P, 0.001).




Figure 3: Post-systolic index between the two groups

IV. DISCUSSION

The major findings of this study can be summarized as follows: (i) although global RS, CS, and LS were significantly reduced in diabetic patients compared with age-matched control subjects, the reduction in LS was more prominent and evenly distributed throughout the LV. (ii) The duration of diabetic disease was the only independent predictor for the decrease in LS. (iii) Diabetic patients had more evidence for PSS index, and its distribution did not match the vascular territory of any coronary artery.

a) *Two-dimensional strain*

Identification of early manifestations of diabetic heart disease would allow the institution of timely medical interventions to prevent the development of heart failure. Although diastolic dysfunction has been described as an early stage in diabetic heart disease progression in patients with normal LVEF,[5,6] isolated diastolic dysfunction is usually rare,[7] and when present, it often associated with subclinical systolic dysfunction. Systolic dysfunction might be initially apparent in the longitudinal direction, because subendocardial fibres, which are the ones more

vulnerable to myocardial ischaemia and fibrosis, are longitudinally oriented.[8,10] Several studies have demonstrated that systolic longitudinal dysfunction can be identified using tissue Doppler imaging in patients with hypertension,[18] diabetes,[19,20] and diastolic dysfunction.[21] However, this method provides information in a single direction from a fixed transducer position. This method is also dependent on the angle between the beam and myocardial motion. In contrast, 2DSTE has the advantage that it allows the measurement of all principal LV strains in an angle independent manner, thus eliminating the major limitation of tissue Doppler imaging. Similar to previous tissue Doppler studies,[19,20] we observed that global and regional LSs were significantly reduced in diabetic patients, with 40% of the patients showing abnormal LS values compared with the normal range obtained from our control group of age-matched subjects. In addition, global RS and CS were also reduced in diabetic patients, a finding which is in agreement with a previous magnetic resonance imaging study.[22] On the contrary, Fang *et al*, [11] using tissue Doppler imaging reported that reduced longitudinal function was compensated by the augmentation of radial function in diabetic patients. Differences between study populations and in the method of measuring strain could have accounted for the discrepancies between Fang's and our study. Our results suggest that abnormal function is more widespread than just in the longitudinal direction in diabetic patients. We found that the reduction in global LS was independently associated with diabetic duration as well as early diastolic indices ( $E$ -wave velocity and  $E'$ ), relative wall thickness, and albuminuria. Significant correlation between global LS and  $E'$  confirms the link between systole and diastole, which has been confirmed in previous studies.[23,24] Albuminuria is independently associated with systolic and diastolic dysfunction in diabetic patients.[25,26] The present study showed that diabetic duration was the only independent predictor for the reduction in LS. This highlights the relationship between long-term hyperglycaemia and the impairment of LS. Although global LS was reduced, regional RS and CS were paradoxically increased in diabetic patients of long-term duration. This augmentation in regional RS and CS might reflect a compensation to maintain LVEF in diabetic patients with a long history of disease.

#### b) *Post-systolic shortening*

Myocardial shortening after aortic valve closure, i.e. PSS, has been suggested as a sensitive marker of regional myocardial dysfunction. [16,17] However, PSS may also occur in healthy subjects. To discriminate between pathological and physiological PSS, Voigt *et al*, [17] described that the timing and the magnitude are different between these two situations. The present study showed that PSS was significantly larger in

diabetic patients compared with control subjects. Pathological PSS is usually associated with a reduction in systolic strain. The finding that PSI was negatively correlated with LS in our study supports this concept. Thus, we propose that PSS with reduced LS is a marker of myocardial dysfunction in diabetic patients with preserved LVEF. Interestingly, the distribution of PSS in both diabetic patients and control subjects was mainly observed in the basal myocardium. We also noted that its distribution did not correlate with the vascular territory of the coronary arteries. Although the precise mechanism of why PSS is preferentially observed in the basal myocardium is unknown, PSS observed in this study is not related to myocardial ischaemia induced by epicardial coronary artery stenosis.

#### c) *Study limitations*

The study size was relatively small. Thus, our results cannot be extrapolated to the general diabetic population. The majority of diabetic patients had concomitant hypertension, which also affects longitudinal function. However, exclusion of hypertensive diabetic patients would produce significant bias in our results. Diabetic patients were considered to have a low probability of coronary artery disease based on clinical grounds and normal resting echocardiography.

## V. CONCLUSIONS

LVEF is not a sensitive indicator for the detection of subclinical systolic dysfunction in our study. Diabetic duration was the only independent predictor for the reduction of global LS. 2DSTE has the potential for detecting subclinical LV systolic dysfunction, and it might provide useful information for the risk stratification of an asymptomatic diabetic population.

## REFERENCES RÉFÉRENCES REFERENCIAS

1. Bell DS. Heart failure in the diabetic patient. *Cardiol Clin* 2007; 25: 523–38.
2. Fang ZY, Prins JB, Marwick TH. Diabetic cardiomyopathy: evidence, mechanisms, and therapeutic implications. *Endocr Rev* 2004; 25: 543–67.
3. Fang ZY, Schull-Meade R, Downey M, Prins J, Marwick TH. Determinants of subclinical diabetic heart disease. *Diabetologia* 2005; 48: 394–402.
4. Fang ZY, Yuda S, Anderson V, Short L, Case C, Marwick TH. Echocardiographic detection of early diabetic myocardial disease. *J Am Coll Cardiol* 2003; 41: 611–7.
5. Galderisi M. Diastolic dysfunction and diabetic cardiomyopathy: Evaluation by Doppler echocardiography. *J Am Coll Cardiol* 2006; 48: 1548–51.
6. Poirier P, Bogaty P, Garneau C, Marois L, Dumesnil JG. Diastolic dysfunction in normotensive men with

- well-controlled type 2 diabetes: importance of maneuvers in echocardiographic screening for preclinical diabetic cardiomyopathy. *Diabetes Care* 2001; 24: 5–10.
7. Sanderson JE, Fraser AG. Systolic dysfunction in heart failure with a normal ejection fraction: Echo-Doppler measurements. *Prog Cardiovasc Dis* 2006; 49: 196–206.
  8. Greenbaum RA, Ho SY, Gibson DG, Becker AE, Anderson RH. Left ventricular fibre architecture in man. *Br Heart J* 1981; 45: 248–63.
  9. Henein MY, Gibson DG. Normal long axis function. *Heart* 1999; 81: 111–3.
  10. Henein MY, Gibson DG. Long axis function in disease. *Heart* 1999; 81: 229–31.
  11. Fang ZY, Leano R, Marwick TH. Relationship between longitudinal and radial contractility in subclinical diabetic heart disease. *Clin Sci (Lond)* 2004; 106: 53–60.
  12. Cho GY, Chan J, Leano R, Strudwick M, Marwick TH. Comparison of two dimensional speckle and tissue velocity based strain and validation with harmonic phase magnetic resonance imaging. *Am J Cardiol* 2006; 97: 1661–6.
  13. Langeland S, D’Hooge J, Wouters PF, Leather HA, Claus P, Bijmens B et al. Experimental validation of a new ultrasound method for the simultaneous assessment of radial and longitudinal myocardial deformation independent of insonation angle. *Circulation* 2005; 112: 2157–62.
  14. Reisner SA, Lysyansky P, Agmon Y, Mutlak D, Lessick J, Friedman Z. Global longitudinal strain: A novel index of left ventricular systolic function. *J Am Soc Echocardiogr* 2004; 17: 630–3.
  15. Wang J, Khoury DS, Yue Y, Torre-Amione G, Nagueh SF. Preserved left ventricular twist and circumferential deformation, but depressed longitudinal and radial deformation in patients with diastolic heart failure. *Eur Heart J* 2008; 29: 1283–9.
  16. Voigt JU, Exner B, Schmiedehausen K, Huchzermeyer C, Reulbach U, Nixdorff U et al. Strain-rate imaging during dobutamine stress echocardiography provides objective evidence of inducible ischemia. *Circulation* 2003; 107: 2120–6.
  17. Voigt JU, Lindenmeier G, Exner B, Regenfus M, Werner D, Reulbach U et al. Incidence and characteristics of segmental postsystolic longitudinal shortening in normal, acutely ischemic, and scarred myocardium. *J Am Soc Echocardiogr* 2003; 16: 415–23.
  18. Nishikage T, Nakai H, Lang RM, Takeuchi M. Subclinical left ventricular longitudinal systolic dysfunction in hypertension with no evidence of heart failure. *Circ J* 2008; 72: 189–94.
  19. Andersen NH, Poulsen SH, Eiskjaer H, Poulsen PL, Mogensen CE. Decreased left ventricular longitudinal contraction in normotensive and normoalbuminuric patients with Type II diabetes mellitus: a Doppler tissue tracking and strain rate echocardiography study. *Clin Sci (Lond)* 2003; 105: 59–66.
  20. Ha JW, Lee HC, Kang ES, Ahn CM, Kim JM, Ahn JA et al. Abnormal left ventricular longitudinal functional reserve in patients with diabetes mellitus: Implication for detecting subclinical myocardial dysfunction using exercise tissue Doppler echocardiography. *Heart* 2007; 93: 1571–6.
  21. Yip G, Wang M, Zhang Y, Fung JW, Ho PY, Sanderson JE. Left ventricular long axis function in diastolic heart failure is reduced in both diastole and systole: time for a redefinition? *Heart* 2002; 87: 121–5.
  22. Fonseca CG, Dissanayake AM, Doughty RN, Whalley GA, Gamble GD, Cowan BR et al. Three-dimensional assessment of left ventricular systolic strain in patients with type 2 diabetes mellitus, diastolic dysfunction, and normal ejection fraction. *Am J Cardiol* 2004; 94: 1391–5.
  23. Vinereanu D, Lim PO, Frenneaux MP, Fraser AG. Reduced myocardial velocities of left ventricular long-axis contraction identify both systolic and diastolic heart failure—a comparison with brain natriuretic peptide. *Eur J Heart Fail* 2005; 7: 512–9.
  24. Yip GW, Zhang Y, Tan PY, Wang M, Ho PY, Brodin LA et al. Left ventricular long-axis changes in early diastole and systole: impact of systolic function on diastole. *Clin Sci (Lond)* 2002; 102: 515–22.
  25. Liu JE, Robbins DC, Palmieri V, Bella JN, Roman MJ, Fabsitz R et al. Association of albuminuria with systolic and diastolic left ventricular dysfunction in type 2 diabetes: the Strong Heart Study. *J Am Coll Cardiol* 2003; 41: 2022–8.
  26. Shim CY, Park S, Choi EY, Kang SM, Cha BS, Ha JW et al. Is albuminuria an indicator of myocardial dysfunction in diabetic patients without overt heart disease? A study with Doppler strain and strain rate imaging. *Metabolism* 2008.

