



GLOBAL JOURNAL OF MEDICAL RESEARCH: E
GYNECOLOGY AND OBSTETRICS
Volume 20 Issue 5 Version 1.0 Year 2020
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Prevalence of Acne Associated Gynecological Diseases among Multiethnic Female Medical Students

By Khuraseva A.B & Jayaweera J.A.C.S.

Kursk State Medical University

Abstract- *The purpose of the study:* To investigate and analyze the presence of pathological acne and their correlation between gynecological disorders among young female medical students from different ethnicity.

Material and Methods: The prospective study included 126 female students from Kursk State Medical University aged 18-30 years. This study group consists of six nationalities Indians (27.7%), Nigerians (12.6%), Sri Lankans (15.8%), Malaysians (26.9%), Brazilians (9.5%), and Thai (7.1%) accordingly. The clinical nature of acne, genetic predisposition to acne, gynecological history, lifestyle and habits were recorded in the questionnaire.

Results: After analyzing the questionnaire it was found that 84.7% of students have or had acne in past and 15.3% students never had acne. Among them, 8.26% were diagnosed with PCOS, 65.11% were diagnosed with premenstrual syndrome, 1.8% diagnosis with endometriosis 5.6% with vaginal candidiasis 22.2% diagnosed with hirsutism. 77.5% students with acne got normal menstrual cycle (between 22 to 34) 13% students got oligomenorrhea (menstruation cycle above 34 days) 12.1% with polymenorrhea (menstruation cycle lesser than 22 days).

Keywords: *Acne, hormonal imbalance, PCOS, life style modifications.*

GJMR-E Classification: *NLMC Code: WP 100*



Strictly as per the compliance and regulations of:



Prevalence of Acne Associated Gynecological Diseases among Multiethnic Female Medical Students

Khuraseva A.B ^α & Jayaweera J. A.C.S. ^σ

Abstract- *The purpose of the study:* To investigate and analyze the presence of pathological acne and their correlation between gynecological disorders among young female medical students from different ethnicity.

Material and Methods: The prospective study included 126 female students from Kursk State Medical University aged 18-30 years. This study group consists of six nationalities Indians (27.7%), Nigerians (12.6%), Sri Lankans (15.8%), Malaysians (26.9%), Brazilians (9.5%), and Thai (7.1%) accordingly. The clinical nature of acne, genetic predisposition to acne, gynecological history, lifestyle and habits were recorded in the questionnaire.

Results: After analyzing the questionnaire it was found that 84.7% of students have or had acne in past and 15.3% students never had acne. Among them, 8.26% were diagnosed with PCOS, 65.11% were diagnosed with premenstrual syndrome, 1.8% diagnosis with endometriosis 5.6% with vaginal candidiasis 22.2% diagnosed with hirsutism. 77.5% students with acne got normal menstrual cycle (between 22 to 34) 13% students got oligomenorrhea (menstruation cycle above 34 days) 12.1% with polymenorrhea (menstruation cycle lesser than 22 days)

Conclusion: Many causes of adult acne are due to changes in hormone levels that women experience at certain points during their lives such as before menstrual periods, starting or stopping birth control pills and polycystic ovarian syndrome and their prevalence percentage depend on their nationality.

Keywords: Acne, hormonal imbalance, PCOS, life style modifications.

disorders and also potential skin marker of internal diseases or component of syndromes such as polycystic ovarian syndrome, Hyperandrogenism insulin resistance acanthosis nigricans syndrome (HAIR-AN syndrome) and SAHA syndrome [20].

Women of secondary reproductive age suffer more from acne than men. Psychological stress, diet, smoking, genetic predisposition and hormonal imbalance have been considered as factors that can trigger or worsen acne [15].

I. INTRODUCTION

Acne is the most common type of inflammatory dermatological disease widespread among any age from newborn to menopause [15]. It affects nearly 80% of people at some time between the ages of 11 and 30 years. It can persist for several years and result in disfigurement and permanent scarring, and it can have serious adverse effects on psychosocial development, resulting in emotional problems, withdrawal from society, and depression [28]. Acne is a multifactorial disease which is associated with systemic

Author α: Doctor of Medical Science, Professor Department of Obstetrics and Gynecology Kursk state medical university Kursk, Russian Federation. e-mail: chira.jaya@gmail.com

Author σ: Clinical Resident Kursk state medical university Kursk, Russian Federation.




Figure 1: Pathogenesis of acne and risk factors (Khuraseva AB, Jayaweera JACS. Acne associated gynecological diseases and risk factors in the multiethnic women. *Obstet Gynecol Int J.* 2019; 10(1):42-45. DOI: 10.15406/ogij.2019.10.00411)

II. MATERIAL AND METHOD

This research was conducted in the Department of Obstetrics and Gynecology; Kursk State Medical University on 4th year and 5th year female medical students aged 18 to 30. The number of students enrolled in the study was 126 after application of inclusion and exclusion criteria. The experiment consists of female students from Thailand, Nigeria, Brazil, Malaysia, India and Sri Lanka. They were given a questionnaire about the presence and absence of acne, their location, type of acne, health history, gynaecological history, genetic and life style considerations. All students were thoroughly informed about the study aims and through discussion about the procedure, associated benefits and risks and assigned written consent. The response rate was 95%.

- a) *The inclusion criteria* was history or / and presence of acne, age above 18 below 30 female, non pregnant and non lactating women.
- b) *The exclusion criteria* was absence of acne, age below and above 30, Pregnant and lactating woman

12.6% Nigerian, 26.9% Malaysian, and 15.8% SriLankan students totally. There were 7.1% Thai, 9.5% Brazilian, 17.4% Indian, 12.6% Nigerian 25.4% Malaysian, and 12.7% Sri Lankan students with acne. Whereas 10.3% Indian, 1.6%Malaysian, 3.2% Sri Lankan without acne.

III. RESULTS

After the evaluation of questionnaire following statistical data was obtained. According to the results there were 7.1% Thai, 9.5% Brazilian, 27.7% Indian,

Evaluation methods




Figure 2: Study design

According to the figure 3, 77.5% students with acne got normal menstrual cycle (menstruation cycle between 22 to 34 days), 13% students got oligomenorrhea (menstruation cycle above 34 days), 12.1% with polymenorrhea (menstruation cycle lesser than 22 days). With reference to figure 4, 61.6% of students with acne got 4-6 days of menstruation

duration in one cycle which is considered to be normal. 19.6% students got menorrhagia (7-8 days of menstruation duration in one menstruation cycle). 3.7% students are more prone to hyper menstrual syndrome since more than 8 days of menstruation duration in one cycle. 14.9% students show hypomenorrhea since they have 2-3 days of menstruation duration in one cycle.




Figure 3: Length of menstruation cycle (students with acne)




Figure 4: Duration of menstruation cycle (students with acne)

Table 1: Menstrual history of the participants (students with and without acne)

| | Students with acne | Students without acne |
|------------------------------|--------------------|-----------------------|
| Average age of menarche | 12-13 | 13-14 |
| Regular menstruation cycle | n=75 59.5% | n=13 10.3% |
| Irregular menstruation cycle | n=32 25.3% | n=6 4.7% |

After analyzing the menstruation history of the participants with acne (with reference table 1) 59.5% students with regular menstruation cycle, 25.3% with irregular menstruation. Participants without acne 10.3% with regular menstruation, 4.7% with irregular menstruation. Average age of menarche for students with acne is 12-13 years old whereas 13-14 years students without acne.

When analyzing the gynecological diseases in the participants with acne the below mentioned results were obtained (figure 5). Brazilian participants got highest variety of gynecological diseases, premenstrual syndrome (11.21%), hirsutism (3.7%), polycystic ovarian syndrome (PCOS) (2.8%), vaginal candidiasis (2.8%), and endometriosis (0.9%) compared to other nationalities.

Nigerian and Thai students got the second largest variety of gynecological diseases. Nigerian participants, got premenstrual syndrome (10.2%), hirsutism (5.6%), polycystic ovarian syndrome (PCOS) (1.86%), and vaginal candidiasis (2.8%). Thai participants got premenstrual syndrome (7.4%), hirsutism (0.9%), polycystic ovarian syndrome (PCOS) (0.9%), and endometriosis (0.9%). Whereas Indian, Malaysian, SriLankan students got only premenstrual syndrome, hirsutism and polycystic ovarian syndrome.

Totally from the students with acne, 8.26% were diagnosed with PCOS, 65.11% were diagnosed with premenstrual syndrome, 1.8% diagnosis with endometriosis 5.6% with vaginal candidiasis and 22.2% diagnosed with hirsutism.




Figure 5: Prevalence of gynecological diseases among students with acne

IV. CONCLUSION

Acne is a common skin condition which mostly affects woman of secondary reproductive age. It is not only a dermatological problem but also affects woman in socially and psychosocial aspects. Acne can be the sign of many hormone related gynecological diseases.

This study consisted of 126 female medical students which have understandable knowledge about

dermatology and gynecology which could help us in increasing the success rate of answers in the study questionnaire. According to research analysis prevalence of acne was found in 84.7% students who participated in this research work. From them considerable number of students with acne have more prevalence to gynecological disorders such as polycystic ovarian syndrome, premenstrual syndrome, endometriosis, vaginal candidiasis, hirsutism,

oligomenorrhea, polymenorrhea, menorrhagia and hypomenorrhea. Brazilian students got highest variety of gynecological disorders where as Srilankan, Malaysian, Indian students got least variety of gynecological disorders.

Being medical students they had to lead a stressful life with more unhealthy foods, lack of physical exercises due to busy schedule with their studies. Unhealthy lifestyle of students might leads to obesity, diabetes mellitus, hormonal imbalance and psychological stress issues which can lead to future severe form of gynecological disorders.

As health care providers it is our main responsibility to pre diagnose and screen the hormonal imbalance, endocrine disorders and gynecological disorders. And take measures to alter healthy life style and stress among young female medical students. These measures could lead to healthy population of young female doctors.

LITERATURE REVIEW

- Anamaria Jović, Branka Marinović, Krešimir Kostović, Romana Čević, Aleksandra Basta-Juzbašić, Zrinka Bukvić Mokos. The Impact of Psychological Stress on Acne. University Hospital Centre Zagreb, Department of Dermatology and Venereology, University of Zagreb School of Medicine, Zagreb, Croatia. 2017; 25(2): 133-14.
- Arora M.K., Seth S, Dayal S., "The relationship of lipid profile and menstrual cycle with acne vulgaris." *Clinical Biochemistry*. 2010; 43(18): 1415-1420.
- Bowe, W. P., Joshi, S. S. & Shalita, A. R. Diet and acne. *J Am Acad Dermatol* 63, 124–141 (2010).
- Bowe W, Patel NB, Logan AC. Acne vulgaris, probiotics and the gut-brain-skin axis: from anecdote to translational medicine. *Benef Microbes* 2014; 5: 185-99.
- Carnevale, R. et al. Acute Impact of Tobacco vs Electronic Cigarette Smoking on Oxidative Stress and Vascular Function. 2016;; *Chest* 150, 606–612.
- Chen WC, Zouboulis CC. Hormones and the pilosebaceous unit. *Dermato-Endocrinology*. 2009; 1(2); 81-86.
- Chen W, Obermayer-Pietsch B, Hong J.B., Melnik BC, Yamasaki O, Dessinioti C et al. Acne-associated syndromes: models for better understanding of acne pathogenesis. *J Eur Acad Dermatol Venereol*. 2011; 25(6): 637–46.
- Collier C. N., Harper J. C., Cantrell W. C. The prevalence of acne in adults 20 years and older // *JAm Acad Dermatol*. 2008; 58: 56–59 [PubMed].
- Dalamaga M, Papadavid E, Basios G, Vaggopoulos V, Rigopoulos D, Kassanos D, et al. Ovarian SAHA syndrome is associated with a more insulin-resistant profile and represents an independent risk factor for glucose abnormalities in women with polycystic ovary syndrome: a prospective controlled study. *J Am Acad Dermatol*. 2013; 69(6): 922–30.
- Davidovici, B. B. & Wolf, R. The role of diet in acne: facts and controversies. *Clin Dermatol* 2010. 28, 12–16.
- Dédjan AH, Chadli A, El Aziz S, Farouqi A. Hyperandrogenism-Insulin resistance-acanthosis nigricans syndrome. *Case Rep Endocrinol*. 2015; 193097.
- Escobar-Morreale HF, Carmina E, Dewailly D, Gambineri A, Kelestimur F, Moghetti P, et al. Epidemiology, diagnosis and management of hirsutism: a consensus statement by the androgen excess and polycystic ovary syndrome society. *Hum Reprod Update*. 2012; 18(2):146–70.
- Housman E, Reynolds RV. Polycystic ovary syndrome: a review for dermatologists: Part I. Diagnosis and manifestations. *J Am Acad Dermatol*. 2014; 71(5): 847.
- Kapoor, D. & Jones, T. H. Smoking and hormones in health and endocrine disorders. *Eur J Endocrinol* 152, 491–499 (2005). 62 (2014).
- Khuraseva A.B, Jayaweera J.A.C.S. Acne associated gynecological diseases and risk factors in the multiethnic women. *Obstet Gynecol Int J*. 2019; 10(1): 42-45 DOI: 10.15406/ogij.2019.10.004 11.
- Knowles SR, Nelson EA, Palombo EA. Investigating the role of perceived stress on bacterial flora activity and salivary cortisol secretion: a possible mechanism underlying susceptibility to illness. *Biol Psychol* 2008; 77: 132-7.
- Lee W.J, et al. "Influence of substance-P on cultured sebocytes." *Archives of Dermatological Research*. 2008; 300(6): 311-317.)
- Legro R.S, Arslanian SA, Ehrmann DA, Hoeger KM, Murad MH, Pasquali R. Diagnosis and treatment of polycystic ovary syndrome: an endocrine society clinical practice guideline. *J Clin Endocrinol Metab*. 2013; 98(12): 4565–92.
- Makrantonaki E, Ganceviciene R and Zouboulis CC. *Dermatoendocrinology*. 2011; 3(1): 41-49.
- Nazan Emiroglu. *Acne Associated Syndromes*. 2017; DOI10.5772/65635.
- Norman RJ, Dewailly D, Legro RS, Hickey TE. Polycystic ovary syndrome. *Lancet*. 2007; 370 (9588): 685–97.
- Perkins, A. C., Cheng, C. E., Hillebrand, G. G., Miyamoto, K. & Kimball, A. B. Comparison of the epidemiology of acne vulgaris among Caucasian, Asian, Continental Indian and African American women. *J Eur Acad Dermatol Venereol* 25, 1054–1060 (2011).
- Schmidt TH, Shinkai K. Evidence-based approach to cutaneous hyperandrogenism in women. *J Am Acad Dermatol*. 2015; 73(4): 672–90.

24. Slominski AT, Zmijewski MA, Zbytek B, Tobin DJ, Theoharides TC, Rivier J. Key role of CRF in the skin stress response system. *Endocr Rev* 2013; 34: 827- 84.
25. Slominski A. On the role of the corticotropin-releasing hormone signalling system in the aetiology of inflammatory skin disorders. *Br J Dermatol* 2009; 160: 229-32.
26. Suh DH, Kim BY, Min SU, Lee DH, Yoon MY, Kim NI, et al. A multicenter epidemiological study of acne vulgaris in Korea. *Int J Dermatol* 2011; 50: 673-81.
27. Szabo, K. et al. Interleukin-1A +4845(G> T) polymorphism is a factor predisposing to acne vulgaris. *Tissue Antigens* 76, 411–415 (2010).
28. Wang YY, Li SW, Luo S, et al. How to Evaluate Acne in Reproductive-Age Women: An Epidemiological Study in Chinese Communities. *Biomed Res Int.* 2019; 2019: 6126808. Published 2019 Feb 3. doi:10.1155/2019/6126808
29. Хурасева А.Б.Поиски путей коррекции син-дрома гиперандрогении у девочек, рожден-ных с большой массой тела // Акушерство и гинекология. 2009; (5): 68–71.
30. Хурасева А. Б. Акне: не только косметическая проблема Современная гинекология Том № 2 Медицинский алфавит № 13 (350) 2018: 31-34.

