



GLOBAL JOURNAL OF MEDICAL RESEARCH: K
INTERDISCIPLINARY
Volume 20 Issue 7 Version 1.0 Year 2020
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Results of Salt Cognition Test using SALSAVE since 67 Female University Students

By Naomi Katayama, Akemi Ito & Mayumi Hirabayashi

Nagoya Women's University

Abstract- In Japan, salt reduction is encouraged to prevent high blood pressure. However, it is difficult to reduce salt, and good salt intake per day did not fail to the target value. In this study, we report a saltiness cognitive threshold test using female university students. The participant is 67 female university students. Participants were subjected to a salty cognitive threshold test using SALSAVE (manufactured by Advantech). The saltiness test started from a light taste and tried a strong taste in order. The salt concentration is 0.6%, 0.8%, 1.0%, 1.2%, 1.4%, 1.6%. We also conducted a questionnaire survey on eating habits. As a result, 62 out of 67 female university students who felt salty at a concentration of 0.6% were 92% of all participants. However, two female university students did not feel taste even with a salt concentration of 1.6%. They are 3% of all participants. As a result of the questionnaire survey, female university students answered that they had a good taste and secreted saliva well, and they usually had a rather light diet. In the future, we would like to increase the number of participants and compare more detailed dietary habits with SALSAVE results.

Keywords: saltiness test, cognition, threshold, salsave, university student.

GJMR-K Classification: NLMC Code: QU 84



Strictly as per the compliance and regulations of:



Results of Salt Cognition Test using SALSAVE since 67 Female University Students

Naomi Katayama ^α, Akemi Ito ^σ & Mayumi Hirabayashi ^ρ

Abstract- In Japan, salt reduction is encouraged to prevent high blood pressure. However, it is difficult to reduce salt, and good salt intake per day did not fail to the target value. In this study, we report a saltiness cognitive threshold test using female university students. The participant is 67 female university students. Participants were subjected to a salty cognitive threshold test using SALSAVE (manufactured by Advantech). The saltiness test started from a light taste and tried a strong taste in order. The salt concentration is 0.6%, 0.8%, 1.0%, 1.2%, 1.4%, 1.6%. We also conducted a questionnaire survey on eating habits. As a result, 62 out of 67 female university students who felt salty at a concentration of 0.6% were 92% of all participants. However, two female university students did not feel taste even with a salt concentration of 1.6%. They are 3% of all participants. As a result of the questionnaire survey, female university students answered that they had a good taste and secreted saliva well, and they usually had a rather light diet. In the future, we would like to increase the number of participants and compare more detailed dietary habits with SALSAVE results.

Keywords: saltiness test, cognition, threshold, salsave, university student.

I. INTRODUCTION

Japanese Dietary Intake Standards in the 2020 version, the daily intake of salt for females is recommended to be 6.5g or less. Salt intake is

Table 1: Average \pm Standard deviation (SD) of participant's age and body composition (n=67)

Participants	Age	Height	Weight
Average	20.6	158.9	50.3
S D	0.6	5.7	4.5

b) Assessment of salt taste identification

Participants were subjected to a salty cognitive threshold test using SALSAVE (manufactured by Advantech). The saltiness test started from a light taste and tried a strong taste in order. The saltiness test starts from 0.6%, and the concentration increases by 0.2% in 6 steps up to 1.6%. Participants put a filter paper impregnated with salt in their mouth to check the taste,

decreasing with each revision of dietary standards. In the near future, it is expected that salt intake in Japanese dietary intake standards will lower to the international standard of 6.0g. By reducing salt, it is expected to prevent illness from various diseases (high blood pressure, kidney disease, heart disease, etc.). Therefore, the purpose of this study was to conduct a salt concentration cognition test on Japanese people to understand the actual condition of the cognitive threshold for salt, and to use it as future data. To begin with, we report that a healthy female university student underwent a salt concentration recognition test.

II. MATERIALS AND METHODS

a) Participants

The participant is 67 female university students. Their average age \pm standard deviation (SD) was 29.6 ± 0.6 , average height \pm SD was 158.9 ± 5.7 cm, and average weight \pm SD was 50.3 ± 4.5 kg (Table 1).

and then answered to the inspector what the taste was. The inspector recorded the answers of the participants.

We also conducted a questionnaire survey on dietary habits. There are four questions, 1) Does saliva come out? 2) Do you feel the taste? 3) Frequency of purchase of restaurants and commercial food, 4) Regular seasoning (for food was salty or thin) (Table 2).

Author α σ ρ : Graduate School of Nagoya Women's University, Nagoya City, Japan.

Corresponding Author α : Nagoya Women's University, Nagoya City, Japan. e-mail: naomik@nagoya-wu.ac.jp

Table 2: Questionnaire survey items

	Question 1	Question 2	Question 3	Question 4
	Saliva secretion	Taste perception	Use of restaurants and commercial food	Favorite food salt taste
1	Very well	Very well	every day	Strong salt taste
2	Well	Well	four or five times a week	rather strong salt taste
3	Not good	Not good	two or three times a week	rather light salt taste
4	Do not know		once a week	light salt taste
5			two or three times a month	
6			Hardly used	

c) *Ethical review board*

This study conducted with the approval of the Ethics Committee (Nagoya women's university 'hitowo mochiita kennkyuuni kansuru iinnkai'). The approval number is 30-14.

SALSAVE. As a result, 62 out of 67 female university students who felt salty at a concentration of 0.6% were 92% of all participants. And, three female university students felt a salt concentration of 0.8%, were 4.5% of all participants. However, two female university students did not feel taste even with a salt concentration of 1.6%. This is 3% of all participants (Table3).

III. RESULTS

a) *Saltiness recognition test result*

Sixty-seven female university students underwent a saltiness cognitive threshold test using

Table 3: Female University students Saltiness cognitive threshold test results (n=67)

	0.60%	0.80%	1.00%	1.20%	1.40%	1.60%	1.6%以上
Participants (number of students)	62	3	0	0	0	0	2
Participants (%)	92.5	4.5	0.0	0.0	0.0	0.0	3.0

b) *Questionnaire results*

Table 4 shows the results of the questionnaire survey. When asked if they had enough saliva, most female university students replied that they had enough or normal secretions. None of the female university students answered that they had none or poorsaliva. In response to the question of feeling the taste, most female university students answered that they could well or understand the taste. None of the female university

students answered that they didn't understand the question of whether they would feel the taste. When asked about the frequency of use of restaurants and groceries, female university students answered that they would use 2-3 times a week or once a week. When asked if the usual seasoning for food was salty or not too much salty, female university students answered that they were the lightly salted.

Table 4: Results of Questionnaire survey conducted on female university students (n=67)

	Saliva secretion	Taste perception	Use of restaurants and commercial food	Favorite food salt taste
Average	1.3	1.5	3.8	2.3
S D	0.7	0.5	1.0	0.7

IV. DISCUSSION

Many researchers have reported the relationship between salt intake and blood pressure ¹⁾ and the relationship between hypertension ^{2,3,4)}. There are also reports on diet and salt intake in the younger generation ⁵⁾. We also reported the results of the saltiness cognition threshold test conducted by Yakumo Study on the elderly ^{6,7)}. There are also reports on the relationship between salt intake and blood pressure overseas ^{8,9)}, as well as educational effects on salt intake ^{10,11)}. Currently, many new drugs for lowering blood pressure have developed for people with high blood pressure ¹²⁾, but dietary food habits also desirable can be improved their blood pressure. The drug also

has side effects and that costs a lot to keep taking it. In Alzheimer's dementia, the taste is not known, and the value of the salty cognitive threshold test also deteriorates ¹³⁾. Therefore, it is highly likely that the dietary intake will be high, which may raise blood pressure. A low saltiness recognition threshold means that the saltiness of the meal can reduced. Keeping the salty cognitive threshold low makes sense for a healthy diet. We conducted a saltiness cognition threshold test on 67 young female university students. As a result, they recognized 0.6% saltiness, and with included 0.8% recognized students, the total number of recognized was 96.5%.The result of the dietary habit questionnaire showed that saliva was well secreted, the taste of the meal was well understood, and the dietary intake was

light. However, two female university students did not feel taste even with a salt concentration of 1.6%. This is 3% of all participants.

In the future, we would like to conduct a more detailed questionnaire survey on dietary intake and compare it with the salty cognitive threshold test results.

V. CONCLUSIONS

The results of this test show that most female university students (92%) feel salty at 0.6%. Also, female university students were eating a light taste in their daily life. Moreover, the use of restaurants and the use of commercially available foods were once or 2-3 times a week. The effect of the salt reduction awareness campaign in Japan, which has continued for over 20 years, is good for the younger generation. However, on the other hand, among the female university students who participated in this study, two students could not recognize 1.6% saltiness, so it is necessary to investigate the dietary habits of the participants in more detail.

ACKNOWLEDGEMENTS

This study was supported by the research aid of Choju-iryō-kenkyū-kaihatsuhi 30-14 and the Japanese Society of Taste Technology, 2019.

REFERENCES RÉFÉRENCES REFERENCIAS

1. Volos BE, Vasil'ev lu M, Masliaeva LV and Snegurskaia IA (1994). Taste receptor response to sodium chloride and natriuresis in the workers and employees of an industrial enterprise. *Likars'ka sprava* 3-4.
2. Olayemi SO and Mabadeja AF (2003). Comparative study of salt taste threshold of hypertensives, their normotensive relatives and non-relatives. *The Nigerian postgraduate medical journal* 10 (2).
3. Elias SO, Azinge EC, Umoren GA, Jaja ST and Sofola OA (2011). Salt-sensitivity in normotensive and hypertensive Nigerians. *Nigerian quarterly journal of hospital medicine* 21 (1).
4. Volkov VS, Poseliugina OB, Nilova SA, Vinogradova TS, Rokkina SA and Svistunov OP (2010). Impaired gustatory sensitivity of the tongue to table salt as a risk factor of arterial hypertension. *Klinicheskaia meditsina* 88(1).
5. Malaga S, Diaz JJ, Arguelles J, Perillan C, Malaga I and Vijande M (2003). *Pediatric nephrology* 18(5).
6. Katayama N, Kondo S, Ootake H et al (2018). Odour and Salt Taste Identification in Older Adults: Evidence from the Yakumo Study in August, 2018. *Acade. J. Med. Plants* 7(3) 066-071.
7. Nishimoto K, Ohhori J, Shimomugi T, Kurono Y (2005). Reproducibility of taste examination with Salsave: Control study for healthy volunteers. *Japan Society of Stomato-pharyngology*. pp.309-315.
8. Cho H, Kinm SM, Jeong SS and Kim SB (2016). Comparison of salt taste thresholds and salt usage behaviours between adults in Myanmar and Korea. *Asia Pacific journal of clinical nutrition* 25 (4).
9. Okoro EO, Uroghide GE and Jolayemi ET (1998). Salt taste sensitivity and blood pressure in adolescent school children in southern Nigeria. *East African medical journal* 75 (4).
10. Kusabe U, Mori Y, Okagaki M, Neriya H, Adachi T, Sugishita C, Aonomura K, Kimura T, Kishimoto N, Nakagawa H, Okigaki M, Hatta T and Matsubara H (2009). Sodium restriction improves the gustatory threshold for salty taste in patients with chronic kidney disease. *Kidney international* 76 (6).
11. Ferrante D, Apro N, Ferreira V, Virgolini M, Aguilar V, Sosa M, Perel P and Casas J (2011). Feasibility of salt reduction in processed foods in Argentina. *Pan American journal of public health* 29 (2).
12. Petrova TS, Bazhenov ND, Mazur VV and Mazur ES (2012). Gustatory sensitivity threshold to table salt and efficacy of the treatment of newly diagnosed patients with antihypertensive therapy. *Klinicheskaia meditsina* 90 (4).
13. Contri-Degiovanni PV, Degiovanni GC, Ferriolli E, Da Costa LNK and Moriguti JC (2020). Aging clinical and experimental research 2.