



GLOBAL JOURNAL OF MEDICAL RESEARCH: F DISEASES

Volume 20 Issue 11 Version 1.0 Year 2020

Type: Double Blind Peer Reviewed International Research Journal

Publisher: Global Journals

Online ISSN: 2249-4618 & Print ISSN: 0975-5888

COVID-19: Public Health Implications among Healthcare Workers in a Tertiary Health Facility in Enugu State, Nigeria

By Nwoga Hope O, Ajuba Miriam O & Ezeoke Uche E

Enugu State University

Abstract- *Background:* Coronavirus is one of the emerging respiratory viruses that are known to cause diseases in humans. It causes a range of illnesses ranging from the common cold to Severe Acute Respiratory Syndrome (SARS).

Objectives: To assess the knowledge, attitude, and practice of healthcare workers in a tertiary health facility towards COVID 19 pandemic and also the perceived barrier to the control of the infection in a healthcare setting.

Methodology: The study was a descriptive cross-sectional study that involved all the healthcare workers (HCWs); (doctors, nurses, laboratory scientists) working in a tertiary health facility. Data from the doctors and laboratory scientists were gathered with an online questionnaire formulated in Google form.

Keywords: COVID-19, healthcare workers, Enugu State, Nigeria, tertiary health facility.

GJMR-F Classification: NLNC Code: W 84



COVID-19 PUBLIC HEALTH IMPLICATIONS AMONG HEALTHCARE WORKERS IN A TERTIARY HEALTH FACILITY IN ENUGU STATE NIGERIA

Strictly as per the compliance and regulations of:



COVID-19: Public Health Implications among Healthcare Workers in a Tertiary Health Facility in Enugu State, Nigeria

Nwoga Hope O ^a, Ajuba Miriam O ^a & Ezeoke Uche E. ^b

Abstract- Background: Coronavirus is one of the emerging respiratory viruses that are known to cause diseases in humans. It causes a range of illnesses ranging from the common cold to Severe Acute Respiratory Syndrome (SARS).

Objectives: To assess the knowledge, attitude, and practice of healthcare workers in a tertiary health facility towards COVID 19 pandemic and also the perceived barrier to the control of the infection in a healthcare setting

Methodology: The study was a descriptive cross-sectional study that involved all the healthcare workers (HCWs); (doctors, nurses, laboratory scientists) working in a tertiary health facility. Data from the doctors and laboratory scientists were gathered with an online questionnaire formulated in Google form. It was shared through their different online platforms (Whatsapp). Data from the nurses were collected using a semi-structured questionnaire given to all the nurses through the head nursing services. All the data were analyzed using SPSS version 25. Categorical variables were presented in the form of frequencies and percentages.

Results: Most of the HCWs were within 30-39 years of age (48.1%) and had worked for 11-20 years (45.5%). Majority were females (88.9%), married (86.0%) and nurses (82.1%). About 59.1% of the HCWs had good knowledge of COVID 19, while only 43.4% had a good attitude towards COVID 19. However, the majority (93.2%) of them had good practice towards COVID 19. The commonest perceived barrier to the control of infection was overcrowding in clinics, wards, and emergency rooms.

Conclusion: The knowledge and attitude of the HCWs in this study to COVID 19 was poor, but this did not affect their practice. There should be an orientation of the HCWs for attitudinal change towards the infection.

Keywords: COVID-19, healthcare workers, Enugu State, Nigeria, tertiary health facility.

I. INTRODUCTION

Coronavirus is one of the emerging respiratory viruses that are known to cause diseases in humans.¹ It is a zoonotic infection that can be transmitted from animal-to-human and then human-to-

Author a: Department of community medicine, Enugu State University Teaching Hospital Parklane Enugu, Nigeria.

e-mail: nwogahope@gmail.com

Author a: Department of Community Medicine, Enugu State University College of Medicine Enugu, Nigeria.

Author b: Department of Community Medicine, University of Nigeria College of Medicine Enugu Campus, Enugu, Nigeria.

human.² The most recent outbreak of coronavirus disease 2019 (COVID-19) in Wuhan City, Hubei Province, China, has emerged as a global outbreak and significant public health issue.³

Coronaviruses outbreaks have been recorded previously such as Severe Acute Respiratory Syndrome-Coronavirus (SARS CoV) and Middle East Respiratory Syndrome-Coronavirus (MERS-CoV) in 2003 and 2015, which are similar to the present novel coronavirus SARS CoV 2 (COVID-19). COVID-19 was first reported in December 2019 among patients with viral pneumonia symptoms in Wuhan City, Hubei Province; China.^{2,4} Due to the rapid spread of the virus from the source to many other countries of the world, the World Health Organization (WHO) declared it a Public Health Emergency of International Concern (PHEIC) on January 30, 2020. The first case of COVID-19 in Nigeria was recorded on 27 February 2020, and as of the 29th of September 2020, there were about 58,647 confirmed cases with 1,111 confirmed deaths. Globally there were about 33, 249, 563 cases with 1,000,040 deaths as of 29th September 2020.⁵

However, on the 11th of March 2020 the WHO declared COVID-19 a global pandemic due to its fast spread, increase in the number of countries affected, the severity of illness, and the continuous escalation in the number of cases and causalities.⁶

Coronavirus spreads mainly from person to person through close contact with infected person through respiratory droplets (coughs or sneezes) or by touching a surface or object with the virus on.⁷ The WHO reported that more than 80% of COVID-19 patients showed mild symptoms and recovered without any medical intervention, approximately 20% of infected cases had a severe illness such as shortness of breath, septic shock, and multi-organ failure while an estimated 2% of the cases can be fatal.⁶ The elderly, especially those with underlying chronic diseases are at an increased risk of having severe disease.

Coronavirus is most contagious during the first three days after onset of symptoms, however, spread may be possible before symptoms appear and in later stages of the disease.⁸ It can survive on surfaces for up to 72 hours.⁸ The best prevention for COVID-19 is to avoid being exposed to it. The recommended measures to achieve this include frequent hand washing,



maintaining physical distance from others (especially from those with symptoms), covering coughs and sneezes with a tissue or inner elbow, keeping unwashed hands away from the face using face masks, isolating confirmed and suspected cases.⁹

Healthcare Workers (HCWs) are primarily involved in caring for patients with this highly contagious virus (COVID-19), and it has posed a serious occupational health risks to the HCWs owing to their frequent exposure to infected individuals in the course of their duty.¹⁰ Protection of HCWs and prevention of intra-hospital transmission of infection are important aspects in epidemic response, and this requires that HCWs must have updated knowledge regarding the source, transmission, symptoms, and preventive measures.¹¹ Literature suggests that lack of knowledge and misunderstandings among HCWs leads to delayed diagnosis, the spread of disease, and poor infection control practice.¹² Thousands of HCWs have already been infected. Knowledge, attitude, and practice survey like the present study provide a suitable format to evaluate existing programs and protocols and to identify effective strategies for behavioral change both in the hospital setting and in the society at large.¹³

II. METHODOLOGY

The study was a descriptive cross-sectional design conducted at Enugu State University of Science and Technology, Enugu, Nigeria. It is one of the tertiary health institutions in Enugu State. It is located at the heart of Enugu town.

a) Study population

All the consenting HCWs (doctors, nurses, and laboratory scientists) in the hospital were used for the study. There were about 426 HCWs (doctors, nurses, and laboratory scientists) in the hospital as at the time of data collection and, they were reached for the study.

b) Data collection

The data for the doctors and laboratory scientists were collected using an online questionnaire formulated with Google form. This was shared through the different doctors' and laboratory scientists' online platform (WhatsApp). The data for the nurses was collected using the same semi-structured self -

administered questionnaire shared through the head nursing services to all the nurses. This was so because there was no online platform where all the nurses in the hospital belong.

The questionnaire was formulated after extensive literature review and, according to WHO recommendations on modes of transmission and prevention of COVID-19.⁷ The questionnaire had five sections. The first section was on socio-demographic characteristics of the HCWs, the second section was on knowledge, the third was on attitude, the fourth on practice, and the fifth on perceived barrier to control of COVID-19 in healthcare settings. Eleven questions were used to access the HCWs knowledge of COVID-19. A correct answer scores one while a wrong answer scores 0. The higher the score the, more knowledgeable the HCWS is. Twenty-one questions were used to assess the attitude of the HCWs. A correct answer scores one while an incorrect answer scored 0. Some of the questions were reversed to eliminate the bias of giving a single similar response in all the questions. Thirteen questions were used to access the practice score using 3 Likert-scale questions. The responses were never, sometimes, and always weighing 1, 2, and ,3 respectively.

The perceived barrier to control of COVID-19 was accessed using nine questions. A correct answer scores one while a wrong answer scores 0. The higher the score the, more the barrier to control of COVID-19 in hospital settings

c) Data analysis

All the questionnaires were examined for completeness and all the completed ones were entered into SPSS version 25. All the responses in Google form were also entered into SPSS version 25. The data were edited for errors by generating frequencies. The categorical variables were summarized using frequencies and percentages. The significance level was placed at ≤ 0.05 . The knowledge, attitude, and practice scores were categorized into good and poor. Blooms cut-off criteria¹⁴ of 80% were used to indicate good knowledge (≥ 8.8), attitude (≥ 16.8) and practice (≥ 10.4) towards COVID-19, while less than these cut-offs were categorized as poor.

III. RESULTS

Table 1: Socio-demographic characteristics of the healthcare workers

Variable	Frequency	Percentage
Age		
20-29	13	5.5
30-39	113	48.1
40-49	79	33.6
50-59	28	11.9
Above 59	2	0.9
Gender		

Male	26	11.1
Female	209	88.9
Marital status		
Single	33	14.0
Married	202	86.0
Ethnicity		
Igbo	231	98.3
Yoruba	3	1.3
Edo	1	0.4
Religion		
Christianity	231	98.3
Islam	3	1.3
Traditional Religion	1	0.4
Occupation		
Doctor	31	13.2
Nurse	193	82.1
Laboratory Scientist	11	4.7
Years of service		
1-10	93	39.6
11-20	107	45.5
21-30	24	10.2
Above 30	11	4.7

Out of the 426 HCWs that were reached out for the study, only 235 responded, giving a response rate of 55.2%. Table 1 above shows the socio-demographic characteristics of the studied HCWs. Most of the HCWs were within the 30-39 years age group (48.1%) and have

worked for 11-20 years (45.5%). Majority were females (88.9%), married (86.0%) and nurses (82.1%). Almost all were of the Igbo ethnic group (98.3%) and Christians (98.3%).

Table 2: Knowledge of COVID-19 among healthcare workers

Variable	Yes	
	Frequency	Percentage
Have heard of COVID-19	235	100
Sources of information*		
Government	191	81.3
Internet	213	90.6
Mass media	219	93.2
Family/friends	178	75.7
Social media	200	85.1
Colleagues	202	86.0
Hospital management	170	72.3
Church	3	1.3
COVID-19 is a viral infection	213	90.6
Its incubation period is 14days	212	90.2
COVID-19 can be spread by*		
Droplets from infected persons	234	99.6
Surfaces touched by infected persons	221	94.0
Touching banknotes	167	71.1
Asymptomatic persons	201	85.5
Shaking of hands	220	93.6
Hugging	176	74.9
Staying together in a crowded room	224	95.3
Goods imported from China	65	27.7
Common symptoms of COVID-19 includes*		
Fever	234	99.4
Dry cough	229	97.4
Body aches	177	75.3
Loss of taste	53	22.6
Difficulty in breathing	230	97.9

Vomiting	110	46.8
Sneezing	203	86.4
Diarrhoea	130	55.3
Vaccine for COVID-19 is available	7	3.0
Effective treatment for COVID-19 is available	55	23.4
People with co-morbidities like hypertension/DM are more at risk	205	87.2
Children have no risk of death from COVID-19	33	14.0
COVID-19 may be more dangerous for the elderly	226	96.2
COVID-19 spreads through close contact	219	93.2
All COVID-19 patients develop severe acute respiratory illness	181	77.0
Influenza vaccine also gives protection against COVID-19	20	8.5
COVID-19 can be prevented by*		
Proper and regular hand washing with soap	235	100
Use of hand sanitizers	229	97.4
Social distancing	231	98.3
Wearing face mask	231	98.3
Eating garlic, ginger and local herbs	93	39.6
Avoid touching of the eyes, nose and mouth	228	97.0
Good cough etiquette	225	95.7
Drinking/inhaling hot water	114	48.5
Drinking alcohol	16	6.8
Taking chloroquin tablets	62	26.4
Taking antibiotics	60	25.5
Knowledge categorized		
Poor	96	40.9
Good	139	59.1

*multiple answers allowed

Table 2 above shows the knowledge of the HCWs towards COVID-19. All the HCWs have heard about COVID-19, and their major source of information was mass media (93.2%) and the internet (90.6%). About 90.2% of the HCWs knew the incubation period of the virus and, the majority also knew the modes of spread. However, about 27.7% believe that the virus can be spread by goods imported from China. The majority knew the common symptoms, but only 22.6% knew that loss of taste could be a symptom of COVID-19. About 77.0% asserted that all COVID-19 patients develop a

severe acute respiratory illness. On the knowledge of the preventive measures, all the HCWs knew that proper and regular hand washing with soap could prevent COVID-19. The majority also knew the other preventive measures like the use of hand sanitizers, social distancing, wearing of face mask, and good cough etiquette. However, 39.6% believe that eating garlic, ginger, and other local herbs can prevent COVID-19.

On the categorization of knowledge scores, only 59.1% of the HCWs had good knowledge of COVID-19.

Table 3: Attitude of the healthcare workers towards COVID-19

Variable	Yes	
	Frequency	Percentage
You think you can contact COVID-19	82	34.9
Worried that a family member may contact COVID-19	133	56.6
COVID-19 infection is associated with stigma	195	83.0
Media coverage is exaggerated	93	39.6
The virus was initially designed as a biological weapon	119	50.6
It is a plague caused by sin and unbelief	44	18.7
Designed to control population	47	20.0
Designed by pharmaceutical companies to sell drugs	21	8.9
COVID-19 can be successfully controlled	179	76.2
Self-protection is necessary for the protection of others	225	95.7
Lockdown is an effective control measure	183	77.9
Nigeria can win the battle against COVID-19	190	80.9
If I contact a person infected with COVID-19 I will inform NCDC	216	91.9
If I have symptoms of COVID-19 I will inform NCDC	208	88.5
If I have contact with an infected person I agree to be isolated at home	210	89.4

If I have contact with an infected person I agree to be isolated at an isolation centre	177	75.3
If vaccine for COVID-19 is available, I will be willing to take it	144	61.3
I follow updates on COVID-19 in my country	212	90.2
I follow updates about COVID-19 worldwide	196	83.4
Willing to attend any lecture about COVID-19 around me	212	90.2
COVID-19 can be controlled by following standard precautions for infection control	226	96.2
Attitude categorized		
Poor	133	56.6
Good	102	43.4

Table 3 above shows the attitude of the HCWs towards COVID-19. Only a minority of the HCWs think they can contact COVID-19 (34.9%), while about half (56.6%) were worried that a family member could contact it. The majority agree that self-protection is necessary for protecting others (95.7%) and that lockdown is an effective control measure (77.9%). Only

about 61.3% of the HCWs are willing to take a vaccine for COVID-19. Almost all of the respondents (96.2%) agree that COVID-19 can be controlled by following standard precautions for infection control. Only 43.4% of the HCWs had a good attitude toward COVID-19 on categorization.

Table 4: Healthcare workers practice towards COVID-19

Variable	Yes	
	Frequency	Percentage
I have been to crowded places in recent time	139	59.1
Have cancelled attending events due to COVID-19	174	74.0
Wear face mask always while leaving home	227	96.6
Wear mask in correct way	231	98.3
Dispose mask when it becomes moist or after 8 hours	194	82.6
Follow WHO hand washing technique	218	92.8
Practice social distancing	220	93.6
Follow lockdown regulations	208	88.5
Cover mouth with tissue while sneezing/coughing	199	84.7
Dispose used tissue in covered trash	215	91.5
Avoid touching eyes, mouth and nose as much as possible	204	86.8
I educate my patients about COVID-19	226	96.2
I follow standard precautions for infection control always	224	95.3
Practice categorized		
Poor	16	6.8
Good	219	93.2

Table 4 shows the HCWs practice towards COVID-19. The practice scores of the HCWs were high, as 93.2% of them had good practice towards COVID-19. The majority of them follow the WHO hand washing

technique (92.8%), follow lockdown regulations (88.5%), and follow standard precautions for infection control always (95.3%).

Table 5: Perceived barriers to effective control of COVID-19 in hospital settings

Variable	Yes	
	Frequency	Percentage
Lack of knowledge about mode of transmission of COVID-19	141	60.0
Unavailability of infection control supplies*		
Gloves	179	76.2
Face mask	182	77.4
Face shield	207	88.1
Splash proof apron	214	91.1
Goggles	211	89.8
Carelessness on the part of the healthcare workers	138	58.7
Insufficient training on infection control	190	80.9
Lack of policies on infection control	175	74.5
Less commitment of healthcare workers to policies	139	59.1

Overcrowding in the emergency rooms	206	87.7
Overcrowding in clinics and wards	207	88.1
Not washing hands before and after contact with patients	174	74.0

*multiple answers allowed

Table 5 shows the perceived barrier to control of COVID-19 in a hospital settings. The major perceived barrier to effective control of COVID-19 in the hospital setting was the unavailability of PPEs like splash proof aprons (91.1%) and goggles (89.8%) followed by overcrowding in emergency rooms (87.7%), clinics, and wards (88.1%). Others were insufficient training on infection control (80.9%) and a lack of policies on infection control (74.5%).

IV. DISCUSSION

COVID-19 is a global health problem that is affecting every aspect of human life. HCWs being at the forefront of the fight against the infection, are faced with a high likelihood of being infected with the infection than the general population.^{15, 16}

This is one of the reasons why the WHO have recommended the adoption and use of proper prevention and control measures for HCWs. This includes the use of proper Personal Protective Equipment (PPE), such as face masks, splash-proof gowns, gloves, eye goggles, and FFP2 respirator masks in aerosol-generating procedures.¹⁷

In this study, about 59.1% of the HCWs had good knowledge of COVID-19 including its incubation period, mode of transmission, common symptoms, and preventive measures. This knowledge score was not optimal as knowledge helps to form one's attitude, promote productive behaviors and affect their coping mechanisms towards specific events.¹⁸ The score was, however, similar to the report of a similar study in Uganda where 69% of the HCWs had good knowledge of COVID-19¹⁹ but lower than the report of other studies in China.^{20, 21} Better knowledge among these HCWs may be due to better training on COVID-19.

Further education, training, and retraining of HCWs through continuous professional education, particularly on the common symptoms, modes of transmission, and prevention will help in improving the knowledge of the HCW about COVID-19. The commonest source of information on COVID-19 was the mass media like television, radio, newspapers (93.2%), followed by the internet (90.6%). This indicates that such media should be used to disseminate information about COVID-19 to reach the majority of the HCWs.

The delivery of rapid, reliable information that addresses critical infection control measures was of key importance, e.g., poor hospital infection prevention and control measures, inadequate training in infection control processes, poor compliance with the use of personal protection equipment (PPE), exposure to high-risk procedures such as tracheal intubation and

exposure to unsuspected SARS patients are known to be associated with a high risk of transmission of viral infections to healthcare workers.^{22, 23} The aim of information dissemination is to transfer knowledge of international best practices to the front-liners who need it most, at a rate equal to or faster than the increasing epidemic. The use of social media have shown to be a faster means as concurred by another literature.²⁴

However, there has been worries about misinformation regarding COVID-19 on the internet, which can misguide HCWs because the internet is not without disadvantages. It is, therefore, necessary that before use, healthcare workers must critically appraise the information contained.²⁵ In particular, health authorities and scientists have warned that widespread misinformation about COVID-19 is a serious concern causing xenophobia worldwide.^{26, 27} In this regard, HCWs should carefully evaluate COVID-19-related information and should use scientific and authentic content as information sources.

The only factor that significantly affected the knowledge of COVID-19 in the present study was occupation where the nurses (64.2%) and the laboratory scientists (63.6%) were found to have better knowledge scores than the doctors (25.8%). This was surprising as doctors, by their training, were supposed to have a better knowledge of COVID-19 compared to other HCWs. However, a similar study reported that doctors had better knowledge than the other HCWs.^{21, 28}

One notable point on the HCWs knowledge of the mode of spread of COVID-19 is that about 27.7% asserted that COVID-19 could be contacted from goods imported from China. This can affect the use of aids and other products from China for the prevention of COVID-19 and have a negative impact on the general control measures. Also, 39.6% agree that eating ginger, garlic, and local herbs can prevent COVID-19 infection. This misconception can delay seeking medical advice and result in fatal outcomes for COVID-19 cases. Awareness of the mode of spread and prevention should be heightened among the HCWs as the general public also depend on the information given by the HCWs.

In the present study, only 43.4% of the HCWs had good attitude toward COVID-19. This poor attitude of HCWs towards COVID-19 was reported in a similar studies in Uganda, where only 21% of them had a good attitude toward COVID-19.¹⁹ The similarity may be because both studies were done in Africa with similar beliefs and attitudes.

About half (50.6%) of the HCWs believe that the virus was initially designed as a biological weapon. This finding was similar to the report of a Nigerian study

where 46.9% of the respondents had a similar view.²⁹ About 56.6% are worried that their family member may contact COVID-19. This was lower than the report of a similar study where 79.8% of the HCWs are worried that their family member may contact COVID-19.²⁰

However, there was a good attitude towards some of the individual elements of attitude. About 95.7% believe that self-protection is necessary for the protection of others, 90.2% follow update about COVID-19 in Nigeria, 90.2% are willing to attend lectures on COVID-19 if organized around them, and 96.2% believe that COVID-19 can be controlled by following standard precautions for infection control. A similar study in Egypt reported a similar finding where 95.6% of the studied HCWs asserted that following standard precautions can prevent infection transmission.³⁰

The majority of the HCWs (93.2%) in the present study had good preventive practices towards COVID-19. They reported to follow standard precautions for infection control and follow the WHO hand washing technique. Good knowledge and practice of standard precautions including good cough etiquette, have been shown to reduce infection transmission in hospitals.³¹ The finding in this study was, however, higher than what was reported in another study in Uganda.¹⁹

The commonest perceived barriers to effective infection control were overcrowding in the clinics, wards, and emergency rooms; thus, the HCWs were not able to maintain a safe distance from their patients. This will lead to an increase in infection transmission. An Egyptian study reported that crowdedness and poor ventilation were factors that make HCWs more susceptible to infection than the general population.³⁰ An Iranian study reported that physicians (67%), nurses (24%), and operating room staff (100%) did not keep a one-meter distance from patients due to lack of space.³² Other barriers were the unavailability of infection control supplies, especially the Personal Protective Equipment's (PPEs). This was reported in similar studies in Iran³² and China³³ where lack of these PPEs was reported to be one of the reasons for infection of HCWs with COVID-19. A Nigerian study conducted in primary and tertiary health facilities showed that there is poor availability of infection control supplies in the studied facilities.³⁴ This strengthens the finding in the present study.

Insufficient training on infection control and lack of knowledge on the mode of transmission of COVID-19 were also reported barriers to its control. This was not surprising as COVID-19 is a novel virus that was not well understood at the onset of the pandemic. It calls for training and retraining of HCWs on such novel diseases once they come up to reduce its transmission in hospital settings. This lack of knowledge by the HCWs has been reported in other studies as a significant factor in the transmission of infection in hospitals.³⁵

a) Limitations

The data presented in our study were self-reported and partially dependent on the participants' integrity and recall ability; thus, they may be subject to recall bias. However, in despite of this limitation, our findings present valuable information on the knowledge, attitudes, and practice of HCWs during a peak period of COVID-19.

V. CONCLUSION

The advancing global threat of COVID-19 indicates that greater efforts through educational campaigns that target HCWs and the wider population are highly needed. This can be done using more effective social media.

ACKNOWLEDGMENT

We thank all the study participants for their voluntary participation and for the provision of essential information.

Authors' contributions: NHO designed the study, developed the questionnaire, collected the data, analysed the data, and prepared the manuscript. MOA also collected data, conducted the pre-test, conducted the literature review. EU contributed to literature review and filtered the data. All the authors read and approved the final manuscript.

Funding: No source of funding

Competing interests: The authors declare that they have no competing interests

Short running head: COVID-19 among healthcare workers

REFERENCES RÉFÉRENCES REFERENCIAS

1. Yin Y, Wunderink RG. MERS, SARS and other corona viruses as causes of pneumonia. *Respirology.* 2018 Feb 20; 23(2): 130-137
2. Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, et al. Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus-Infected Pneumonia. *N Engl J Med.* 2020 Mar 26; 382 (13): 1199-1207. doi: 10.1056/NEJMoa2001316
3. Lai C, Shih T, Ko W, Tang H, Hsueh P. Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and coronavirus disease-2019 (COVID-19): The epidemic and the challenges. *Int J Antimicrob Agents.* 2020 Mar; 55(3): 105924. doi: 10.1016/j.ijantimicag.2020.105924
4. Lu H, Stratton CW, Tang YW. Outbreak of pneumonia of unknown etiology in Wuhan, China: the mystery and the miracle. *J Med Virol.* 2020; 92:401-2. doi: 10.1002/jmv.25678
5. Nigeria Centre for Disease Control. NCDC 2020. Available at www.ncdc.gov.ng. Accessed 1st October 2020

6. WHO Director-General's opening remarks at the media briefing on COVID-19 - 11 March 2020. Available at <https://www.who.int/dg/speeches/detail/who-director-general-s-openingremarks-at-the-mediabriefing-on-covid-19---11-march-2020> [Accessed April 10, 2020]
7. CDC. Coronavirus disease 2019 (COVID-19). 2020. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/about/transmission.html> [Accessed on 1st March 2020].
8. World Health Organization (2020). 'Modes of transmission of virus causing COVID-19: implications for IPC precaution recommendations', Available at: <https://www.who.int/newsroom/commentaries/detail/modes-of-transmission-of-virus-causing-covid-19-implications-for-ipc-precaution-recommendations>. [Accessed 20/5/2020]
9. Salehi S, Abedi A, Balakrishnan S, Gholamrezaeizad A. 'Coronavirus Disease 2019 (COVID-19): A Systematic Review of Imaging Findings in 919 Patients', American Journal of Roentgenology. 2020; 1-7.
10. Gan WH, Lim JW, Koh D. Preventing intra-hospital infection and transmission of COVID-19 in healthcare workers. Saf Health Work 2020.
11. Nemati M, Ebrahimi B, Nemati F. Assessment of Iranian Nurses' Knowledge and Anxiety to ward COVID-19 During the Current Outbreak in Iran. Arch Clin Infect Dis. 2020; In Press. <https://doi.org/10.5812/archcid.102848>.
12. Omrani AS, Shalhoub S. Middle East respiratory syndrome coronavirus (MERS-CoV): what lessons can we learn?. J Hosp Infect. 2015; 91: 188-96.
13. WHO. A Guide to Developing Knowledge, Attitude and Practice Surveys 2008.
14. Kaliyaperumal K. Guideline for conducting a knowledge, attitude and practice (KAP) study. Community Ophthalmology. 2004 Jan -March; 4 (1):7-9.
15. MedScape. In Memoriam: Healthcare Workers Who Have Died of COVID-19. 2020. Available online at: www.medscape.com/viewarticle/927976. [Accessed September 06, 2020]
16. Koh D. Occupational risks for COVID-19 infection. Occupational medicine (Oxford, England). 2020 Mar;70(1):3.
17. World Health Organization. Rational use of personal protective equipment (PPE) for coronavirus disease (COVID-19): interim guidance, 19 March 2020. World Health Organization; 2020.
18. McEachan R, Taylor N, Harrison R, Lawton R, Gardner P, Conner M. Meta-analysis of the reasoned action approach (RAA) to understanding health behaviors. Annals of Behavioral Medicine. 2016 Aug 1; 50(4): 592-612.
19. Olum R, Chekwech G, Wekha G, Nassozi DR, Bongomin F. Coronavirus Disease-2019: Knowledge, Attitude, and Practices of Health Care Workers at Makerere University Teaching Hospitals, Uganda. Front. Public Health. 2020; 8: 181. doi: 10.3389/fpubh.2020.00181
20. Huynh G, Nguyen TNH, Tran VK, Vo KN, Vo VT, Pham LA. Knowledge and attitude toward COVID-19 among healthcare workers at District 2 Hospital, Ho Chi Minh City. Asian Pac J Trop Med. 2020; 13(6): 260-265.
21. Zhang M, Zhou M, Tang F, Wang Y, Nie H, Zhang L et al. Knowledge, attitude, and practice regarding COVID-19 among healthcare workers in Henan, China. Journal of Hospital Infection. 2020; 105: 183-187 <https://doi.org/10.1016/j.jhin.2020.04.012>
22. Fowler RA, Guest CB, Lapinsky SE, Sibbald WJ, Louie M, Tang Petal. Transmission of severe acute respiratory syndrome during intubation and mechanical ventilation. American Journal of Respiratory and Critical Care Medicine. 2004 Jun 1; 169(11): 1198-202.)
23. Tran K, Cimon K, Severn M, Pessoa-Silva CL, Conly J. Aerosol generating procedures and risk of transmission of acute respiratory infections to healthcare workers: a systematic review. PloS one. 2012 Apr 26; 7(4): e35797)
24. Ng FK, Wallace S, Coe B, Owen A, Lynch J, Bonvento Betal. From smartphone to bed-side: exploring the use of social media to disseminate recommendations from the National Tracheostomy Safety Project to front-line clinical staff. Anaesthesia. 2020 Feb; 75(2): 227-33.
25. Nickson CP. Free Open-Access Medical education (FOAM) and critical care. ICU Manag Pract. 2017;17(4):1
26. Shimizu K. 2019-nCoV, fake news, and racism. Lancet. 2020; 395: 685-6. [https://doi.org/10.1016/S040-6736\(20\)30357-3](https://doi.org/10.1016/S040-6736(20)30357-3).
27. Mejova Y, Kalimeri K. Advertisers jump on coronavirus bandwagon: Politics, news, and business. March 2020. Preprint arXiv: 2003.00923v1. Available at <https://www.researchgate.net/publication/339642155>. Accessed 1/10/2020.
28. Bhagavathula AS, Aldhaleei WA, Rahmani J, Mahabadi MA, Bandari DK. Knowledge and Perceptions of COVID-19 Among Health Care Workers: Cross-Sectional Study. JMIR Public Health Surveill. 2020; 6(2): e19160 URL: <http://publichealth.jmir.org/2020/2/e19160/> doi: 10.2196/19160
29. Olapegba PO, Ayandele O, Kolawole SO, Oguntayo R, Gandi JC, Dangiwa AL et al. A preliminary assessment of novel coronavirus (COVID-19) Knowledge and perceptions in Nigeria. SSRN Electronoc Journal. Jan 2020. doi:10.2139/ssrn.3584408. [Assessed 23/9/2020].
30. Wafaa YAW, Enas MH, MonaiA, Nashwa SH. Assessment of knowledge, attitudes and perception of healthcare workers regarding COVID 19, a cross-

sectional study from Egypt. *J. Comm Health*. 2020. Available at <https://doi.org/10.1007/s10900-020-00882-0>. [Assessed 13/9/2020]

31. Nwoga HO, Ajuba M, Ogugua I. Assessment of the knowledge, attitude and practice of standard precautions among healthcare workers in primary government health facilities in Enugu, South-East Nigeria. *Nigerian Research Journal of Clinical Sciences*. 2019; 9 (1 and 2) :16-30.
32. Shokofeh M, Farid N, Khosro F, Mahmoud F, Fatemeh H, Mehdi N. Knowledge, Attitude and Behavior of Health Care Workers in the Prevention of COVID-19. *BMC Medical Education*. DOI: 10.21203/rs.3.rs-23113/v1 [Assessed 13/9/2020]
33. Wang J, Zhou M, Liu F. Reasons for healthcare workers becoming infected with novel coronavirus disease 2019 (COVID-19) in China. *J Hosp Infect*. 2020. <http://dx.doi.org/10.1016/j.jhin.2020.03.002>
34. Nwoga HO, Ajuba M, Onwasigwe CN. Availability of infection control supplies in primary and tertiary government health facilities in Enugu Metropolis, Nigeria. *Nigerian Research Journal of Clinical Sciences*. 2019; 9(1 and 2): 43-52.
35. ChangXu H, Rebaza A, Sharma L, Dela Cruz CS. Protecting health-care workers from subclinical coronavirus infection. *Lancet Respir Med*. 2020; 8(3): e13. [http://dx.doi.org/10.1016/S2213-2600\(20\)30066-7](http://dx.doi.org/10.1016/S2213-2600(20)30066-7).

