



GLOBAL JOURNAL OF MEDICAL RESEARCH: E
GYNECOLOGY AND OBSTETRICS
Volume 21 Issue 4 Version 1.0 Year 2021
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Analysis of Cesarean Section Rate using Robson 10 Group Classification System in a Tertiary Hospital: An Observational Study

By Abhilasha Yadav, Rachna Agrawal, K. Romila Chawang & Ruchika Garg

Sarojini Naidu Medical College

Abstract- Background: The Cesarean Section rate has increased from 12% in 2000 to 21% in 2015. WHO has recommended a rate between 10% and 15% and introduced The Robson's criteria in the year 2015 as a standardized approach to determine the rate and indications of Cesarean section. To understand this and to implement effective measures to reduce cesarean section rates, WHO issued Robson's implementation manual to monitor and compare cesarean section rates in the same setting over time and among different settings.

Methods: This is a cross-sectional study of 5744 women delivering in the Obstetric Department of S.N. Medical College, Agra from April 1st through March 31st, 2020. In this study, demographic data, parity, obstetric history, fetal lie, fetal presentation, gestational age, and number of newborns for each woman were collected, grouped using a flowchart, and categorized into 10 groups. In accordance with recommendations from WHO, data were reported using "Robson's classification Report Table" and quality of data, types of population, and the CS rates were analyzed.

Keywords: cesarean section rate, robson's criteria, groups.

GJMR-E Classification: NLMC Code: WQ 430



Strictly as per the compliance and regulations of:



© 2021. Abhilasha Yadav, Rachna Agrawal, K. Romila Chawang & Ruchika Garg. This research/review article is distributed under the terms of the Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0). You must give appropriate credit to authors and reference this article if parts of the article are reproduced in any manner. Applicable licensing terms are at <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

Analysis of Cesarean Section Rate using Robson 10 Group Classification System in a Tertiary Hospital: An Observational Study

Abhilasha Yadav ^α, Rachna Agrawal ^σ, K. Romila Chawang ^ρ & Ruchika Garg ^ω

Abstract- Background: The Cesarean Section rate has increased from 12% in 2000 to 21% in 2015. WHO has recommended a rate between 10% and 15% and introduced The Robson's criteria in the year 2015 as a standardized approach to determine the rate and indications of Cesarean section. To understand this and to implement effective measures to reduce cesarean section rates, WHO issued Robson's implementation manual to monitor and compare cesarean section rates in the same setting over time and among different settings.

Methods: This is a cross-sectional study of 5744 women delivering in the Obstetric Department of S.N. Medical College, Agra from April 1st through March 31st, 2020. In this study, demographic data, parity, obstetric history, fetal lie, fetal presentation, gestational age, and number of newborns for each woman were collected, grouped using a flowchart, and categorized into 10 groups. In accordance with recommendations from WHO, data were reported using "Robson's classification Report Table" and quality of data, types of population, and the CS rates were analyzed.

Result: There was an overall CS rate of 26 percent, which is higher than the national average. The size of group 5 (15% in our institution) may be responsible for that. Group 5's contribution is very high (40%), which is another reason for the overall higher CS rate.

Conclusion: The reduction of the primary CS rate in groups 1 and 2 will result in a reduction of the overall CS rate.

Keywords: cesarean section rate, robson's criteria, groups.

I. BACKGROUND

Over the last few decades, there has been a progressive increase in the rate of cesarean section deliveries in our country. WHO has recommended that this rate should be between 10% and 15% but the driver behind this trend is not completely understood especially in developing countries.^{1,2} Immediate and long-term complications of CS included increased risk of maternal mortality and morbidity, increased need for blood transfusion, longer hospitalization,

postpartum infections, retained placenta, stillbirths, postpartum hemorrhage.

Over the last few decades, there has been a progressive increase in the rate of cesarean section deliveries in our country. WHO has recommended that this rate should be between 10% and 15% but the driver behind this trend is not completely understood especially in developing countries. (1,2). The rising cesarean section trend is a major public health concern due to potential maternal and perinatal risks associated with this. Immediate and long-term complications of CS included increased risk of maternal mortality and morbidity, increased need for blood transfusion, longer hospitalization, postpartum infections, retained placenta, stillbirths, postpartum hemorrhage.

To understand this and to implement effective measures to reduce cesarean section rates, a tool is required to monitor and compare cesarean section rates in the same setting over time and among different settings. Traditionally, at the facility level, we monitor cesarean section rates using the overall percentage of deliveries by cesarean section. But because of some intrinsic differences in hospital factors and infrastructure, a difference in characteristics of the population, and differences in clinical management protocols, this "overall cesarean section rate" becomes difficult to interpret and compare. Ideally, there should be a classification system to monitor and compare cesarean section rates. Such a system should be simple, clinically relevant, accountable, replicable, and verifiable (3,4,5). In 2015, WHO introduced Robson's criteria in the year 2015 as a standardized method to determine the rate and indications of cesarean section and issued an implementation manual(6).

- Obstetric history (parity and previous cesarean section)
- The onset of labor (spontaneous, induced, or cesarean section before the onset of labor)
- Fetal presentation or lie (cephalic, breech, or transverse)
- Number of neonates
- Gestational age (preterm or term).

Author α: (MBBS, MD) A consultant and Assistant Professor Department of Obstetrics and Gynaecology, Sarojini Naidu Medical College, Agra, Uttar Pradesh, India 282002. e-mail: yabhilasha02@gmail.com

Author σ: (MD, FICMCH). e-mail: rachanaagrawal15@gmail.com

Author ρ: (MBBS, MS). e-mail: romichawang@gmail.com

Corresponding Author ω: (MD, FICMCH).

e-mail: ruchikagargagra@gmail.com

II. MATERIALS AND METHODS

This study is a cross-sectional study of 5744 women delivered in the department of obstetrics of S.N. Medical College, Agra performed from 1st April 2020 to 31st March 2021. S.N. Medical College is a tertiary referral hospital where around 6000 deliveries take place annually.

All women who underwent cesarean section in the hospital during the specified period were included in the study.

Cases with incomplete information, doubtful gestational age, and laparotomy for uterine rupture were excluded from the study. The identity of women who underwent cesarean section was obtained from the delivery register, admission and discharge register, and operation register. The admission and

discharge register and delivery register contained information about all women who delivered in the hospital regardless of the mode of delivery (vaginal, Caesarean section) while the operation register contained only information about women who underwent Caesarean section. Using the medical registration number of each woman, we accessed all Caesarean section files performed during the study. 10 groups and their characteristics are shown in table 1. Gestational age was categorized as a term (≥ 37 weeks) or preterm (< 37 weeks). Patient's demographic data, patient's parity, obstetric history, the onset of labor, fetal presentation or lie, number of neonates, gestational age were collected. The indications for cesarean section were grouped using a flowchart (figure 1).

Table 1: The Robson Classification

1	Nulliparous women with a single cephalic pregnancy, ≥ 37 weeks gestation in spontaneous labor
2	Nulliparous women with a single cephalic pregnancy, ≥ 37 weeks gestation who had labor induced or were delivered by cesarean section before labor
2a	Cesarean section performed after induction of labor
2b	Cesarean section performed before onset of labor
3	Multiparous women without a previous cesarean section, with a single cephalic pregnancy, ≥ 37 weeks gestation in spontaneous labor
4	Multiparous women without a previous cesarean section, with a single cephalic pregnancy, ≥ 37 weeks gestation who had labor induced or were delivered by cesarean section before labor
4a	Cesarean sections performed after induction of labor
4b	Cesarean sections performed before onset of labor
5	All multiparous women with one or more previous cesarean sections, with a single cephalic pregnancy, ≥ 37 weeks gestation
5.1	With one previous cesarean section
5.2	With two or more previous cesarean sections
6	All nulliparous women with a single breech pregnancy
7	All multiparous women with a single breech pregnancy including women with previous cesarean section
8	All women with multiple pregnancies including women with previous cesarean section
9	All women with a single pregnancy with a transverse or oblique lie, including women with previous cesarean section
10	All women with a single cephalic pregnancy < 37 weeks gestation, including women with previous cesarean section




Figure 1: Flow chart for the classification of women in the Robson Classification

III. RESULT

To make the most of the information provided by the Robson Classification in local settings and to

allow comparisons between settings, the data is best reported in a standardized way (the "Robson Classification Report Table")⁷

Table 2: The Robson Classification Report Table

Group	Number of CS in group	Number of women in the group	Group Size1 (%)	Group CS rate2 (%)	Absolute group contribution to overall CS rate3 (%)	The relative contribution of a group to overall CS rate4 (%)
1	214	1750	30.43	12.23	3.73	14.21
2	204	603	10.49	33.83	3.55	13.55
3	48	1346	23.43	3.57	0.84	3.19
4	58	259	4.51	22.39	1.01	3.85
5	603	873	15.12	69.07	10.50	40.04
6	60	77	1.34	77.92	1.04	3.98
7	36	82	1.43	43.90	0.63	2.39
8	27	71	1.24	38.02	0.47	1.79
9	39	42	0.73	92.86	0.68	2.59
10	217	741	12.90	29.28	3.78	14.41
Total*	Total number CS 1506	Total number women delivered 5744	100%	Overall CS rate 26.22%	Overall CS rate 26.22%	100%

Unclassifiable: Number of cases and % [Number unclassifiable cases/(Total Number women delivered classified+unclassified)X100]

* These totals and percentages come from the data in the table.

1. Group size(%)=no of women in the group/total no of women delivered in the hospital x 100
2. Group CS rate(%)=no of CS in the group/total no of women in the group x 100
3. Absolute contribution (%)=no of CS in the group/total no of women delivered in the hospital x 100
4. Relative contribution(%)=no of CS in the group/total no of CS in the hospital x 100

A. Steps to assess the quality of data

Table 3: Steps to assess the quality of data using the Robson Classification Report Table

Steps	Robson guideline	Result	Interpretation
1. The total number of cesarean sections and women delivered in our hospital	These numbers should be the same as the total number of cesarean sections performed and of women delivered in the hospital.	57	We excluded the cases with incomplete data
2. Look at the size of Group 9 i.e. all women with single ton transverse or oblique lie.	It should be less than 1%.	0.73%	It is less than 1%
3. Look at the CS rate of Group 9	It should be 100% by convention.	92.86	3 extremely preterm pregnancies with intrauterine death of fetus were delivered vaginally

B. Steps to assess the type of population

Table 4: Steps to assess the type of population using the Robson Classification Report Table

Steps	Robson guideline	Result	Interpretation
1. Look at the size of Group1+ 2i.e. All nulliparous women ≥37 weeks gestationsing letoncephalic	This usually represents 35-42% of the obstetric population of most hospitals.	40.9%	It is within the acceptable range
2. Look at the size of Groups 3+4 i.e. All multiparous women ≥37 weeks gestation single ton cephalic, without previous CS	This usually represents about 30% of women.	26.2%	The reason for low size of Groups 3 and 4 could be that the size of Group 5 is very high which is accompanied by a high overall CS rate.
3. Look at the size of Group 5. (Multiparous women with previous cesarean section ≥37 weeks gestation with singleton cephalic pregnancy).	It is related to the overall CS rate. Group 5 usually contributes to about half of the total CS rate. In settings with low overall CS rates it is usually under10%.	15.12%	Overall CS rate is usually related to the size of group 5 and the size of this group is larger (>15%) if the institute has high CS rate in the pasty ears mainly in Groups 1and 2.
4. Look at the size of Groups6+7ie. Breeches in nulliparous women + breeches in multiparous women	It should be 3-4%	2.77%	It is within the acceptable range
5. Look at the size of Groups 8 Multiples	It should be 1.5 -2%	1.43%	It is nearly within the acceptable range
6. Look at the size of Groups 10 Preterm cephalic and singletons	It should be less than 5% in most normal risk settings.	12.9%	Can be higher in tertiary hospitals as women with high-risk factors are being referred to our hospital. These women require induction of labor, so it is accompanied by a high rate of cesarean section in this group.
7. Look at the Ratio of the size of Group1 versus Group2	It is usually 2:1 or higher	2.90	It is acceptable according to Robson.
8. Look at the Ratio of the size of Group3 versus Group 4	It is always higher than the ratio of Group1/Group2 in the same institution. This is a very reliable finding in confirming data quality and culture of the organization.	5.2	The Ratio of group3:group4 is larger than the ratio of group 1:group2 which strongly signifies the reliability of data

9. Look at the Ratio of the size of Group 6 and Group7.	The ratio of nulliparous breech/multiparous breech is usually 2:1 because breeches are more frequent in nulliparous women than in multiparous women.	Ratio 0.94	It can be because of an unusual nullipara/multipara ratio.
---	--	------------	--

C. Steps to assess the rate of cesarean section

Table 5: Steps to assess cesarean section rates using the Robson Report Table

Steps	Robson guideline	Result	Interpretation
1. Look at the CS rate for Group1	Rates under 10% are achievable	12.23%	It is slightly higher in our hospital.
2. Look at the CS rate for Group2	Consistently around 20-35%	33.83%	It is within the appropriate range.
3. Look at the CS rate for Group3	Normally, no higher than 3.0%.	3.57%	Slightly higher than 3%
4. Look at the CS rate for Group 4	It rarely should be higher than15%	22.39%	CS rates in Group 4 reflect the size and rates in 4a and 4b. If the size of Group4 is large (there are more cases of prelabour cesarean sections), the overall CS rate in Group 4 is also to be high.
5. Look at the CS rate for Group 5	Rates of 50-60% are considered appropriate.	69.07%	Rates are higher, this is due to a large number of women with 2 or more previous CS.
6. Look at the CS rate for Group8	It is usually around 60%.	38.02%	The lower rate can be due to more number of multiparous twins or due to nulliparous/multiparous with or without a previous scar.
7. Look at the CS rate in Group10	In most populations, it is usually around 30%	29.28%	It is around 30% in our institution
8. Look at the relative contribution of Groups 1,2, and 5 to the overall CS rate	These three groups combined normally contribute to 2/3rd (66%) of all Caesarean sections done in most hospitals.	These three groups combined contributed to 67.8% of all CS	These three groups are contributing to 2/3rd of all CS.
9. Look at the absolute contribution of Group 5 to the overall C.S rate		This group was responsible for 40% of all CS	This indicates that C.S. rates in Groups 1 and 2 have generally been high in previous years, and should be investigated further.

IV. DISCUSSION

Cesarean sections are becoming more common over the world, with rates rising from 12% in 2000 to 21% in 2015.⁸ In India, the caesarean section rate has been steadily growing from 8.5 percent in 2005-2006 to 17.2 percent in 2015-2016, following global trends.⁹ Many authors have expressed their satisfaction with this classification and have suggested that it be utilised more widely.^{10,11} Madhav Prasad wrote in a 2015 article that now is an ideal moment for India to use the 10-group Robsons classification to rate caesarean sections.¹² The bulk of caesarean sections were found in

groups 2 and 5, according to Deepika Jamwal et al. Group 5 was responsible for 40.3 percent of all caesarean sections, while group 2 was responsible for 29.2%.¹³ At a study conducted in a private tertiary care centre in northwest India, Priyanka D. Jogia et al. discovered that group 5 (women with a history of CS) contributed the most (37 percent) to overall surgical deliveries, with group 2 being the second highest contributor (21 percent).¹⁴ Pratima et al conducted a study of 81,784 deliveries (62,336 vaginal and 19,448 Cesarean deliveries) over 3 years. The year-wise CS rate was 22.4%, 23.5% and 25.5%, respectively. The largest contributor was by group 5 followed by group 2 and

group 1. Based on 3-year data, it was predicted in the study that the CS rate will increase by 0.905% annually in the coming 3 years. (15). During the study period, our institution's overall caesarean section rate was 26.22 percent, which looks to be higher than the national norm. Robson group 5 was the most significant single contribution to our institution's CS rate. Caesarean section rate in group 1 was higher than recommended i.e.12.23% (rate <10% are achievable) and in group 3 CS rate was 3.57% against the recommended rate of <3%. In group 5 the caesarean section rate between 50-60% is recommended but in our institution CS rate in group 5 was 69% as a large number of women with previous 2 or more caesarean sections were being admitted. The absolute contribution of group 5 is very high (40%), this indicates high CS rates. The size of group 5 is frequently connected to the overall caesarean section rate. The size of group 5 in our study was 15%, which explains our institution's overall increased caesarean section rate. It's also worth noting that Robson group 10 has grown in size (preterm deliveries). Because women with high-risk conditions are referred to our hospital, Robson group 5 can be greater in tertiary hospitals. Induction of labour is necessary for these women, hence there is a high rate of caesarean section in this group. n group 1 and group 2 in previous years.

V. CONCLUSION

Reducing the primary caesarean section rate in group 1 and group 2 will reduce the overall caesarean section rate in the institution.

BIBLIOGRAPHY

- Betrán AP, Ye J, Moller AB, et al. The increasing trend in caesarean section rates: global, regional and national estimates: 1990-2014. *PLoS One* 2016; 11: e0148343.
- World Health Organization Human Reproduction Programme, 10 April 2015. WHO statement on caesarean section rates. *Reprod Health Matters* 2015; 23: 149–50.
- Central Statistical Agency (CSA) [Ethiopia] and ICF. Ethiopia Demographic and Health Survey 2016. Addis Ababa, Ethiopia, and Rockville, Maryland, USA. Ethiopia Demographic and Health Survey: CSA and ICF, 2016.
- Fesseha N, Getachew A, Hiluf M, et al. A national review of caesarean delivery in Ethiopia. *Int J GynaecolObstet* 2011; 115:106–11.
- WHO Statement on Caesarean Section Rates. Geneva: World Health Organization; 2015 (WHO/RHR/15.02).
- WHO Robson classification Implementation manual https://www.who.int/reproductivehealth/publications/maternal_perinatal_health/robson-classification/en/
- Example of Robson report table with interpretation <https://apps.who.int/iris/bitstream/handle/10665/259513/WHO-RHR-17.22-eng.pdf?sequence=1&isAllowed=y>.
- BoermaT, et al. Global epidemiology of use of and disparities in caesarean sections. *Lancet*.2018; 392: 1341-8
- International Institute for Population Sciences. (2017). NFHS-4(National Family Health Survey-4)-2015-2016 - India fact sheet. Retrieved from URL <http://rchiip.org/nfhs/pdf/NFHS4/TG FactSheet.pdf>.
- Torloni MR, Betran AP, Souza JP, et al. Classifications for caesarean section: a systematic review. *PLoS One*. 2011; 6(1): e14566. doi: 10.1371/journal.pone.0014566. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
- Betra'n AP, Vindevoghel N, Souza JP, et al. A systematic review of the Robson classification for caesarean section: what works, doesn't work and how to improve it. *PLoS One*. 2014; 9(6):e97769. DOI: 10.1371/journal.pone.0097769. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
- Madhav Prasad et al. It is a good time to adopt 10 group Robsons classification to rate caesarean sections in India: *J ObstetGynaecol India*. 2015 Jul; 65(4): 286–287. <https://dx.doi.org/10.1007%2Fs13224-014-0656-7>
- Jamwal D et al. *Int J Reprod ContraceptObstet Gynecol*. 2021 Jun;10(6):2281-2285
- Priyanka D. Jogia et al .Analysis of caesarean sections according to modified Robson's ten group classification system at a tertiary care center in Western India. <https://dx.doi.org/10.18203/2320-1770.ijrcog20190264>.
- Pratima Mittal et al. *J ObstetGynaecol India*. 2020 Apr; 70(2): 111-118. DOI: 10.1007/s13224-019-01275-7. Epub 2019 Oct 11