



GLOBAL JOURNAL OF MEDICAL RESEARCH: H
ORTHOPEDIC AND MUSCULOSKELETAL SYSTEM
Volume 21 Issue 2 Version 1.0 Year 2021
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4618 & Print ISSN: 0975-5888
Received Date: 24th March 2021, Acceptance Date: 2nd April 2021

Managing Orthopaedic Injuries in Covid-19 Pandemic: A Concise Review

By Dr. Sagar Chaudhari, Dr. Ajit Govind Jangale,
Dr. Sheetal Ajit Jangale & Dr. Sharayu Dhande

Abstract- Conservative, non-operative therapeutic approach may provide an alternative in non-obligatory fractures in the current COVID-19 pandemic and perhaps later on as well. It may serve as a route for us to manage orthopaedic injuries till we tide over the peak of the pandemic and resume elective surgeries. Perhaps, the Coronavirus crises has given us this unique opportunity to rethink and revisit traditional methods of treating fractures and the tolerance to operate every limb fracture must be risen. We must realize that all the fractures do not always need operative management and the conservative management still has a certain place in our armamentarium of fracture management, in an evolving world.

Keywords: orthopaedic injuries, trauma, COVID-19, non-operative management, fracture, dislocation, road traffic accidents, fragility fractures, paediatric injuries.

GJMR-H Classification: NLMC Code: WE 168



Strictly as per the compliance and regulations of:



© 2021. Dr. Sagar Chaudhari, Dr. Ajit Govind Jangale, Dr. Sheetal Ajit Jangale & Dr. Sharayu Dhande. This research/review article is distributed under the terms of the Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0). You must give appropriate credit to authors and reference this article if parts of the article are reproduced in any manner. Applicable licensing terms are at <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

Managing Orthopaedic Injuries in Covid-19 Pandemic: A Consise Review

Covid - 19 and Orthopaedic Injuries

Dr. Sagar Chaudhari ^α, Dr. Ajit Govind Jangale ^σ, Dr. Sheetal Ajit Jangale ^ρ & Dr. Sharayu Dhande ^ω

Abstract- Conservative, non-operative therapeutic approach may provide an alternative in non-obligatory fractures in the current COVID-19 pandemic and perhaps later on as well. It may serve as a route for us to manage orthopaedic injuries till we tide over the peak of the pandemic and resume elective surgeries. Perhaps, the Coronavirus crises has given us this unique opportunity to rethink and revisit traditional methods of treating fractures and the tolerance to operate every limb fracture must be risen. We must realize that all the fractures do not always need operative management and the conservative management still has a certain place in our armamentarium of fracture management, in an evolving world.

Keywords: orthopaedic injuries, trauma, COVID-19, non-operative management, fracture, dislocation, road traffic accidents, fragility fractures, paediatric injuries.

I. INTRODUCTION

Coronavirus disease (COVID-19) is a novel severe acute respiratory syndrome.^{1,2} The virus was first isolated from three people with pneumonia connected to the cluster of cases in Wuhan. It was first identified in December 2019 in Wuhan, China and has spread to the rest of the world creating a global pandemic.³⁻⁸

Extensive measures have been taken by different countries, in order to reduce person-to-person transmission of COVID-19 in a variety of ways, in which the social distancing, lockdowns, curfew and self-isolation remaining common across the whole world.⁹⁻¹³

Hence, the rate of Road Traffic Accidents (RTA's) are drastically declined due to extensive lockdown, but the rate of fragility fractures continues to be unaffected, due to more prevalent osteopenia and osteoporosis with progressive ageing. The fragility fractures and traumatic fractures require robust intensive care.^{11,13-16}

Few other injuries include – falling from standing height, fall from less than 1m height like from stool, chair

Corresponding Author α: M.B.B.S, D.Orthopaedics, Consulting Orthopaedic Surgeon. e-mail: Drsagarchaudhari@Yahoo.Com

Author σ: Ms Orthopaedics, Mrcs, Assistant Professor, Dr Vasant Rao Pawar Medical College, Nashik. e-mail: Ajitjangale86@Gmail.Com

Author ρ: Bds, Mds, Pedodontist and Preventive Dentist, Private Practitioner. e-mail: Sheetalchaudhari25@Gmail.Com

Author ω: Bds, Mds Pg Student (Final Year), Department of Periodontology and Oral Implantology, M A Rangoonwala College of Dental Sciences and Research Centre, Pune. e-mail: Dhandesham1234@Gmail.Com

or bed, fall from a bicycle or a bike, fall from a height more than 1m height like roof of the house, tree etc, high energy fracture from high height, road traffic accidents.¹⁷

These accidental injuries not only increase the susceptibility of COVID-19 transmission but also aid in consumption of medical resources that have declined during the SARS-CoV-2 caused pandemic via the way of transmission through hospital.¹⁸

Simplifying the management of injuries with the use of braces and boots rather than a plaster casts along with consideration for uncemented implants and un-reamed nails, have resulted in potential advantage in decreasing the respiratory complications in patients who were infected with COVID-19. Such treatments help reduce number of visits to the hospital and also exposure to ultra-dense waiting rooms which could be a breeding ground for SARS-CoV-2.¹⁹⁻²¹

Few patients with lower limb fracture have been more susceptible to pulmonary infections with limited ambulatory capacity.^{22,23}

The COVID-19 pandemic exposed orthopaedic surgeons to manage traumatic injuries with limited resources and in safe manner whilst guarding all other healthcare professionals.²⁴⁻³⁰

According to Fineberg 2020, the patients who have to be treated should be categorized based on the COVID-19 exposure -³¹

- 1) A patient who is not known to be exposed or infected at any time
- 2) A patient who was exposed but is currently asymptomatic
- 3) A patient who has recovered from COVID and could be adequately immune
- 4) A patient who is possibly infected (persons with sign and symptoms consistent with infection who initially test negative)
- 5) A patient who is currently infected



On the basis of urgency of surgical procedures patients are categorized as:³²

Category type	Procedures should occur within
1a	24 hours
1b	72 hours
2	1month
3	3month
4	>3months

Steps in Managing a Trauma Patient with Covid-19 Symptoms or History of Contact

An Orthopaedic surgeon has to be vigilant at all times during providing pre-operative, intra-operative as well as post-operative care to refrain cross-infection amongst surgeons as well as other healthcare professionals.³³ Thermal screening for both the patient as well their attendee should be carried out, appropriate travel history, history of any previous contact should be undertaken. A three-layer surgical mask, hand sanitizer and a pair of disposable gloves should be provided at the entry point to patients along with their attendants who require emergency care.³⁴⁻³⁶ The door handles, chair handles, tables and other necessary material in the waiting areas should be regularly cleaned with 1% sodium hypochlorite atleast 4 times a day.³⁷⁻⁴⁰

A separate specialized area should be kept ready in the triage to treat COVID patients with trauma. The respective CMO's and the SMO's in the hospital should be informed immediately, in case a symptomatic patient is encountered.

Each orthopaedic surgeon along with attendants, are advised to donned PPE kits before examining every single patient, which later on should be carefully doffed off after use.⁴¹⁻⁴⁵

Resuscitate the patient and rule out all the injuries, also ask the patient to fill Informed consent, along with splintage of fracture limb.³³

All the necessary pre-operative investigations along with COVID-19 testing are advised.

If possible, the portable X-rays and ultrasound should be shifted to consulting room to avoid contamination of the radiology area and it also helps in decreasing movement of symptomatic patients.³⁵

For investigations like CT scan or MRI, we have to sterilize the respective area after investigating every patient as per centres for disease control and prevention guidelines.³⁵

Patients with closed fractures are advised to wait for surgical interventions until the COVID-19 results are out.

All cases that need urgent management like an open fracture, vascular injuries, compartment syndrome or mangled limb; and cannot wait until COVID reports. These patients should be treated as COVID positive patients unless proven otherwise and strict precautions should be taken while treating them so as to avoid transmission to healthcare professionals or to other patients.³³⁻³⁵

If the reports are positive keep the patient in the COVID isolation ward until the results are negative and take the help of the COVID response team of the hospital. If the results are negative shift the patient to the orthopaedic ward and then discharge as early as possible.³³⁻³⁵

Care must be taken during the hospital stay to physiotherapy, bedsores and DVT prevention.

II. EMERGENCY TRIAGE

Patients presented to the emergency triage with an orthopaedic emergency such as joint dislocations, compartment syndrome, open fractures, mangled extremity, polytrauma with Full Endoscopic Spine Surgery (FESS) should be managed according to a specific guidelines during global health emergencies like a pandemic of COVID-19.⁴⁶⁻⁴⁹

These orthopaedic emergencies require effective outpatient, inpatient and surgical care besides avoiding transmission of infection to fellow patients and health care givers.

The injuries that cannot be managed by the non-operative approach, should be corrected immediately with the surgical approach, with minimum usage or if possible by, completely avoiding Aerosol-Generating Procedures and with proper usage of Personal Protective Equipments with minimum assistants in the operatory.^{20,50-53}

Patient Triaging Guidelines for Orthopaedic Surgeries:⁵⁴

Orthopaedic Subspecialty	Operative Management		Non-Operative Management Indications
	Absolute Indications	Relative Indications	
Trauma & General Orthopaedics	Open Fractures Polytrauma Trauma With Neurovascular Injuries Irreducible Fracture Dislocations Compartment Syndrome Crush Injuries Septic Arthritis Acute Osteomyelitis Amputations For Gangrene	Femur Fractures (Shaft/Neck/Distal Femur) • Unstable Pelvic/ Acetabular Fractures • Intraarticular/ Forearm Fractures • Unstable Tibial Shaft Fractures • Communited/Complex Fractures • Unstable Upper Limb Fractures • Diabetic Foot	Stable Tibial Shaft Fractures • Clavicle Fractures • Stable Upper Limb Fractures • Non Unions • Malunions • Infected Non Unions • Chronic Osteomyelitis
Hand	Crush Hand • Replantation Surgeries • Infections	Tendon Injuries • Communited/ Unstable Fractures • Fracture -Dislocation • Irreducible Dislocations	Compressive Neuropathies • Tendinitis • Stable Fractures
Spine	Cauda Equina Syndrome • Epidural Abscess • Discitis Pyogenic • Spine Fracture Unstable With Paraplegia • Acute/Progressive Compressive Myelopathy	Unstable Spine Fracture With Neural Deficit • Scoliosis With Neural Deficit • Acute Radiculopathy	Low Back Pain • Neck Pain • Flat Back Syndrome • Scoliosis Without Neural Deficit • Spine Fracture Stable
Arthroplasty	Prosthetic Joint Infections • Prosthetic Joint Dislocations • Periprosthetic Fractures		Chronic Hip/Knee Pains
Orthopaedic Oncology	Infection Including Infected Joints	Sarcoma/Malignancy In Chemo/Radiation Window • Benign Aggressive Tumours Like Gct • Impending Pathological Fractures	Benign Soft Tissue Tumors • Benign Bone Tumours
Sports	Multiligamentous Injuries With Neurovascular Deficit	Multiligamentous Injury • Rotator Cuff Repairs (Young) • Major Muscle Tear	Chronic Knee, Elbow, Shoulder, Wrist, Hip Pains • Recurrent Sprains/ Dislocations • Acl/Pcl Tear

The Expert group from the Chinese Orthopaedic Association and Chinese Association of Orthopaedic Surgeons formulated an expert consensus on the diagnosis and treatment of orthopaedic emergency surgery during the outbreak of COVID-19, which has been published within the Chinese Journal of Orthopaedic Trauma in Chinese.⁵⁵

The expert consensus categorized the orthopaedic patients into four types:⁵⁵

Type I- Patients had not travelled within the in the epidemic area within 14 days and had no history of

direct or indirect contact with suspected or confirmed cases.

Type II- Patients had a history of direct or indirect contact with people from the epidemic area within 14 days, or patients board same neighbourhoods with suspected or confirmed cases. However, the patients had no clinical symptoms and every examinations were negative.

Type III- Patients are diagnosed as a suspected case of COVID-19.

Type IV- Patients were diagnosed as a confirmed case.

On the advent of COVID-19 pandemic, it has been acknowledged by the British Orthopaedic Association (BOA) emergency COVID-19 and the

National Health Service England (NHSE) guidelines to manage urgent orthopaedic and trauma conditions pragmatically balancing optimum treatment of patients against clinical safety with resource utilization.^{56,57}

Non-Operative Management of Paediatric Fractures and Dislocations during Coronavirus Crises:^{56,57}

Limb injuries in children and adolescents	Preferred Indications	Equivocal Indications	Contraindications
Hand Trauma Limb	Clavicle Fractures Proximal humerus fractures Shaft-humerus fractures with angulation of less than 45° Supracondylar fractures (Undisplaced/ minimally displaced) Extra-articular distal radius fracture Hand Fractures Reducible dislocations	Displaced Fractures. Eg. Supracondylar humerus, lateral condyle humerus Dislocations Fracture-dislocation	Unreduced dislocations Fractures with vascular deficits Compound Fractures Crush Injuries Compartment Syndrome
Lower Limb Trauma Limb	Shaft-femur fractures with acceptable angulation and displacements Fractures around Knee Shaft-tibial fractures Phalanx Fractures Metatarsal Fractures Calcaneus Fractures Reducible dislocations	Fracture – dislocations Dislocations Displaced Fractures. Eg. Neck of Femur, Shaft Femur, Fractures around the ankle	Unreduced dislocations Fractures with neurovascular deficits Compound Fractures Crush Injuries Compartment Syndrome
Pelvic Acetabular Trauma	Stable/ Minimally displaced fractures	Open book type of injuries Unstable fractures	Compound Fractures

Non-Operative Management of Fractures and Dislocations in Adults during Coronavirus Crises:^{56,57,58}

Limb and Spinal Injuries in Adults	Preferred Indications	Equivocal Indications	Contraindications
Upper limb	Clavicle fractures AC joint dislocations Scapula fractures Fractures without gross displacements; eg; proximal humerus, humeral shaft, humeral-supracondylar Extra-articular distal radius fractures Scaphoid Fractures Metacarpal Fractures Phalanx Fractures Reducible Dislocations	Fractures with significant displacement; Eg., Proximal humerus, humeral shaft, intercondylar humerus, olecranon Radius and Ulna shaft fractures Pathological Fractures Peri-prosthetic Fractures	Severe compound fractures Severe Crush Injuries Fractures involving vascular injuries Irreducible dislocations Grossly comminuted and displaced intra-articular fractures Compartment syndrome
Spine	Stable fractures	Unstable spine Fractures with neural deficit	Stable Spine Fractures Fractures with Progressive/ acute neurologic deficit
Lower Limb	Pubic rami fractures Undisplaced pelvis-acetabular Fractures Undisplaced Fractures around knee Calcaneous Fractures without gross displacement	Pelvic – acetabular Fractures with significant displacement Inter-trochanteric Fractures Intercondylar Fractures Tibial-shaft Fractures Tibial-Condyle Fractures	Fracture of neck of Femur Femur Shaft Fractures Severe Compound Fractures Severe Crush Injuries Fractures with vascular injuries Compartment syndrome Grossly comminuted and



	Metatarsal Fractures Phalanx Fractures	Patella Fractures Talar Fractures Calcaneous Fractures with gross displacement Lis Franc Injuries Peri-prosthetic Fractures Pathological Fractures	displaced intra-articular fractures Irreducible dislocations
--	---	--	--

III. CONCLUSION

The COVID-19 pandemic has substantially led to decrease in operative management of trauma, in order to optimize medical resource allocation and also to help prevent the spread of COVID. The coronavirus crises has led to depletion in the surgical volume, and preference of non-operative management of trauma over operative. On the contrary, orthopaedic surgeons must remain vigilant all the time and be prepared to provide optimal care to the injured patients.

Conflict of Interest: None

Author Contribution: All authors have equally contributed for completion of this manuscript.

REFERENCES RÉFÉRENCES REFERENCIAS

1. Di Gennaro F, Pizzol D, Marotta C, Antunes M, Racalbutto V, Veronese N, Smith L. Coronavirus diseases (COVID-19) current status and future perspectives: a narrative review. *International journal of environmental research and public health*. 2020 Jan; 17(8): 2690.
2. Kumar D, Malviya R, Sharma PK. Corona virus: a review of COVID-19. *EJMO*. 2020; 4(1): 8-25.
3. Perlman S, Netland J. Coronaviruses post-SARS: update on replication and pathogenesis. *Nature reviews microbiology*. 2009 Jun; 7(6): 439-50.
4. World Health Organization Director-General's Opening Remarks at the Media Briefing on COVID-19–11 March 2020. Available online: <https://www.who.int/dg/speeches/detail/who-director-general-s-openingremarks-at-the-media-briefing-on-COVID---11-march-2020>
5. Hsu LY, Chia PY, Lim JF. The Novel Coronavirus (SARS-CoV-2) Pandemic. *Annals Academy of Medicine Singapore*. 2020 Mar 16; 49(105): 105-7.
6. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, Zhang L, Fan G, Xu J, Gu X, Cheng Z. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The lancet*. 2020 Feb 15; 395(10223): 497-506.
7. Lu H, Stratton CW, Tang YW. Outbreak of pneumonia of unknown etiology in Wuhan, China: the mystery and the miracle. *Journal of medical virology*. 2020 Apr; 92(4): 401-2.
8. Yin Y, Wunderink RG. MERS, SARS and other coronaviruses as causes of pneumonia. *Respirology*. 2018 Feb; 23(2):130-7.
9. Rothan HA, Byrareddy SN. The epidemiology and pathogenesis of coronavirus disease (COVID-19) outbreak. *Journal of autoimmunity*. 2020 May 1; 109: 102433.
10. Zhu Y, Chen W, Xin X, Yin Y, Hu J, Lv H, Li W, Deng X, Zhu C, Zhu J, Zhang J. Epidemiologic characteristics of traumatic fractures in elderly patients during the outbreak of coronavirus disease 2019 in China. *International orthopaedics*. 2020 Aug; 44(8): 1565-70.
11. Heckman JD, McKee M, McQueen MM, Ricci W, Tornetta III P. *Rockwood and Green's fractures in adults*. Lippincott Williams & Wilkins; 2014 Sep 4.
12. Westbury LD, Syddall HE, Fuggle NR, Dennison EM, Cauley JA, Shiroma EJ, Fielding RA, Newman AB, Cooper C (2020) Long-term rates of change in musculoskeletal aging and body composition: findings from the health, aging and body composition study. *Calcif Tissue Int* (Mar 3). <https://doi.org/10.1007/s00223-020-00679-2> 5.
13. Zhang Q, Cai W, Wang G, Shen X (2020) Prevalence and contributing factors of osteoporosis in the elderly over 70 years old: an epidemiological study of several community health centers in Shanghai. *Ann. Palliat Med*. <https://doi.org/10.21037/apm.2020.02.09>.
14. Lv H, Zhang Q, Yin Y, et al. Epidemiologic characteristics of traumatic fractures during the outbreak of coronavirus disease 2019 (COVID-19) in China: A retrospective & comparative multi-center study. *Injury*. 2020; S0020-1383(20)30522-2.11.
15. Chen W, Lv H, Liu S, et al. National incidence of traumatic fractures in China: a retrospective survey of 512-187 individuals. *Lancet Glob Health*. 2017; 5(8): e807-e817.
16. Court-Brown CM, McQueen MM. Global forum: fractures in the elderly. *J Bone Joint Surg Am*. 2016; 98(9): e36-e42.
17. Zhu Y, Chen W, Xin X, Yin Y, Hu J, Lv H, Li W, Deng X, Zhu C, Zhu J, Zhang J. Epidemiologic characteristics of traumatic fractures in elderly patients during the outbreak of coronavirus disease 2019 in China. *International orthopaedics*. 2020 Aug; 44(8): 1565-70.
18. Morgan C, Ahluwalia AK, Aframian A, Li L, Sun SN. The impact of the novel coronavirus on trauma and orthopaedics in the UK. *British Journal of Hospital Medicine*. 2020 Apr 2:1-6.

18. Morgan C, Ahluwalia AK, Aframian A, Li L, Sun SN. The impact of the novel coronavirus on trauma and orthopaedics in the UK. *British Journal of Hospital Medicine*. 2020 Apr 2;1-6.
19. Hernigou J, Morel X, Callewier A, Bath O, Hernigou P. Staying home during “COVID-19” decreased fractures, but trauma did not quarantine in one hundred and twelve adults and twenty eight children and the “tsunami of recommendations” could not lockdown twelve elective operations. *International orthopaedics*. 2020 Aug; 44: 1473-80.
20. Iyengar K, Vaish A, Vaishya R. Revisiting conservative orthopaedic management of fractures during COVID-19 pandemic. *Journal of Clinical Orthopaedics and Trauma*. 2020 Jul 1; 11(4): 718-20.
21. Puzzitiello RN, Pagani NR, Moverman MA, Moon AS, Menendez ME, Ryan SP. Inflammatory and coagulative considerations for the management of orthopaedic trauma patients with COVID-19: a review of the current evidence and our surgical experience. *Journal of orthopaedic trauma*. 2020 Aug 1.
22. Chen YC, Lin WC. Risk of long-term infection-related death in clinical osteoporotic vertebral fractures: a hospital-based analysis. *PLoS One*. 2017 Aug 9; 12(8): e0182614.
23. Samsami M, Zebarjadi Bagherpour J, Nematihonar B, Tahmasbi H (2020) COVID-19 pneumonia in asymptomatic trauma patients; report of 8 cases. *Arch AcadEmerg Med* 8(1):e46.
24. Mackay ND, Wilding CP, Langley CR, Young J. The impact of COVID-19 on trauma and orthopaedic patients requiring surgery during the peak of the pandemic: a retrospective cohort study. *Bone & joint open*. 2020 Sep 1; 1(9):520-9.
25. Ambrosio L, Vadalà G, Russo F, Papalia R, Denaro V. The role of the orthopaedic surgeon in the COVID-19 era: cautions and perspectives. *Journal of Experimental Orthopaedics*. 2020 Dec; 7: 1-9.
26. Wallace CN, Kontoghiorghe C, Kayani B, Chang JS, Haddad FS. The impact of COVID-19 on trauma and orthopaedic surgery in the United Kingdom. *Bone & joint open*. 2020 Jul 1; 1(7): 420-3.
27. Oussedik S, Zagra L, Shin GY, D’Apolito R, Haddad FS. Reinstating elective orthopaedic surgery in the age of COVID-19. *The bone & joint journal*. 2020 Jul; 102(7): 807-10.
28. Awad ME, Rumley JC, Vazquez JA, Devine JG. Perioperative considerations in urgent surgical care of suspected and confirmed COVID-19 orthopaedic patients: operating room protocols and recommendations in the current COVID-19 pandemic. *JAAOS-Journal of the American Academy of Orthopaedic Surgeons*. 2020 Jun 1; 28(11): 451-63.
29. Megaloikonomos PD, Thaler M, Igoumenou VG, Bonanzinga T, Ostojic M, Couto AF, Diallo J, Khosravi I. Impact of the COVID-19 pandemic on orthopaedic and trauma surgery training in Europe. *International orthopaedics*. 2020 Sep; 44(9): 1611-9.
30. Guo X, Wang J, Hu D, Wu L, Gu L, Wang Y, Zhao J, Zeng L, Zhang J, Wu Y. Survey of COVID-19 disease among orthopaedic surgeons in Wuhan, People’s Republic of China. *The Journal of bone and joint surgery. American volume*. 2020 Apr 8.
31. Fineberg HV 2020. Ten weeks to crush the curve.
32. Re-starting non-urgent trauma and Orthopaedic care: Full guidance. *British Orthopaedic Association*.
33. Liang ZC, Chong MS, Sim MA, Lim JL, Castañeda P, Green DW, Fisher D, Ti LK, Murphy D, Hui JH. Surgical considerations in patients with COVID-19: what orthopaedic surgeons should know. *The Journal of bone and joint surgery. American volume*. 2020 Jun 3.
34. Neradi D, Hooda A, Shetty A, Kumar D, Salaria AK, Goni V. Management of orthopaedic patients during COVID-19 Pandemic in India: a guide. *Indian journal of orthopaedics*. 2020 May; 54(3): 402-7.
35. Kumar, R., & Pal, R. (2018). India achieves WHO recommended doctor population ratio: A call for paradigm shift in public health discourse! *J Fam Med Prim Care.*, 7(5), 841–844.
36. Mossa-Basha, M., Meltzer, C. C., Kim, D. C., Tuite, M. J., Kolli, K. P., & Tan, B. S. (2020). Radiology department preparedness for COVID-19: Radiology scientific expert panel. *Radiology*, 16, 200988.
37. Rutala WA, Weber DJ. Disinfectants used for environmental disinfection and new room decontamination technology. *Am J Infect Control*. 2013; 41: S36–41.
38. Donskey CJ. Does improving surface cleaning and disinfection reduce health care-associated infections? *Am J Infect Control*. 2013; 41:S12–9.
39. Rutala WA, Gergen MF, Weber DJ. Efficacy of improved hydrogen peroxide against important healthcare-associated pathogens. *Infect Control Hosp Epidemiol*. 2012; 33: 1159–61.
40. Rutala WA, Weber DJ. Disinfection, sterilization, and antisepsis: An overview. *American journal of infection control*. 2016 May 2; 44(5): e1-6.
41. Cheng L, Chen L, Xiao L, Zhang J, Cheng Y, Zhou L, Peng Y, Liu L. Problems and solutions of personal protective equipment doffing in COVID-19. *Open Medicine*. 2020 Jul 2; 15(1): 605-12.
42. Fu Q, Zhang X, Li S. Risk management strategies for occupational exposure of medical staff with novel coronavirus infections. *Chinese J Nosocomiol*. 2020; 6: 1–5. <http://kns.cnki.net/kcms/detail/11.3456.r.20200219.1629.008>.
43. Gharaibeh MA, Al Mohammad B, Gharaibeh B, Khasawneh L, Al-Naser S. Guidelines for the

- orthopedic surgeon in the era of COVID-19. *Orthopedic Reviews*. 2020 Dec 31; 12(4).
44. Chhabra HS, Bagaria V, Keny S, Kalidindi KK, Mallepally A, Dhillon MS, Malhotra R, Rajasekharan S. COVID-19: Current knowledge and best practices for orthopaedic surgeons. *Indian journal of orthopaedics*. 2020 Jul; 54: 411-25.
 45. Clement ND, Hall AJ, Makaram NS, Robinson PG, Patton RF, Moran M, Macpherson GJ, Duckworth AD, Jenkins PJ. IMPACT-Restart: the influence of COVID-19 on postoperative mortality and risk factors associated with SARS-CoV-2 infection after orthopaedic and trauma surgery. *The bone & joint journal*. 2020 Oct 1; 102(12):1774-81.
 46. Randelli PS, Compagnoni R. Management of orthopaedic and traumatology patients during the Coronavirus disease (COVID-19) pandemic in northern Italy.
 47. Rodrigues-Pinto R, Sousa R, Oliveira A. Preparing to perform trauma and orthopaedic surgery on patients with COVID-19. *The Journal of bone and joint surgery. American volume*. 2020 Apr 10.
 48. Keny S, Bagaria V, Chaudhary K, Dhawale A. Emergency and urgent orthopaedic surgeries in non-covid patients during the COVID 19 pandemic: perspective from India. *Journal of orthopaedics*. 2020 Jul 1; 20: 275-9.
 49. Liang ZC, Wang W, Murphy D, Hui JH. Novel coronavirus and orthopaedic surgery: early experiences from Singapore. *The Journal of bone and joint surgery. American volume*. 2020 Mar 20.
 50. Rodrigues-Pinto R, Sousa R, Oliveira A. Preparing to perform trauma and orthopaedic surgery on patients with COVID-19. *The Journal of bone and joint surgery. American volume*. 2020 Apr 10.
 51. Hirschmann MT, Hart A, Henckel J, Sadoghi P, Seil R, Mouton C. COVID-19 coronavirus: recommended personal protective equipment for the orthopaedic and trauma surgeon. *Knee Surgery, Sports Traumatology, Arthroscopy*. 2020 Jun; 28(6):1690-8.
 52. Simpson AH, Dall G, Haas JG. COVID-19: potential transmission through aerosols in surgical procedures and blood products. *Bone & Joint Research*. 2020 Apr; 9(4): 200-1.
 53. de Caro F, Hirschmann TM, Verdonk P. Returning to orthopaedic business as usual after COVID-19: strategies and options. *Knee Surgery, Sports Traumatology, Arthroscopy*. 2020 Jun; 28(6): 1699-704.
 54. Guidelines by Indian Orthopaedic Association: Suggestions for Orthopaedic Practice during COVID-19. Realised on 14th April 2020, 1st Update – 5th May 2020, 2nd Update – 8th June 2020.
 55. Tang PF, Hou ZY, Wu XB, Zhang CQ, Wang JW, Xing X, et al. Orthopedic Trauma Branch of Chinese Orthopaedic Association, External Fixation and Limb Reconstruction Branch of Chinese Orthopaedic Association, Trauma Expert Committee of Chinese Association of Orthopedic Surgeons. Expert consensus on management principles of orthopedic emergency in the epidemic of corona virus disease 2019. *Chin J Orthop Trauma* 2020; 22: 8–11. doi: 10.3760/cma.j.issn.1671-7600.2020.02.000.
 56. British Orthopaedic Association. Management of Patients with Urgent Orthopaedic Conditions and Trauma during the Coronavirus Pandemic; 2020. Accessed <https://www.boa.ac.uk/uploads/assets/ee39d8a8-945745339774e973c835246d/COVID-19-BOASTs-Combined-v1FINAL.pdf>. Accessed May 8, 2020.
 57. NHS England and NHS Improvement Coronavirus. Specialty Guides-Major Trauma; 2020. <https://www.england.nhs.uk/corona-virus/wp-content/uploads/sites/52/2020/03/C0070-specialty-guide-major-trauma-clinicalguide-27-march-2020.pdf>.
 58. Bram JT, Johnson MA, Magee LC, Mehta NN, Fazal FZ, Baldwin KD, Riley J, Shah AS. Where have all the fractures gone? The epidemiology of pediatric fractures during the COVID-19 pandemic. *Journal of Pediatric Orthopaedics*. 2020 Sep 1; 40(8): 373-9.