



GLOBAL JOURNAL OF MEDICAL RESEARCH: H
ORTHOPEDIC AND MUSCULOSKELETAL SYSTEM
Volume 24 Issue 1 Version 1.0 Year 2024
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Morphological Irregularity of Tracheal Rings in Horses Associated or not with Cervical Malformation (Initial Study)

By Vasconcellos, L. A. S.

Abstract- With the purpose of verifying the existence of irregularities in the tracheal rings of horses and correlating it with the presence of equine cervical vertebral malformation, 45 horses were used, including males and females of different breeds, distributed in two groups, the first being horses with lesions cervical lesion of congenital origin (GI 10 animals) and the second group composed of horses with non-congenital cervical lesions (GII 35 animals), animals from different parts of Brazil, treated in private practice over a period of 2019-2023 (4 years). It was observed that both horses without congenital lesions and those with congenital lesions had tracheal rings of different shapes that did not compromise air flow in the animals studied, and also that in all animals with congenital lesions in the cervical vertebrae, they had tracheal rings in the dorsal portion of the tracheal rings a "ridge." This was an initial study and we believe that further investigation should be carried out.

Keywords: equine, cervical radiography, tracheal ring anomalies, equine cervical vertebral malformation.

GJMR-H Classification: LCC Code: SF951, RC925, QM25



Strictly as per the compliance and regulations of:



© 2024. Vasconcellos, L. A. S. This research/review article is distributed under the terms of the Attribution-Non Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0). You must give appropriate credit to authors and reference this article if parts of the article are reproduced in any manner. Applicable licensing terms are at <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

Morphological Irregularity of Tracheal Rings in Horses Associated or not with Cervical Malformation (Initial Study)

Vasconcellos, L. A. S.

Abstract- With the purpose of verifying the existence of irregularities in the tracheal rings of horses and correlating it with the presence of equine cervical vertebral malformation, 45 horses were used, including males and females of different breeds, distributed in two groups, the first being horses with lesions cervical lesion of congenital origin (G1 10 animals) and the second group composed of horses with non-congenital cervical lesions (GII 35 animals), animals from different parts of Brazil, treated in private practice over a period of 2019-2023 (4 years). It was observed that both horses without congenital lesions and those with congenital lesions had tracheal rings of different shapes that did not compromise air flow in the animals studied, and also that in all animals with congenital lesions in the cervical vertebrae, they had tracheal rings in the dorsal portion of the tracheal rings a "ridge." This was an initial study and we believe that further investigation should be carried out.

Keywords: equine, cervical radiography, tracheal ring anomalies, equine cervical vertebral malformation.

I. INTRODUCTION

The trachea of a medium-sized adult horse is made up of a cartilaginous tube (including cartilaginous rings) 70 to 80 cm long with approximately 48 to 60 rings, with the capacity for dorsal expansion thanks to its muscles (tracheal muscle), conducting air from the larynx to the bronchioles. This anatomical structure bifurcates into two bronchi (Carina region), right and left, which in turn branch more and more within the lungs until the unit called pulmonary alveolus or alveolar sac (Vasconcellos, 2019; Budras Et Al, 2011, Clayton Et Al, 2007).

Obstructive tracheal diseases or malformations are observed in neonates with non-productive cough, without other hematological and biochemical changes, as well as in middle-aged horses (tracheal collapse) with low athletic performance (Simmons et al, 1988).

Because it is also classified as a congenital disease, equine cervical vertebral compressive malformation (ECVCM or ECVM) is a developmental defect of the neck vertebrae (cervical vertebrae) that causes narrowing of the spaces in the cervical spinal canal. It results from injuries that lead to compression of

the spinal cord and damage to the spinal cord tracts. One of the most common non-infectious causes of spinal ataxia in horses, CMVC is commonly called "wobbler syndrome", as affected horses are often unstable or uncoordinated, and may be called other names such as cervical vertebral malformation, cervical vertebral instability, cervical spondylotic myelopathy and cervical spinal stenosis (Pezanite Et Al, 2019; Bedenice Et Al, 2022; Vasconcellos, 2021).

There are also spinal compressions resulting from cervical dislocations, compressions due to fractures, lateral deviations due to osteochondritis dissecans (OCD), trauma to soft tissues (muscular and cervical ligament ruptures) but these pathologies are of traumatic origin and acquired (Manna, et al, 2023, Vasconcellos, 2021).

II. MATERIALS AND METHODS

The study was carried out over a period of 4 years (2019-2023), jockey clubs, equestrian stables, stud farms, boarding houses for horses in Brazilian territory, where all horses treated had reduced muscle strength in one or more limbs, muscular atrophies focal, hyposensitivity of the skin and limbs, ataxic (28 males and 17 females, aged between 3 and 12 years, 5 Thoroughbred horses, 30 Quarter Horses, 10 Mangalarga Marchador horses, a group with neurological problems based in the region cervical). In this study group (45 animals), 10 horses had neurological problems with non-traumatic etiology (congenital-G1) and 35 horses had acquired neurological problems (traumas, OCD, dislocations, fractures-GII), but both groups with exclusively localized neurological damage in the cervical region.

After clinical neurological examination (Vasconcellos, 2021), horses with neurological problems based in the cervical region were initially x-rayed (left or right latero-lateral position) for screening, with the purpose of separating the acquired cases (35 horses, traumas, OCD, dislocations, fractures) of the congenital cases (10 horses), and after obtaining the images, the cervical tracheal rings were also analyzed in order to verify their morphology, equality and homogeneity. Images from cervical radiographs of animals with

acquired problems and those with congenital problems were used, with the aim of comparing the tracheal rings and associating changes in them with the group with congenital problems. In atypical cases of tracheal rings identified by radiography, videoendoscopy images were taken in an attempt to visualize changes in the cartilaginous rings and record them when present.

III. RESULTS AND DISCUSSION

According to the images in figures 1 and 2, both in the group with congenital problems and in the group without congenital problems, there is a clear irregularity

of the tracheal cartilaginous rings, which does not represent any problem of a functional nature, as the important thing is that there is no resistance to the air passed through the trachea both during exhalation and inspiration and that this finding represents only an anatomical confirmation, a fact that is very little reported whether in anatomy textbooks or findings in this scientific essay.

There is indeed a change in the dorsal region of the tracheal rings seen in the radiographs of G1 (fig, 3) where there is the presence of a "ridge", seen in all animals in this group, not occurring in the GII group.



Fig. 1, 2: There is Irregularity of the Tracheal Rings (Blue Arrow)

The image recorded by videoendoscopy (fig. 4) clearly shows the change in the spaces between the rings, visualized and marked, showing a greater space between the rings, clearly showing this irregularity, but only with a visual and superficial assessment, we believe there is no interference in respiratory physiology specifically in this case. The presence of this "ridge" in the dorsal region of the tracheal rings, associated with the cervical malformation in the C6C7 vertebrae (GII), could together with the longus colli muscle, compress the recurrent laryngeal nerve (left or right) causing cases

of laryngeal hemiplegia (MAY-DAVIS et al, 2015), which we did not confirm with upper respiratory videoendoscopy, because we did not perform this procedure in all horses in GII..



Fig. 3: The Tracheal Rings Seen in the Radiographs of GI where there is the Presence of A "Ridge"(Blue Arrows)



Fig. 4: Videoendoscopy Image Showing Increased Space between the Tracheal Rings of a Horse with Cervical Problems, Non-Congenital Problems (Blue Arrow)

Another important clinical finding in GI and GII were brachial plexus compressions, muscle atrophies in the anterior limbs (right and left), hyposensitivity of the skin (fig. 5) and cervical muscles (fig. 6), (MAY-DAVIS 2015).



Fig. 5: Image showing complete skin hypoesthesia/anesthesia in a horse with a non-congenital cervical problem



Fig. 6: Cervical region with complete anesthesia resulting, in this case, from a congenital lesion

IV. CONCLUSIONS

The simple lateral projection radiographs initially showed that the tracheal rings in all horses from both groups are of different sizes and shapes, regardless of whether the animal has or does not have a neurological problem located in the cervical region. Irregularities in the cartilaginous rings ("ridge"), were present in all horses with congenital cervical malformations, that is, we found this lesion in all horses with congenital cervical problems in this study, but we suggest further studies using better investigation techniques.



With these simple findings we can indicate the importance of radiographic examination whether for the purchase of animals that will begin high-performance athletic activity or not, in agreement with DASH (2024), and in the case of animals with findings that indicate congenital problems should be prohibited from entering into a reproductive program.

horses: Case management and long-term treatment. *Equine Veterinary Education* v. 20.: p. 80–90.

REFERENCES RÉFÉRENCES REFERENCIAS

1. Bedenice D, Johnson AL. (2022) Neurologic conditions in the sport horse. *Animal Frontier*. v.14; 12(3): 37-44.
2. Budras, K. D.; Sack, W. O.; Rock, S. (2011) *Anatomy of the Horse*. Schlütersche Verlagsgesellschaft mbH & Co. KG. Hanover. Germany. 209 p.
3. Clayton, H. M.; Flood, P. F.; Resenstien, D. S. (2007) *Anatomia Clínica del Caballo*. Elsevier. Madrid. Espana. 135 p.
4. Dash, R. F.; Morgan, R. E. (2024) Should cervical radiographs be included in a pre-purchase examination? *Equine Veterinary Education*. v.00, p.1-9.
5. Mannaa M, Shamaa A. A, Shawky A, Hassan I. M., Refaey A. M., Abu-Seida A. M. (2023). A novel surgical technique for treatment of cervical vertebral stenotic myelopathy (wobbler syndrome) in a filly (2023). *Journal of Equine Veterinary Science*. v. 126: 104493.
6. May-Davis, S.; Walker, C. (2015) Variations and Implications of the gross morphology in the Longus colli muscle in Thoroughbred and Thoroughbred derivative horses presenting with a congenital Malformation of the 6th and 7th cervical vertebrae. *Journal of Equine Veterinary Science*. v.37, i.7, p.560-568.
7. Pezzanite L, Easley J. (2019) Update on Surgical Treatment of Wobblers. (2019). *Veterinary Clinics of North America: Equine Practice*. v.35 (2): 299-309.
8. Siger, L., Hawkins, J., Andrews, F., Henry, R. (1998) Tracheal stenosis and collapse in horses. *Compendium on Continuing Education for the Practising Veterinarian* 20: 628–635.
9. Simmons, T.; Petersen M.; Parker J, Dietze, A; Rebhun W. (1988) Tracheal collapse due to chondrodysplasia in a miniature horse foal. *Equine Practice* 10: 39–41.
10. Vasconcellos, L. A. S. (2019). *Problemas Respiratórios nos Equinos e seus Métodos de Diagnóstico por Imagens*. Editora MedVet. São Paulo. Brasil. 107p.
11. Vasconcellos, L. A. S. (2022). *Neurologia e Neurocirurgia Equina: Princípios gerais*. Editora Lusófona. Lisboa, Portugal. 216 p.
12. Wong, D. M.; Sponseller, B. A.; Riedesel, E. A.; Couetil, L. L.; Kersh. (2008) the use of intraluminal stents for tracheal collapse in two