



Posterior Reversible Encephalopathy Syndrome Secondary to Arterial Hypertension: A Case Report from Lubumbashi, Democratic Republic of Congo

Article Record

Rivain Iteke Fefe^{§*}

*Corresponding Author



Bugeme Marcellin^{‡*}



Jacques Musung Mbaz[¶]

Franck Omangelo Shongo[‡]

Muyumba Kiyana
Emmanuel[¶]

[§] Department of Anesthesia and Intensive Care, University of Lubumbashi, Lubumbashi, Cd

[‡] Department of Neuropsychiatry, University of Lubumbashi, Lubumbashi, Cd

[¶] Department of Internal Medicine, University of Lubumbashi, Lubumbashi, Cd

RECEIVED

2026-03-30

ACCEPTED

2026-04-09

ONLINE PUBLISHED

2026-07-14

PEER REVIEW

Double Blind

Abstract

Posterior reversible encephalopathy syndrome (PRES) is a clinical-radiological entity characterized by the association of neurological clinical signs. The most common causes are hypertensive encephalopathy, renal failure, immunosuppressants, infections and systemic diseases. The clinical course is most often favorable with a more or less rapid and usually complete improvement of clinical signs and a disappearance of radiological images. We report a case of a 61-year-old chronically hypertensive patient who presented with a clinical picture of confusion, in whom brain CT (computed tomography) revealed images suggestive of an PRES. The evolution was quickly favorable under antihypertensive treatment.

Posterior reversible encephalopathy

confusion

hypertension

Lubumbashi

AI USE STATEMENT

No generative AI was used for analysis or results.

FUNDING

No external funding was declared for this work.

CONFLICT OF INTEREST

no conflict of interest is declared by the authors

DATA AVAILABILITY

Not applicable for this article.

ETHICS

No ethics committee approval was required for this article type.

CONSENT

Not applicable for this article.

TRIAL REG.

Not applicable.

Crossref DOI: 10.34257/GJMRA255356

How to Cite: Fefe et al. (2026). Posterior Reversible Encephalopathy Syndrome Secondary to Arterial Hypertension: A Case Report from Lubumbashi, Democratic Republic of Congo. Global Journal of Medical Research, 26(1), 16-17.

DOI: 10.34257/GJMRA255356

LICENSE

© 2026 Global Journals. Open-access article under CC BY-NC-ND 4.0 International License.



Print ISSN 0975-5888



Online ISSN 2249-4618



Under the strict compliance and defined process of



METADATA CONTINUATION

AUTHOR CONTACT QR LEDGER

Rivain Iteke Fefe\$*	
Bugeme Marcellin\$*	

ARCHIVAL RECORD

Posterior Reversible Encephalopathy Syndrome Secondary to Arterial Hypertension: A Case Report from Lubumbashi, Democratic Republic of Congo

Rivain Iteke Fefe^{§*}, Bugeme Marcellin[‡], Jacques Musung Mbaz[¶], Franck Omangelo Shongo[‡], and Muyumba Kiyana Emmanuel[¶]

Affiliations

[§] Department of Anesthesia and Intensive Care, University of Lubumbashi, Lubumbashi, Cd

[‡] Department of Neuropsychiatry, University of Lubumbashi, Lubumbashi, Cd

[¶] Department of Internal Medicine, University of Lubumbashi, Lubumbashi, Cd

Abstract

Posterior reversible encephalopathy syndrome (PRES) is a clinical-radiological entity characterized by the association of neurological clinical signs. The most common causes are hypertensive encephalopathy, renal failure, immunosuppressants, infections and systemic diseases. The clinical course is most often favorable with a more or less rapid and usually complete improvement of clinical signs and a disappearance of radiological images. We report a case of a 61-year-old chronically hypertensive patient who presented with a clinical picture of confusion, in whom brain CT (computed tomography) revealed images suggestive of an PRES. The evolution was quickly favorable under antihypertensive treatment.

Keywords: *Posterior reversible encephalopathy, confusion, hypertension, Lubumbashi*

* Corresponding Author
Rivain Iteke Fefe

DOI
10.34257/GJMRA255356

1. Introduction

Posterior reversible encephalopathy syndrome (PRES), also known as posterior reversible encephalopathy syndrome, is a radioclinical entity characterized by reversible central nervous system dysfunction and typical brain MRI or CT imaging [1]. This syndrome was first described in 1992 and brought to light by Hinchey et al. in 1996 [2,3,4]. PRES is generally a rare complication of a sudden increase in blood pressure occurring in a chronically hypertensive patient [5]. We report a case of a patient who developed PRES secondary to hypertension.

The patient received a treatment of Amlodipina at a dose of 10 mg per day. One week into treatment, his condition was marked by a normal neurological examination. We concluded that he had posterior progressive encephalopathy syndrome secondary to hypertension.

2. Case Report

This is a 61-year-old patient transferred from an intensive care unit for a behavioral disturbance characterized by monologue, visual hallucinations, agitation, verbal aggression, and acting out. These symptoms had been developing rapidly for five days. He had a history of poorly controlled hypertension. Neurological examination revealed disorientation, anterograde and retrograde amnesia, acalculia, and a vacant stare. The remainder of the neurological examination was normal. His blood pressure was 170/112 mmHg. Other vital signs were normal. A diagnosis of delirium was initially considered. Brain CT scans performed without contrast (Fig. 1) and with contrast (Fig. 2) revealed a hypodense edematous infiltrate within the cortical sulci in the parietal, posterior temporal, and occipital lobes, bilaterally and symmetrically. Minimal hypodensity of the periventricular corneal white matter and the centrum semiovale was also observed, indicating leukoaraiosis.




Figure 1. Brain CT scan without contrast showing hypodensity in the posterior regions, symmetrically and bilaterally.



Figure 2. Brain CT scan with contrast, shows hypodensity in the posterior regions symmetrically and bilaterally.

3. Discussion

Posterior reversible encephalopathy syndrome (PRES) is a rare but severe disease of the central nervous system. Its onset and presentation are characterized by significant clinical polymorphism, sometimes making diagnosis difficult [1]. Numerous predisposing factors exist, including hypertension. This is a classic finding and was the first factor described. Moderate hypertension has been observed in 75

In our case, hypertension was the identified predisposing factor. Other circumstances may include eclampsia, chemotherapy, septicemia, chronic renal failure, autoimmune diseases, etc. [1,8]. The clinical presentation is nonspecific, neuroradiological signs are characteristic, and the triggering factors or associated pathophysiological conditions are numerous [5]. Neurological manifestations are varied and frequently include seizures, sometimes complicated by status epilepticus, headaches, confusion, nausea, and vomiting [6]. These clinical manifestations depend on the associated pathology. Hypertension with a diastolic blood pressure greater than 120 mmHg is usually observed [9]. The clinical presentation of our patient was mental confusion. The therapeutic strategy depends on the etiology and clinical presentation of PRES. Discontinuation of the triggering or aggravating factor is the first therapeutic measure [5]. Controlling hypertension is the primary aspect of treatment. This involves the usual antihypertensive agents, calcium channel blockers, beta-blockers, and diuretics. The therapeutic goal is to maintain a mean arterial pressure between 105 and 125 mmHg, without reducing this pressure by more than 25

4. Conclusion

Diagnostic workup must absolutely identify the cause of PRES in order to implement appropriate corrective measures. Therefore, PRES should be considered in any case of acute encephalopathy. The diagnosis is generally appropriate in the presence of hypertension or other predisposing factors. Knowledge of this syndrome should encourage the avoidance of unnecessary repeat imaging studies when the clinical course is favorable.

■ REFERENCES

- [1] Hugonnet, E., Da Ines, D., Bobby, H., Claise, B., Petitcolin, V., Lannareix, et al. Syndrome d'encéphalopathie postérieure réversible (EPR): aspects en imagerie TDM et IRM. *Journal de Radiologie diagnostique et interventionnelle*. 2013; 94(1): 44-51.
- [2] Schwartz RB, Jones KM, Kalina P, et al. Hypertensive encephalopathy: findings on CT, MR imaging, and SPECT imaging in 14 cases. *AJR Am J Roentgenol*. 1992; 159:379-83
- [3] Schwartz RB, Bravo SM, Klufas RA, et al Cyclosporine neurotoxicity and its relationship to hypertensive encephalopathy: CT and MR findings in 16 cases. *AJR Am J Roentgenol*. 1995;165: 627-31
- [4] Hinchey J, Chaves C, Appignani B, et al. A reversible posterior leukoencephalopathy syndrome. *N Engl J Med*. 1996; 334: 494-500
- [5] Amoli, A. M., Mégarbane, B., Chabriat, H. La leucoencéphalopathie postérieure réversible. *Réanimation* 2007; 16(6), 490-497.
- [6] Bartynski WS. Posterior reversible encephalopathy syndrome, part 1: fundamental imaging and clinical features. *AJNR Am J Neuroradiol*. 2008; 29: 1036-1042.
- [7] Lee VH, Wijidicks EF, Manno EM, Rabinstein AA. Clinical spectrum of reversible posterior leukoencephalopathy syndrome. *Arch Neurol* 2008; 65: 205-210.
- [8] Amagada JO, Kondagunta H, Afshan N, Watermeyer S, Jones R. Posterior reversible encephalopathy syndrome secondary to eclampsia. *J Obstet Gynaecol* 2008; 28: 646-7.
- [9] Schwartz RB. Hyperperfusion encephalopathies: hypertensive encephalopathy and related conditions. *Neurologist*. 2002; 8: 22-34
- [10] Servillo G, Bifulco F, De Robertis E, Piazza O, Striano P, Tortora F, et al. Posterior-reversible encephalopathy syndrome in intensive care medicine. *Intensive Care Med* 2007; 33: 230-6.
- [11] Gokce M, Dogan E, Nacitarhan S, Demirpolat G. Posterior reversible encephalopathy syndrome caused by hypertensive encephalopathy and acute uremia. *Neurocrit Care* 2006; 4: 133-6