



GLOBAL JOURNAL OF SCIENCE FRONTIER RESEARCH
ENVIRONMENT & EARTH SCIENCES
Volume 12 Issue 3 Version 1.0 Year 2012
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Environmental Impacts of Oil Exploration and Exploitation in the Niger Delta of Nigeria

By Kadafa, Adati Ayuba

Universiti Putra Malaysia

Abstract - Problem Statement : Oil exploration and exploitation has been on-going for several decades in the Niger Delta. It has had disastrous impacts on the environment in the region and has adversely affected people inhabiting that region. The Niger Delta consist of diverse ecosystems of mangrove swamps, fresh water swamps, rain forest and is the largest wetland in Africa and among the ten most important wetland and marine ecosystems in the world, but due to oil pollution the area is now characterized by contaminated streams and rivers, forest destruction and biodiversity loss in general the area is an ecological wasteland. This affects the livelihood of the indigenous people who depend on the ecosystem services for survival leading to increased poverty and displacement of people. The oil industry located within this region has contributed immensely to the growth and development of the country which is a fact that cannot be disputed but unsustainable oil exploration activities has rendered the Niger Delta region one of the five most severely petroleum damaged ecosystems in the world. Studies have shown that the quantity of oil spilled over 50 years was a least 9-13 million barrels, which is equivalent to 50 Exxon Valdez spills.

Keywords : ecosystem, oil, pollution, niger delta.

GJSFR-H Classification : FOR Code: 850104, 850103



Strictly as per the compliance and regulations of :



© 2012. Kadafa, Adati Ayuba. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License (<http://creativecommons.org/licenses/by-nc/3.0/>), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Environmental Impacts of Oil Exploration and Exploitation in the Niger Delta of Nigeria

Kadafa, Adati Ayuba

Abstract - Problem Statement : Oil exploration and exploitation has been on-going for several decades in the Niger Delta. It has had disastrous impacts on the environment in the region and has adversely affected people inhabiting that region. The Niger Delta consist of diverse ecosystems of mangrove swamps, fresh water swamps, rain forest and is the largest wetland in Africa and among the ten most important wetland and marine ecosystems in the world, but due to oil pollution the area is now characterized by contaminated streams and rivers, forest destruction and biodiversity loss in general the area is an ecological wasteland. This affects the livelihood of the indigenous people who depend on the ecosystem services for survival leading to increased poverty and displacement of people. The oil industry located within this region has contributed immensely to the growth and development of the country which is a fact that cannot be disputed but unsustainable oil exploration activities has rendered the Niger Delta region one of the five most severely petroleum damaged ecosystems in the world. Studies have shown that the quantity of oil spilled over 50 years was a least 9-13 million barrels, which is equivalent to 50 Exxon Valdez spills.

Approach : This paper aims to investigate the various environmental problems associated with oil exploration and spillage in specifically the Niger Delta in Nigeria, as well as bring into perspective the environmental impact occurring in an important, reproductive wetland and marine ecosystem. Secondary data was used and analysed using descriptive method. Towards obtaining logical deductions, sequential presentation of facts from the data and a clear picture of the problem.

Results : There is no doubt that oil has been of great benefit to the Nigerian state and the people as a whole but the oil pollution caused by spillages from the oil industry located primarily in the Niger Delta region has caused the massive destruction to farmlands, sources of drinking water, mangrove forest, fishing grounds and declination of fish, crabs, molluscs, periwinkles and birds. Large areas of mangrove forest have been destroyed over a wide area affecting terrestrial and marine resources. Some past spills have necessitated the complete relocation of some communities, loss of ancestral homes, pollution of fresh water, loss of forest and agricultural land, destruction of fishing grounds and reduction of fish population, which is the major source of income for the Niger Delta people.

Conclusion : The activities that come with the oil exploration and exploitation causes alterations to the environment. Which significantly have negative effects; some of the effects that come with petroleum development can be reduced or prevented basically by taking some steps in terms of prevention. Monitoring is also essential, but is lacking in the

Niger Delta region. Monitoring the location of the oil companies; the terrain, the accessibility, revenue, man power availability for the monitoring agency, qualified personnel are not available. This restricts the ability and efficiency of monitoring by the government.

Recommendation : Updating and revising the legislations, reviewing the license of the oil companies and reviewing the fines will go a long way in ensuring compliance, even though the government cannot systematically or frequently monitor these sites

Keywords : ecosystem, oil, pollution, niger delta.

I. INTRODUCTION

August 1859, Colonel Drake drilled a 70 feet well in Titusville, Pennsylvania and discovered oil. By the 1800's a number of wells were drilled in Pennsylvania, Kentucky and California. The birth of the modern oil industry is credited to the discovery oil at Spindletop in 1901 atop a salt dome near Beaumont Texas (Knowles, 1983). Oil and natural gas are dominant fuel sources in the U.S economy it provides 62% of the nation's energy and about 100% of its transportation fuels this is also similar for many other nations (NEPDG, 2001). Oil spillage is a global issue that has been occurring since the discovery of crude oil, which was part of the industrial revolution. In 1956, Shell British Petroleum (now Royal Dutch Shell) discovered crude oil at a village Oloibiri in Bayelsa state located within the Niger Delta of Nigeria (Onuoha, 2008; Anifowose, 2008) and commercial production began in 1958. Oil exploration and exploitation has been on-going for several decades in the Niger Delta. It has had disastrous impacts on the environment in the region and has adversely affected people inhabiting that region. The Niger Delta is among the ten most important wetland and marine ecosystems in the world. The oil industry located within this region has contributed immensely to the growth and development of the country which is a fact that cannot be disputed but unsustainable oil exploration activities has rendered the Niger Delta region one of the five most severely petroleum damaged ecosystems in the world. Studies have shown that the quantity of oil spilled over 50 years was a least 9-13 million barrels, which is equivalent to 50 Exxon Valdez spills (FME, et. al. 2006). The Niger Delta consist of diverse ecosystems of mangrove swamps, fresh water swamps, rain forest and is the largest wetland in Africa but due to oil pollution the area is now characterized by contaminated streams and rivers, forest

Author : Department of Environmental Management, Faculty of Environmental Studies, Universiti Putra Malaysia, 43400 DarulEshan, Serdang, Selangor, Malaysia. E-mail : kwaala@yahoo.com

destruction and biodiversity loss in general the area is an ecological wasteland. This affects the livelihood of the indigenous people who depend on the ecosystem services for survival.

This paper aims at the various environmental problems associated with oil exploration and spillage in specifically the Niger Delta in Nigeria. As well as bring into perspective the environmental impact occurring in an important, reproductive wetland and marine ecosystem. Contribute to the Federal Government of Nigeria, States, Local Governments of the Niger Delta region, Oil Companies, Nigerian National Petroleum Cooperation, Institutions, Host Communities, Researchers and the existing body of knowledge on oil spillage and environmental degradation in the Niger Delta region of Nigeria. It will enlighten and sensitize relevant authorities on the problem within the Niger Delta region of Nigeria with recommendations that would be made arising from the study, which will help policy makers on future plans. Secondly, bring into perspective the environmental impact occurring in an important, reproductive wetland and marine ecosystem and assess the extent of the environmental degradation that has occurred to date. Thirdly, serve as a useful reference material and catalyst that will stimulate future researchers.

II. MATERIALS AND METHODS

This involved obtaining data from past and present studies, government and non-government bodies and existing literature (Twumasi and Merem, 2006; Uyigüe and Agho, 2007; Uyigüe and Ogbeibu, 2007). The study relied on secondary data, data was obtained from The Nigerian National Petroleum Cooperation, World Bank Reports, National Bureau of Statistics, United Nations Environmental Protection Programme, Amnesty International, International Monetary Fund, Published and Unpublished materials, Books, Newspapers, Conference and Seminar Papers, Journals and the internet. The data obtained was analysed using descriptive method to obtain logical deductions and sequential presentation of facts from the data obtained to give a clear picture of the problem.

a) Background on the Niger delta region

Nigeria has a coastal line of approximately 85km towards the Atlantic Ocean lying between latitude 4°15' to 4°50' and longitude 5°25' to 7°37' with a land mass of about 28000sq/km area within the coastal region. The surface area of the continental shelf is 46300sq/km. The coastal areas consist of freshwater swamp, mangrove swamp, beach ridges, sand bars, lagoons marshes and tidal channels. Nigeria has a total land mass of 923,768sq/km; 918,768sq/km being terrestrial land and 13000 sq. /km being aquatic (CIA World Fact Book). The coastal area is humid with a mean average temperature of 24-32°C and coastal area has an average annual rainfall ranging between 1,500-

4,000m (Kuruk, 2004). Nigeria has two large rivers; the Niger-Benue and the Chad River. There are several rivers that channel into the Atlantic Ocean directly, all other flowing waters flow into the Chad basin or into the lower Niger to the sea eventually (Kuruk, 2004).

The Niger Delta is located in the Atlantic coast of Southern Nigeria and is the world's second largest delta with a coastline of about 450km which ends at Imo river entrance (Awosika, 1995). The region is about 20,000sq/km as it is the largest wetland in Africa and among the third largest in the world (Powell, et al., 1985; CLO, 2002; Anifowose, 2008; Chinweze and Abiola-Oloke, 2009). 2,370sq/km of the Niger Delta area consists of rivers, creeks, estuaries and stagnant swamps cover approximately 8600sq/km, the Delta mangrove swamp spans about 1900sq/km as the largest mangrove swamp in Africa (Awosika1995). The Niger Delta is classified as a tropical rainforest with ecosystems comprising of diverse species of flora and fauna both aquatic and terrestrial species. The region can be classified into four ecological zones; coastal inland zone, freshwater zone, lowland rainforest zone, mangrove swamp zone and this region is considered one of the ten most important wetlands and marine ecosystems in the world (FME, et al., 2006; ANEEJ, 2004). As of 1991, the National Census estimated about 25% of the entire Nigerian population lives within the Niger Delta region (Twumasi and Merem, 2006; Uyigüe and Agho, 2007). The Niger Delta region has a steady growing population of approximately 30 million people as of 2005, accounting for more than 23% of Nigeria's total population Twumasi and Merem, 2006; Uyigüe and Agho 2007).

b) Oil Production in Nigeria

Nigeria has been a member of Organization of Petroleum Exporting Countries (OPEC) since 1971. It has the largest natural gas reserve in Africa, has the second largest oil reserve in Africa and is the African continent's primary oil producer. As of the 1980s oil revenue provided 90% of Nigeria foreign exchange earnings and 85% of the government revenue (Odeyemi and Ogunseitan 1985), with estimated reserves extending beyond 20-30 years (NNPC, 1984). Shell D'Arcy the pioneer oil company in Nigeria, which started commercial production in 1958 with a production rate of 5100 barrels per day and a peak production of 2.44 million barrels per day over the next few years (Amu, 1997). According to NNPC (1984) through OPEC, production rates dropped to 1.5 million barrels per day from the activities of 10 international companies working 122 fields, containing over 970 oil wells. Nigeria has four oil refineries with an estimated total refining capacity of 445,000 barrels per day (Onuoha, 2008; Anifowose, 2008). The first and oldest being the Port Harcourt refinery, commissioned in 1965. It had an initial capacity of 35,000 barrels per day, which was later expanded to

60,000 barrels per day of light crude oil. The Port Harcourt refinery has a second refinery with a capacity of 150,000 barrels per day (Odeyemi and Ogunseitan 1985; Ukoli 2005). Anifowose (2008) and Onuoha (2008) cited in their studies that the region has about 606 oil fields with 355 situated onshore; 251 situated offshore with 5,284 drilled oil wells and 7,000km of oil and gas pipelines.

c) *Biodiversity in the Niger Delta*

The ecosystem of the area is highly diverse and supportive of numerous species of terrestrial and aquatic fauna and flora as well as human life (Uyigue 2009). The Niger Delta has been declared as a key zone for the conservation of the western coast of Africa on the basis of its extraordinary biodiversity (Nenibarini, 2004). It is estimated, that in Nigeria there are more than 46,000 plant species of which about 205 are endemic, and approximately 484 plants in 112 families are threatened with extinction as well as many animal and bird species (Salau, 1993). Another estimate is that 24 out of 274 mammal, 10 out of 831 birds and 2 out of 114 reptiles known to exist in Nigeria are endangered (WRI, 1992). The larger population of the Niger Delta survive on services provided by the ecosystem; agriculture, industry, fishing, food, drinking water, wood, shelter, medicine, employment and aesthetics. All aspect of oil exploration and exploitation has adverse effects on the ecosystem and the local biodiversity. Oil exploration by seismic oil companies involves clearing of seismic lines, dynamiting for geological excavation, which affects the aquatic environment. It causes mortality in fauna, turbidity in the water that blockage of gills of the filter feeders in the benthic fauna, reduction of photosynthetic activity caused by the water turbidity that reduces the amount of sunlight penetration.

Oil and gas pipeline have been installed covering 7,000km to enhance the distribution crude oil products to other parts of the country (Onuoha, 2008). The installation of these pipelines involved clearing large areas of habitat to make pipeline tracks. These pipelines run across the rainforests and mangroves with incidences of leakage and rupture and accidental discharges. These discharges are caused by vandalism, failure of pipeline integrity due to aging and defects in material. Most incidences of the reported oil spillages have occurred in the mangrove swamp forest, which is one of the most reproductive ecosystems rich in fauna and flora (Nenibarini, 2004).

d) *Oil Spillages*

An estimated 9 million- 13 million (1.5 million tons) of oil has been spilled in to the Niger Delta ecosystem over the past 50 years; 50 times the estimated volume spilled in Exxon Valdez oil spill in Alaska 1989 (FME, NCF, WWF UK, CEESP-IUCN 2006). The first oil spill in Nigeria was at Araromi in the present Ondo state in 1908 (Tolulope, 2004). In July 1979 the

Forcados tank 6 Terminal in Delta state incidence spilled 570,000 barrels of oil into the Forcados estuary polluting the aquatic environment and surrounding swamp forest (Ukoli, 2005; Tolulope, 2004). The Funiwa No.5 Well in Funiwa Field blew out an estimate 421,000 barrels of oil into the ocean from January 17th to January 30th 1980 when the oil flow ceased (Ukoli, 2005; Gabriel, 2004; Tolulope, 2004), 836 acres of mangrove forest within six miles off the shore was destroyed. The Oyakama oil spillage of 10th may 1980 with a spill of approximately 30,000bbl (Ukoli, 2005).

In August 1983 Oshika village in River state witnessed a spill of 5,000 barrels of oil from Ebocha-Brass (Ogada-Brass 24) pipeline which flooded the lake and swamp forest, the area had previously experienced an oil spill of smaller quantity; 500 barrels in September 1979 with mortality in crabs, fish and shrimp. Eight months after the occurrence of the spill there was high mortality in embryonic shrimp and reduced reproduction due to oil in the lake sediments (Gabriel, 2004). The Ogada-Brass pipeline oil spillage near Etiama Nembe in February 1995 spilled approximately 24,000 barrels of oil which spread over freshwater swamp forest and into the brackish water mangrove swamp. The Shell Petroleum Development Company (SPDC) since 1989 recorded an average of 221 spills per year in its operational area involving 7,350 barrels annually (SPDC Nigeria Brief, May 1995:3). From 1976-1996 a total of 4647 oil spill incidences spilling approximately 2,369,470 barrels of oil into the environment of which 1,820,410.5 (77%) were not recovered. Most of these oil spill incidences in the Niger Delta occur on land, swamp and the offshore environment (Nwilo and Badejo 2005a, 2005b, 2004; Twumasi and Merem, 2006; Uyigue and Agho 2007). NNPC estimates 2,300 cubic meters of oil has spilled in 300 separate incidences annually between 1976-1996 (Twumasi and Merem, 2006). Table I below show some of the oil polluted sites in the Niger Delta region.

Table 1: Some Severely Oil Polluted Sites in the Niger Delta

| Location | Environment | Impacted Area (ha) | Nature of Incidence |
|---------------|-------------------------------------|--------------------|------------------------------|
| Bayelsa State | | | |
| Biseni | Freshwater Swamp Forest | 20 | Oil Spillage |
| Etiama/Nembe | Freshwater Swamp Forest | 20 | Oil Spillage & Fire Outbreak |
| Etelebu | Freshwater Swamp Forest | 30 | Oil Spill Incidence |
| Peremabiri | Freshwater Swamp Forest | 30 | Oil Spill Incidence |
| Adebawa | Freshwater Swamp Forest | 10 | Oil Spill Incidence |
| Diebu | Freshwater Swamp Forest | 20 | Oil Spill Incidence |
| Tebidaba | Freshwater Swamp Forest Mangrove | 30 | Oil Spill Incidence |
| Nembe creek | Mangrove Forest | 10 | Oil Spill Incidence |
| Azuzuama | Mangrove | 50 | Oil Spill Incidence |
| 9 sites | | | |
| Delta State | | | |
| Opuekebe | Barrier Forest Island | 50 | Salt Water Intrusion |
| Jones Creek | Mangrove Forest | 35 | Spillage & Burning |
| Ugbeji | Mangrove | 2 | Refinery Waste |
| Ughelli | Freshwater Swamp Forest | 10 | Oil Spillage-Well head leak |
| Jesse | Freshwater Swamp Forest | 8 | Product leak/Burning |
| Ajato | mangrove | | Oil Spillage Incidence |
| Ajala | Freshwater Swamp Forest | | Oil Spillage Incidence |
| Uzere | Freshwater Swamp Forest | | Oil Spillage Incidence |
| Afiesere | Freshwater Swamp Forest | | Oil Spillage Incidence |
| Kwale | Freshwater Swamp Forest | | Oil Spillage Incidence |
| Olomoro | Freshwater Swamp Forest | | QC |
| Ughelli | Freshwater Swamp Forest | | Oil Spillage Incidence |
| Ekakpare | Freshwater Swamp Forest | | Oil Spillage Incidence |
| Ughuvwughe | Freshwater Swamp Forest | | Oil Spillage Incidence |
| Ekerejegbe | Freshwater Swamp Forest | | Oil Spillage Incidence |
| Ozoro | Freshwater Swamp Forest | | Oil Spillage Incidence |
| Odimodi | Mangrove Forest | | Oil Spillage Incidence |
| Ogulagha | Mangrove Forest | | Oil Spillage Incidence |
| Otorogu | Mangrove Forest | | Oil Spillage Incidence |
| Macraba | Mangrove Forest | | Oil Spillage Incidence |
| 20 sites | | | |
| Rivers State | | | |
| Rumuokwurusi | Freshwater Swamp | 20 | Oil Spillage |
| Rukpoku | Freshwater Swamp | 10 | Oil Spillage |

Source: FME, NCF, WWF UK, CEEP-IUCN 2006 Niger Delta Resource Damage Assessment and Restoration Project.

The Punch Newspaper on February 20, 1991:2 reported a total of 2,796 oil spill incidences recorded between the periods of 1976-1990 leading to 2,105,393 barrels of oil spilled. The UNDP 2006:181 also reported that between the period of 1976-2001, 3 million barrels of oil were lost in 6,817 oil spill incidences of which over 70% of the spilt oil was not recovered. In 2001 the western operations of the Shell Petroleum Development Company (SPDC) recorded a total of 115 incidences of oil spills in which 5,187.14 barrels of oil were spilled and 734,053 barrels of the spilt oil representing 14.2% were

recovered (SPDC Nigeria Brief, May 1995). In January 1998, 40,000 barrels of crude oil was spilled by Mobil in Eket but the largest spill in Nigeria was the offshore well blowout in January 1980 with a spill of approximately 200,000 barrels of oil into the Atlantic Ocean from an oil facility which damaged 340 hectares of mangrove forest (Nwilo and Badejo 2005b). Table below gives the number of oil spills into the marine environment reported between the periods of 1997-2001. The Niger Delta has a complex and extensive system of pipelines running across the region and large amounts of oil spill

incidences have occurred through the pipelines and storage facility failures, these failures could be caused by material defect, pipeline corrosion, ground erosion but the oil companies blame most of the spills on sabotage. The Department of Petroleum Resources contends that 88% of the oil spill incidences are traceable to equipment failure, main causes of oil spills in the Niger Delta are vandalism, oil blowouts from the flow stations, accidental and deliberate releases and oil tankers at sea (Nwilo and Badejo 2004, 2005a).

III. GAS FLARING

The Energetic Solution Conference (2004) estimates that the Niger Delta region has about 123 gas flaring sites. Agbola and Olurin (2003) stated that about 45.8 billion kilo watts of heat is discharged into the atmosphere from 1.8 billion cubic feet of gas daily in the Niger Delta region, leading to temperatures that render large areas inhabitable. Complete utilization of produced associated gas, reduction of flaring and production greenhouse gas is one of the policies that oil companies are expected to comply, with the stoppage of gas flaring completely by 2004 or 2008. Still 84.60% of total gas produced is still flared with 14.86% only being used locally (Ukoli, 2005). From 1970-1986 a total of 125.5 cubic meters approximately of gas was produced in the Niger Delta region, 102.3 (81.7%) million cubic meters was flared, 2.6 million cubic meter was used as fuel by the oil producing companies and about 14.6 million cubic meters was domestically

consumed (Awosika, 1995). In 2004 Nigerian Liquefied Natural Gas pipeline transversing through Kala-Akama, Okrika mangrove forest leaked and set ablaze and burnt for three days. The local plant and animals within the areas where engulfed (Nenibarini, 2004). Apart from this fire incidence over several decades there have been many well documented cases of fire incidences that have resulted in a large number of human fatalities.

Acid rain is another problem within the Niger Delta region caused by gas flaring which has led to loss in biodiversity, with forest and economic crops being destroyed. The dominance of grasses and shrubs in some parts of the region is indication of loss of natural forest, this may be due to acid rain but other factors maybe the cause such as agricultural activities and the exploration and exploitation of oil companies (Uyigue and Agho, 2007; Opukria and Ibaba, 2008). The concentration of acid in rain water appears to be higher in the Niger Delta region and decreases further away from the region (Uyigue and Agho, 2007). The heat generated from gas flaring kills vegetation around flaring area, destroys mangrove swamps and salt marshes, suppresses the growth and flowering of some plants, induces soil degradation and diminishes agricultural productivity (UNDP, 2006; Mba, 2000). A study by Salau (1993) and Adeyemo (2002) about the impact of gas flaring on agriculture showed a direct relationship between gas flaring and productivity decline in agriculture as shown below in Table II.

Table II : The Impact of Gas Flaring on Agricultural Output

| Distance of Farmland from Flare site | Percentage Loss in Yield of Crops |
|--------------------------------------|-----------------------------------|
| 200 meters | 100 percent |
| 600 meters | 45 percent |
| 1 Kilometer | 10 percent |

Source : Salau, 1993:19-22, Adeyemo, 2002:69.

Gas flaring is related oil spillage and UNDP, 2006 estimates that Nigeria flares 75% of the gas it produces which is more than any other country in the

world. Table III below shows some of the Global gas flaring estimates in some of the major oil producing countries.

Table III : Flaring of National Gas in Major Producing Countries (% of Gross Production in 1991)

| Country | Percentage |
|--|------------|
| United States of America(USA) | 0.6 |
| Holland | 0.0 |
| Britain | 4.3 |
| Former Union of Soviet Socialist Republic (USSR) | 1.5 |
| Mexico | 5.0 |
| OPEC Countries | |
| Nigeria | 76.0 |
| Libya | 21.0 |

| | |
|--------------|------|
| Soudi Arabia | 20.0 |
| Algeria | 19.0 |
| OPEC Total | 18.0 |
| World Total | 4.8 |

Source : World Bank Report, 1995, Vol.1:59

Apart from the above issues the toxicity to humans causing respiratory illness, leading to kidney disease, neurological disease and potential death (Ndubisi and Asia 2007). Oil exploration and exploitation activities such as this have significantly contributed to the environmental degradation of the Niger Delta region in spite of government measures to stop gas flaring by 2008 and the existence of monitoring agencies, regulations and standards, the flaring activities in the area is still a problem. Gas flaring in the area is a major source of Cox, Nox, Sox and particulate matter and the cumulative environmental impact of these flaring activities result in contaminant build up on land, shallow ground water, greenhouse effect and general global warming and have also caused high concentration of acid rain within the region.

a) Impact of Petroleum Hydrocarbon in the Environment

In terms of organisms, they vary greatly in their sensitivity to petroleum hydrocarbons and predicting the environmental impacts of specific releases of a quantity of petroleum hydrocarbon requires much site specific information about the nature of the receiving body. Most of what is known about petroleum hydrocarbons comes from studies of catastrophic oil spills. Effects tend to reflect the amount of toxic hydrocarbon in the environment and the different susceptibility of organisms, population and ecosystems and doses are rarely directly proportional to the amount released; one must consider the type of petroleum hydrocarbon released and the susceptibility of the organisms due to the environmental processes acting on the released petroleum hydrocarbon. The toxicity to the organism will depend on the available dose of petroleum available to an organism.

When petroleum hydrocarbon is released into the environment, processes alter the chemical composition of the petroleum hydrocarbon which alters the toxicity. Physical weathering may transform the petroleum hydrocarbon to a form less available to the organism. The chemical and physical properties of the petroleum hydrocarbon components determine the rate it passes into an organism. The bioavailability and persistence of specific hydrocarbons, the ability of an organism to accumulate and metabolize, fate of the metabolized products, metabolites of the hydrocarbon interphase with the normal metabolic process may alter an organisms chances of survival and reproduction in

the environment. The narcotic effects of hydrocarbon on nerve transmission are the major biological factors in determining the ecological impacts of any release; other factors include photo-degradation and photo-activation. Birds and mammals are vulnerable to oil spills when their habitats become contaminated and this may reduce reproductive rates, survival and physiological impairment (Briggs, et al., 1996). In water, oil film floating on the water surface prevents natural aeration and leads to death of fresh water or marine life and on land lead to retardation of vegetation growth, cause soil in fertility for a long period of time (Ukoli, 2005).

Ukoli, 2005 in this study made a summary of some significant pollutants from the oil industry released into the environment as follows:

1. Exploration and Production activities include: Drilling Muds, Cuttings, Oil and Greases, Salinity, Sulphides, Turbidity, Suspended Solids, Temperature, pH, Heavy metals, Biological Oxygen Demand and COD.
2. Petroleum refining activities include; Oil & Greases, BOD, COD, Phenol, Cyanide, Sulphide, Suspended solids, Toxic Additives, Hydrocarbons and Total Suspended Solids.

Effects can be either direct damage of a resource or the ability of the environment to support a resource, an effect is only said to be over when complete recovery has taken place. To quantify the effects and recovery is difficult; damage to a small area containing highly valued resources can be of greater significance than damage to a much larger area devoid of valued resources. The US DOE has reported that the Niger Delta area has experienced 4,000 oil spill incidences since 1960. This has resulted in the loss mangrove trees due to the inability of the mangrove trees to withstand the high toxicity levels of the petrochemicals spilled into the habitat. The spills have also has adverse effects on the marine habitat which has become contaminated. This poses enormous human health risk from the consumption of contaminated seafood (Twumasi and Merem 2006). The environmental problems of the Niger Delta result in generally land resource degradation, renewable resource degradation and environmental pollution, agricultural land degradation, fisheries depletion, deforestation, biodiversity loss, oil pollution, gas flaring and mangrove degradation.

b) *General Environmental Input on Oil Spill Effect on Mangroves*

Damage to mangrove forests varies with the amount and toxicity of the spilled oil product. The degree of impact is a function of oil type, spill volume, duration or re-oiling, extent of oil coverage on exposed roots, degree of substrate oiling (NRC, 2003). Light oils are acutely toxic while heavier oil can lead to eventual death by smothering. Black mangroves are the most sensitive because they osmoregulate by passing material through the roots and vascular system and then out of the leaves through specialized glands on the leaf surface. When black mangroves are oiled this osmoregulatory process aids oil uptake (Getter, et al., 1985). The Niger Delta is highly susceptible to adverse environmental changes occasioned by climate changes

because it is located in the coastal region of the world (Uyigue and Agho, 2007). The area is environmentally rich it consist of primarily mangrove swamp with areas of fresh water swamp and rainforest (Odeyemi and Ogunseitan, 1985). It has a network of streams and tributaries interlinking, the rivers are the main source of portable water for many towns and villages (Tolulope, 2004). Many species of plants and animals within the mangrove forest have become endangered or at the verge of extinction. Lewis (1981) proposed a generalized response stage of oil mangroves based on his experience at spills in Florida (T/V Howard Star) and Puerto Rico (T/V Zoe Colocotron). Lamparelli, et al., 1997 conducted a nine year research on crude oil spill site along a tidal channel in Brazil as shown below in Table IV.

Table IV : Proposed Stages of Impact and Recovery of Oiled Mangroves

| Author | Stage/Phase | Reponse |
|------------------------|-------------------|--|
| Lewis, 1981 | Acute | |
| | 0-15 days | Death of birds, turtles, fish, and invertebrates |
| | 10-30 days | Defoliation and death of small (<1m) mangroves; loss of aerial root community |
| | Chronic | |
| | 30 days-1year | Defoliation and death of medium (<3m) mangroves, tissues damage to aerial roots |
| | 1-5 years | Death of larger (>3m) mangroves; loss of oiled aerial roots and regrowth of new ones (sometimes deformed); recolonization of oiled damaged areas by new seedlings |
| Lamparelli et al, 1997 | 1-10 years | Reduction in litter fall, reduces reproduction, and reduced survival of seedlings; death or reduced growth of young trees colonizing oiled sites? Increased insect damage? |
| | 10-15 years | Complete recovery |
| | Initial Effect | |
| | 0-1 year | Seedling and saplings die; no structural alterations can be measured |
| | Structural damage | |
| | 1-4 years | High mortality is observed, and the oil impact can be measured in terms of major structural alterations |
| | Stabilization | |
| | 4-9 years | No or few additional alterations to the structural parameters; sapling growth is observed |
| | Recovery | |
| | > 9 years | It is possible to measure improvements in the structural tree parameters; ecosystem may not recover fully to its original state |

Source: NRC, 2003

Wetlands naturally have the ability to break down and assimilate pollutants, which is a resource applicable to the Niger Delta region. The forest cover within the region serves as a climate control, which regulates local climate at macro and micro levels. The canopies within the water catchment areas maintain hydrological cycles and soil stabilization, which prevents erosion and are important in watershed regulations (Chinweze and Abiola-Oloke, 2009). The clearing of vegetation within the region has many adverse effects such as: salination of water table, declination of water

quality, adds silt to the marine ecosystem due to mangrove clearing. A lot of plants found within the region are used for medicine and insect repellent. Also raw material used by many industries in Nigeria such as: starch, ink, rubber, gums, fibres are found within the region.

c) *Environmental Degradation in the Niger Delta and its Effect on the People*

In terms of environmental changes occurring within the region, large areas of mangrove forest have

been destroyed which is a major source of wood to the indigenous people. When oil spills occur, the oil spreads over a wide area affecting terrestrial and marine resources. Some past spills have necessitated the complete relocation of some communities, loss of ancestral homes, pollution of fresh water, loss of forest and agricultural land, destruction of fishing grounds and reduction of fish population, which is the major source of income for the Niger Delta people. Which all constitute massive unquantifiable losses to farmers, fishermen and hunters (Ukoli, 2005). The pollution exposes people also to new risk of diseases.

A study by Twumasi and Merem (2006) about the Niger Delta forest area made assessments using Geo spatial Data processing and Analysis; Two Landsat Thematic Mapper I and Enhanced Thematic Mapper plus (ETM+) images, the analysis was for the period 1985-2005. The results showed a slight decline in water bodies from 343,654 to 343,513 hectares, mangrove and closed forest showed a decline from an initial estimate of 55,410 hectares in 1985 to 37,117 hectares and closed forest from 250,161 hectares in 1985 to 175,609 hectares, the results of the study is shown below in Table V.

Table V: Decline of Mangrove and Closed Forest in the Niger Delta

| Classes | Area (ha) in 1985 | Area (ha) in 2000 | % Change (1985-2000) |
|-----------------------|-------------------|-------------------|----------------------|
| Water | 343,654 | 343,513 | -0.04 |
| Crop land | 16,495 | 23,974 | 45.34 |
| Settlement/bare areas | 52,738 | 108,725 | 106.16 |
| Mangrove | 55,410 | 37,117 | -33.01 |
| Closed forest | 250,161 | 175,609 | -29.80 |
| Mixed forest | 162,916 | 192,436 | 18.12 |

Source: Twumasi and Merem, 2006

IV. RESULTS

From the literature review of the oil exploration and exploitation activities in Nigeria, oil development activities have contributed to the growth and development of the country in general. There is no doubt that oil has been of great benefit to the Nigerian state and the people as a whole but the oil pollution caused by spillages from the oil industry located primarily in the Niger Delta region has caused the massive destruction to farmlands, sources of drinking water, mangrove forest, fishing grounds and declination of fish, crabs, molluscs, periwinkles and birds. Large areas of mangrove forest have been destroyed over a wide area affecting terrestrial and marine resources. Some past spills have necessitated the complete relocation of some communities, loss of ancestral homes, pollution of fresh water, loss of forest and agricultural land, destruction of fishing grounds and reduction of fish population, which is the major source of income for the Niger Delta people (Tolulope, 2004; CAB, 2009).

V. DISCUSSION

In terms of the environmental problems associated with oil exploration and exploitation in the Niger Delta. In the past based on the oil spill occurrence, a lot has happened in terms of degradation of the environment. The problem with oil is that even years after a spillage, without proper clean-up and remediation, oil persists in the environment. Even with

remediation it takes up to 15 year for recovery and factors like the geology of the area and the type of oil, will also determine the recovery period. The quantity of oil spilled is also relevant to recovery period but is not the major determinant of the recovery period and the type of clean-up method also determines recovery period. It is seen from areas like in Ogoniland that oil spills that occurred before the early 1990s with no remediation and subsequent spill from pipeline have rendered area a wasteland.

The government should be commended in that now they are tackling the problem with the UNEP with collaborations with UN (United Nation), which is taking steps towards finding a permanent solution or remediation for Ogoniland. The SPDC has to be commended also for the sponsorship of this activity and their readiness finally in taking steps into the remediation of Ogoniland.

VI. CONCLUSION

Although, the activities that come with the oil exploration and exploitation causes alterations to the environment. Which significantly have negative effects; some of the effects that come with petroleum development can be reduced or prevented basically by taking some steps in terms of prevention. Monitoring is also essential, but is lacking in the Niger Delta region. Monitoring the location of the oil companies; the terrain, the accessibility, revenue, man power availability for the monitoring agency, qualified personnel are not available. This restricts the ability and efficiency of monitoring by the government.

Environmentalists and people generally give blame to the oil companies but the Federal Government provides the laws, legislations and license, which the oil companies must adhere to. The Federal Government has to take steps, which they have started with NOSDRA, NDDC, UNEP, UN SPDC and NGOs. Improvement has begun in terms of achieving sustainable development in the Niger Delta, the government should continue to allocate more revenue into the Niger Delta for steps toward finding a permanent and lasting solution.

VII. RECOMMENDATION

- Updating and revising the legislations, reviewing the license of the oil companies and reviewing the fines will go a long way in ensuring compliance, even though the government cannot systematically or frequently monitor these sites
- Adoption of environmentally friendly technology that will minimize impacts of petroleum development on the environment; gas flaring, the gas can be converted to alcohol for diverse uses or used as an alternative energy source.
- Environmental restoration by government and oil companies
- Development of environmental management and research institutions
- Periodic Environmental Impact Statement (EIS) and Environmental Impact Assessment (EIA)
- Public awareness and education

REFERENCES RÉFÉRENCES REFERENCIAS

1. Adeyemo, A.M. (2002). The Oil Industry Extra-Ministerial Institutions and Sustainable Agricultural Development: A Case Study of Okrika L.G.A. of Rivers State, in Nigeria. *Journal of Oil and Politics*, 2(1).
2. Agbola, T. and Olurin, T.A. (2003). Landuse and Landcover Change in the Niger Delta. Excerpts from a Research Report presented to the Centre for Democracy and Development.
3. Amu, L.A.O. (1997). A Review of Nigeria's Petroleum Industry. NNPC, Lagos.
4. ANEEJ (2004). Oil of Poverty in the Niger Delta. A publication of the African Network for Environment and Economic Justice.
5. Anifowose, B. (2008). Assessing the Impact of Oil & Gas Transport on Nigeria's Environment. U21 Postgraduate Research Conference Proceedings 1, University of Birmingham UK.
6. Awosika, L.F. (1995). Impacts of global climate change and sea level rise on coastal resources and energy development in Nigeria. In: Umolu, J.C., (ed). *Global Climate Change: Impact on Energy Development*. DAMTECH Nigeria Limited, Nigeria.
7. Briggs, K.T., Yoshida, S.H. and Gershwin, M.E. (1996). The Influence of Petrochemicals and Stress on the Immune System of Seabirds. *Regulatory Toxicology and Pharmacology*, 23:145-155.
8. Chinweze, C. and Abiol-Oloke, G. (2009). Women Issues, Poverty and Social Challenges of Climate Changes in the Nigerian Niger Delta Context. 7th International Conference on the Human Dimension of Global Environmental Changes, UN Campus, Bonne, Germany.
9. CLO, Civil Liberties Organization, (2002). *Blood Trail: Repression and Resistance in the Niger Delta*, Ikeja: CLO.
10. Country Analysis Brief <http://www.eia.doe.gov/cab/index.html.pds>
11. Gabriel, A.O. (2004). Women in the Niger Delta: Environmental Issues and Challenges in the Third Millennium. www.jsd_africa.com
12. Federal Ministry of Environment Abuja, Nigerian Conservation Foundation Lagos, WWF UK and CEESP-IUCN Commission on Environmental, Economic, and Social Policy, May 31,(2006). *Niger Delta Resource Damage Assessment and Restoration Project*.
13. Getter, C.D., Ballou, T.G. and Koons, C.B., (1985). Effects of dispersed oil on mangroves: synthesis of a seven-year study. *Marine Pollution Bulletin*, 16:318-324
14. Knowles, R.S. (1983). *The First Pictorial History of the American Oil and Gas Industry 1859-1983*. Ohio University Press, Athens, OH. 169pp.
15. Kuruk, P. (2004). *Customary Water Laws and Practices: Nigeria*. <http://www.fao.org/legal/advserv/FOA/UCNCS.Nigeria.pdf>
16. Lamparelli, C.C., Rodeigues, F.O. and Orgler de Moura, D. (1997). Long-term Assessment of an Oil Spill in a Mangrove Forest in Sao Paulo, Brazil. In: Kjerfve, B., Drude de Lacerda, L., and Salif Diop, W. H. [eds.], *Mangrove Ecosystem Studies in Latin America and Africa*. UNESCO, Paris, France, pp. 191-203.
17. Mba, C.H. (2000). Environmental Protection and National Development: Towards Optimal Resolution of Conflicting Measures and Strategies. in: Aja Akpuru-Aja and Emeribe Augustine C., *Policy and Contending Issues in Nigerian National Development Strategy*. John Jacob's Publishers, Ltd, Enugu, Nigeria.
18. NEPDG, National Energy Policy Development Group, (2001). *National Energy Policy, Report of the National Energy Policy Development Group*. May 2001.

19. (NRC), National Research Council, (2003). Oil in the Sea 3: Inputs, Fates, and Effects. National Academy Press, Washington, D.C.
20. Nenibarini, Z. (2004). Impacts of Extractive Industries on the Biodiversity of the Niger Delta. National Workshop on Coastal and Marine Biodiversity Management.
21. Ndubuisi, A.L. and Asia, I.O. (2007). Environmental Pollution in Oil Producing Areas of the Niger Delta Basin, Nigeria: Empirical Assessment of Trends and People's Perception. Environmental Research Journal 1 (1-4):18-26
22. NNPC, Nigerian National Petroleum Corporation, (1984). Monthly Petroleum Information. September, Lagos, Nigeria, 53pp.
23. Nwilo, C.P. and Badejo, T.O., (2004). Management of Oil Dispersal Along the Nigerian Coastal Areas. Department of Survey & Geoinformatics, University of Lagos, Lagos, Nigeria. www.oceandocs.org/handle/1834/267
24. Nwilo, C.P. and Badejo, T.O., (2005a). Impacts and Management of Oil Spill Pollution along the Nigerian Coastal Areas. Department of Survey & Geoinformatics, University of Lagos, Lagos, Nigeria. www.fig.net/pub/figpub36/chapters/chapter_8.pdf
25. Nwilo, C.P. and Badejo, T.O., (2005b). Oil Spill Problems and Management in the Niger Delta. International Oil Spill Conference, Miami, Florida, USA.
26. Odeyemi, O. and Ogunseitan, O.A. (1985). Petroleum Industry and its Pollution Potential in Nigeria. Oil & Petroleum Pollution 2: 223-229, Elsevier Applied Science Publishers Ltd, England. 0143-7127/85.
27. Onuoha, F.C. (2008). Oil Pipeline Sabotage in Nigeria: Dimensions, Actors and Implications for National Security L/C. African Security Review, 17(3)
28. Opukri, O.C.O. and Ibaba I.S. (2008). Oil Induced Environmental Degradation and Internal Population Displacement in the Nigeria's Niger Delta. Journal of Sustainable Development in Africa, 10(1)
29. Powell, C.B., White, S.A., Ibiebele, D.O., Bara, M., Dut Kwicz, B., Isoun, M. and Oteogbu, F.U. (1985). Oshika Oil Spill Environmental Impact; effect on Aquatic biology. Paper presented at NNPC/FMHE International Seminar on petroleum industry and the Nigerian Environment 11 – 13 Nov. 1985, Kaduna, Nigeria 168 – 178.
30. Salau, A.J. (1993). Environmental Crisis and Development in Nigeria. Inaugural Lecture, No.13 University of Port Harcourt, Choba, Nigeria.
31. (SPDC), Shell Petroleum Development Company, (1995). People and the Environment. Annual Report.
32. (SPDC), Shell Petroleum Development Company, (1989). People and the Environment. Annual Report.
33. The World Bank, (1995). Defining and Environmental Development Strategy for the Niger Delta. 1:59
34. Tolulope, A.O. (2004). Oil Exploration and Environmental Degradation: the Nigerian Experience. International Information Archives, 2: 387-393, International Society for Environmental Information Science
35. Twumasi, Y. and Merem E. (2006). GIS and Remote Sensing Applications in the Assessment of Change within a Coastal Environment in the Niger Delta Region of Nigeria. International Journal of Environmental Research & Public Health, 3(1):98-106
36. Ukoli, M.K. (2005). Environmental Factors in the Management of the Oil and Gas Industry in Nigeria. www.cenbank.org
37. United Nations Development Programme, (2006:185,186). Niger Delta Human Development Report. Abuja, Nigeria.
38. Uyigüe, E. and Agho, M. (2007). Coping with Climate Change and Environmental Degradation in the Niger Delta of Southern Nigeria. Community Research and Development Centre Nigeria (CREDC).
39. Uyigüe, E. and Ogbeibu A.E. (2007). Climate Change and Poverty: Sustainable Approach in the Niger Delta Region of Nigeria. www.2007.amsterdamconference.org/....AC2007_uyigüeogbeibu.pdf
40. WRI, World Resource Institute, (1992). World Resource 1992-93, N.Y. Oxford University Press.