



GLOBAL JOURNAL OF SCIENCE FRONTIER RESEARCH
MATHEMATICS AND DECISION SCIENCES

Volume 12 Issue 6 Version 1.0 June 2012

Type : Double Blind Peer Reviewed International Research Journal

Publisher: Global Journals Inc. (USA)

Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Hypergeometric Form of Certain Indefinite Integrals

By Salahuddin

P.D.M College of Engineering, Bahadurgarh, Haryana, India

Abstract - In this paper we have given Hypergeometric form of some indefinite integrals involving Hypergeometric function. The results represent here are assume to be new.


Keywords : Pochhammer Symbol; Gaussian Hypergeometric Function; Kampé de Fériet double Hypergeometric Function and Srivastava's Triple Hypergeometric Function.

GJSFR-F Classification 2010 MSC NO : 33C05, 33C45, 33C15, 33D50, 33D60



Strictly as per the compliance and regulations of :





Hypergeometric Form of Certain Indefinite Integrals

Salahuddin

Abstract - In this paper we have given Hypergeometric form of some indefinite integrals involving Hypergeometric function. The results represent here are assume to be new.

Keywords and Phrases : Pochhammer Symbol; Gaussian Hypergeometric Function; Kampé de Fériet double Hypergeometric Function and Srivastava's Triple Hypergeometric Function.

1. INTRODUCTION AND PRELIMINARIES

The Pochhammer's symbol or Appell's symbol or shifted factorial or rising factorial or generalized factorial function is defined by

$$(b, k) = (b)_k = \frac{\Gamma(b+k)}{\Gamma(b)} = \begin{cases} b(b+1)(b+2)\cdots(b+k-1); & \text{if } k = 1, 2, 3, \dots \\ 1 & ; \text{if } k = 0 \\ k! & ; \text{if } b = 1, k = 1, 2, 3, \dots \end{cases}$$

where b is neither zero nor negative integer and the notation Γ stands for Gamma function.

Generalized Gaussian Hypergeometric Function

Generalized ordinary hypergeometric function of one variable is defined by

$${}_A F_B \left[\begin{matrix} a_1, a_2, \dots, a_A & ; \\ b_1, b_2, \dots, b_B & ; \end{matrix} \quad z \right] = \sum_{k=0}^{\infty} \frac{(a_1)_k (a_2)_k \cdots (a_A)_k z^k}{(b_1)_k (b_2)_k \cdots (b_B)_k k!}$$

or

$${}_A F_B \left[\begin{matrix} (a_A) & ; \\ (b_B) & ; \end{matrix} \quad z \right] \equiv {}_A F_B \left[\begin{matrix} (a_j)_{j=1}^A & ; \\ (b_j)_{j=1}^B & ; \end{matrix} \quad z \right] = \sum_{k=0}^{\infty} \frac{((a_A))_k z^k}{((b_B))_k k!} \quad (1.1)$$

where denominator parameters $b_1, b_2, \dots, b_B$ are neither zero nor negative integers and A, B are non-negative integers.

Kampé de Fériet's General Double Hypergeometric Function

In 1921, Appell's four double hypergeometric functions F_1, F_2, F_3, F_4 and their confluent forms $\Phi_1, \Phi_2, \Phi_3, \Psi_1, \Psi_2, \Xi_1, \Xi_2$ were unified and generalized by Kampé de Fériet.

We recall the definition of general double hypergeometric function of Kampé de Fériet in slightly modified notation of H.M.Srivastava and R.Panda:

$$F_{E;G;H}^{A:B:D} \left[\begin{matrix} (a_A):(b_B):(d_D) & ; \\ (e_E):(g_G):(h_H) & ; \end{matrix} \quad x, y \right] = \sum_{m,n=0}^{\infty} \frac{((a_A))_{m+n} ((b_B))_m ((d_D))_n x^m y^n}{((e_E))_{m+n} ((g_G))_m ((h_H))_n m! n!} \quad (1.2)$$

where for convergence

- (i) $A + B < E + G + 1$, $A + D < E + H + 1$; $|x| < \infty$, $|y| < \infty$, or
 (ii) $A + B = E + G + 1$, $A + D = E + H + 1$, and

$$\begin{cases} |x|^{\frac{1}{(A-E)}} + |y|^{\frac{1}{(A-E)}} < 1 & , \text{ if } E < A \\ \max \{|x|, |y|\} < 1 & , \text{ if } E \geq A \end{cases}$$

Srivastava's General Triple Hypergeometric Function

In 1967, H. M. Srivastava defined a general triple hypergeometric function $F^{(3)}$ in the following form

$$F^{(3)} \left[\begin{matrix} (a_A) :: (b_B); (d_D); (e_E) : (g_G); (h_H); (l_L); \\ (m_M) :: (n_N); (p_P); (q_Q) : (r_R); (s_S); (t_T); \end{matrix} \middle| x, y, z \right]$$

$$= \sum_{i,j,k=0}^{\infty} \frac{((a_A))_{i+j+k} ((b_B))_{i+j} ((d_D))_{j+k} ((e_E))_{k+i} ((g_G))_i ((h_H))_j ((l_L))_k x^i y^j z^k}{((m_M))_{i+j+k} ((n_N))_{i+j} ((p_P))_{j+k} ((q_Q))_{k+i} ((r_R))_i ((s_S))_j ((t_T))_k i! j! k!} \quad (1.3)$$

Wright's Generalized Hypergeometric Function

$${}_p\Psi_q \left[\begin{matrix} (\alpha_1, A_1), \dots, (\alpha_p, A_p) ; \\ (\lambda_1, B_1), \dots, (\lambda_q, B_q) ; \end{matrix} \middle| x \right] = \sum_{m=0}^{\infty} \frac{\Gamma(\alpha_1 + mA_1)\Gamma(\alpha_2 + mA_2) \cdots \Gamma(\alpha_p + mA_p)x^m}{\Gamma(\lambda_1 + mB_1)\Gamma(\lambda_2 + mB_2) \cdots \Gamma(\lambda_q + mB_q)m!} \quad (1.4)$$

$${}_p\Psi_q^* \left[\begin{matrix} (\alpha_1, A_1), \dots, (\alpha_p, A_p) ; \\ (\lambda_1, B_1), \dots, (\lambda_q, B_q) ; \end{matrix} \middle| x \right] = \sum_{m=0}^{\infty} \frac{(\alpha_1)_{mA_1}(\alpha_2)_{mA_2} \cdots (\alpha_p)_{mA_p}x^m}{(\lambda_1)_{mB_1}(\lambda_2)_{mB_2} \cdots (\lambda_q)_{mB_q}m!} \quad (1.5)$$

II. MAIN INTEGRALS

$$\int \frac{dx}{\sqrt{(1-x \sinh^4 x)}} =$$

$$= \cosh x \sinh^{4m-1} x (-\sinh^2 x)^{\frac{1-4m}{2}} F_{0;1}^{1;2} \left[\begin{matrix} \frac{1}{2} ; \frac{1}{2}, \frac{1-4m}{2} ; \\ - ; \frac{-3}{2} ; \end{matrix} \middle| x, \cosh^2 x \right] + Constant \quad (2.1)$$

$$\int \frac{dx}{\sqrt{(1-x \cosh^4 x)}} =$$

$$= \frac{\sqrt{-\sinh^2 x} \operatorname{cosech} x \cosh^{4m+1} x}{4m+1} F_{0;1}^{1;2} \left[\begin{matrix} \frac{1}{2} ; \frac{1}{2}, \frac{4m+1}{2} ; \\ - ; \frac{4m+3}{2} ; \end{matrix} \middle| x, \cosh^2 x \right] + Constant \quad (2.2)$$

$$\int \frac{dx}{\sqrt{(1-x \tanh^4 x)}} = \frac{\tanh^{4m+1} x}{(4m+1)} F_{0;1}^{1;2} \left[\begin{matrix} \frac{1}{2} ; 1, \frac{4m+1}{2} ; \\ - ; \frac{4m+3}{2} ; \end{matrix} \middle| x, \tanh^2 x \right] + Constant \quad (2.3)$$

$$\int \frac{dx}{\sqrt{(1-x \coth^4 x)}} = \frac{\coth^{4m+1} x}{(4m+1)} F_{0;1}^{1;2} \left[\begin{matrix} \frac{1}{2} ; 1, \frac{4m+1}{2} ; \\ - ; \frac{4m+3}{2} ; \end{matrix} ; x, \coth^2 x \right] + Constant \quad (2.4)$$

$$\int \frac{dx}{\sqrt{(1-x \operatorname{sech}^4 x)}} =$$

$$= \sinh x \cosh^2(x)^{\frac{4m+1}{2}} \operatorname{sech}^{4m+1} x F_{0;1}^{1;2} \left[\begin{matrix} \frac{1}{2} ; \frac{1}{2}, \frac{1+4m}{2} ; \\ - ; \frac{3}{2} ; \end{matrix} ; x, -\sinh^2 x \right] + Constant \quad (2.5)$$

$$\int \frac{dx}{\sqrt{(1-x \operatorname{cosech}^4 x)}} =$$

$$= \cosh x (-\sinh^2(x))^{\frac{4m+1}{2}} \operatorname{cosech}^{4m+1} x F_{0;1}^{1;2} \left[\begin{matrix} \frac{1}{2} ; \frac{1}{2}, \frac{1+4m}{2} ; \\ - ; \frac{3}{2} ; \end{matrix} ; x, \cosh^2 x \right] + Constant \quad (2.6)$$

III. DERIVATION OF INTEGRALS

Derivation of integral (2.1)

$$\int \frac{dx}{\sqrt{(1-x \sinh^4 x)}} = \int (1-x \sinh^4 x)^{-\frac{1}{2}} dx$$

$$= \sum_{m=0}^{\infty} \frac{(\frac{1}{2})_m (x)^m}{m!} \sinh^{4m} x \, dx = \sum_{m=0}^{\infty} \frac{(\frac{1}{2})_m (x)^m}{m!} \int \sinh^{4m} x \, dx$$

$$= \sum_{m=0}^{\infty} \frac{(\frac{1}{2})_m (x)^m}{m!} (\cosh x) \sinh^{4m-1} x (-\sinh^2 x)^{\frac{1-4m}{2}} {}_2F_1 \left[\begin{matrix} \frac{1}{2}, \frac{1-4m}{2} ; \\ \frac{-3}{2} ; \end{matrix} ; \cosh^2 x \right] + Constant$$

$$= \cosh x \sinh^{4m-1} x (-\sinh^2 x)^{\frac{1-4m}{2}} F_{0;1}^{1;2} \left[\begin{matrix} \frac{1}{2} ; \frac{1}{2}, \frac{1-4m}{2} ; \\ - ; \frac{-3}{2} ; \end{matrix} ; x, \cosh^2 x \right] + Constant \quad (3.1)$$

Derivation of integral (2.2)

$$\int \frac{dx}{\sqrt{(1-x \cosh^4 x)}} = \int (1-x \cosh^4 x)^{-\frac{1}{2}} dx$$

$$= \sum_{m=0}^{\infty} \frac{(\frac{1}{2})_m (x)^m}{m!} \cosh^{4m} x \, dx = \sum_{m=0}^{\infty} \frac{(\frac{1}{2})_m (x)^m}{m!} \int \cosh^{4m} x \, dx$$

$$= \sum_{m=0}^{\infty} \frac{(\frac{1}{2})_m (x)^m}{m!} \frac{\sqrt{-\sinh^2 x} \cosh^{4m+1} x \operatorname{cosech} x}{(4m+1)} {}_2F_1 \left[\begin{matrix} \frac{1}{2}, \frac{4m+1}{2} ; \\ \frac{4m+3}{2} ; \end{matrix} ; \cosh^2 x \right] + Constant$$

$$= \frac{\sqrt{-\sinh^2 x} \operatorname{cosech} x \cosh^{4m+1} x}{4m+1} F_{0;1}^{1;2} \left[\begin{matrix} \frac{1}{2} ; \frac{1}{2}, \frac{4m+1}{2} ; \\ - ; \frac{4m+3}{2} ; \end{matrix} ; x, \cosh^2 x \right] + Constant \quad (3.2)$$

Derivation of integral (2.3)

$$\begin{aligned}
 \int \frac{dx}{\sqrt{(1-x \tanh^4 x)}} &= \int (1-x \tanh^4 x)^{-\frac{1}{2}} dx \\
 \int \sum_{m=0}^{\infty} \frac{(\frac{1}{2})_m (x)^m}{m!} \tanh^{4m} x \, dx &= \sum_{m=0}^{\infty} \frac{(\frac{1}{2})_m (x)^m}{m!} \int \tanh^{4m} x \, dx \\
 &= \sum_{m=0}^{\infty} \frac{(\frac{1}{2})_m (x)^m}{m!} \frac{\tanh^{4m+1} x}{(4m+1)} {}_2F_1 \left[\begin{matrix} 1, \frac{4m+1}{2} \\ \frac{4m+3}{2} \end{matrix} ; \tanh^2 x \right] + Constant \\
 &= \frac{\tanh^{4m+1} x}{(4m+1)} F_{0;1}^{1;2} \left[\begin{matrix} \frac{1}{2} ; 1, \frac{4m+1}{2} \\ - ; \frac{4m+3}{2} \end{matrix} ; x, \tanh^2 x \right] + Constant \quad (3.3)
 \end{aligned}$$

Derivation of integral (2.4)

$$\begin{aligned}
 \int \frac{dx}{\sqrt{(1-x \coth^4 x)}} &= \int (1-x \coth^4 x)^{-\frac{1}{2}} dx \\
 \int \sum_{m=0}^{\infty} \frac{(\frac{1}{2})_m (x)^m}{m!} \coth^{4m} x \, dx &= \sum_{m=0}^{\infty} \frac{(\frac{1}{2})_m (x)^m}{m!} \int \coth^{4m} x \, dx \\
 &= \sum_{m=0}^{\infty} \frac{(\frac{1}{2})_m (x)^m}{m!} \frac{\coth^{4m+1} x}{(4m+1)} {}_2F_1 \left[\begin{matrix} 1, \frac{4m+1}{2} \\ \frac{4m+3}{2} \end{matrix} ; \coth^2 x \right] + Constant \\
 &= \frac{\coth^{4m+1} x}{(4m+1)} F_{0;1}^{1;2} \left[\begin{matrix} \frac{1}{2} ; 1, \frac{4m+1}{2} \\ - ; \frac{4m+3}{2} \end{matrix} ; x, \coth^2 x \right] + Constant \quad (3.4)
 \end{aligned}$$

Derivation of integral (2.5)

$$\begin{aligned}
 \int \frac{dx}{\sqrt{(1-x \operatorname{sech}^4 x)}} &= \int (1-x \operatorname{sech}^4 x)^{-\frac{1}{2}} dx \\
 \int \sum_{m=0}^{\infty} \frac{(\frac{1}{2})_m (x)^m}{m!} \operatorname{sech}^{4m} x \, dx &= \sum_{m=0}^{\infty} \frac{(\frac{1}{2})_m (x)^m}{m!} \int \operatorname{sech}^{4m} x \, dx \\
 &= \sum_{m=0}^{\infty} \frac{(\frac{1}{2})_m (x)^m}{m!} \sinh x \cosh^2(x)^{\frac{4m+1}{2}} \operatorname{sech}^{4m+1} x {}_2F_1 \left[\begin{matrix} \frac{1}{2}, \frac{4m+1}{2} \\ \frac{3}{2} \end{matrix} ; -\sinh^2 x \right] + Constant \\
 &= \sinh x \cosh^2(x)^{\frac{4m+1}{2}} \operatorname{sech}^{4m+1} x F_{0;1}^{1;2} \left[\begin{matrix} \frac{1}{2} ; \frac{1}{2}, \frac{1+4m}{2} \\ - ; \frac{3}{2} \end{matrix} ; x, -\sinh^2 x \right] + Constant \quad (3.5)
 \end{aligned}$$

Derivation of integral (2.6)

$$\begin{aligned}
 \int \frac{dx}{\sqrt{(1-x \operatorname{cosech}^4 x)}} &= \int (1-x \operatorname{cosech}^4 x)^{-\frac{1}{2}} dx \\
 \int \sum_{m=0}^{\infty} \frac{(\frac{1}{2})_m (x)^m}{m!} \operatorname{cosech}^{4m} x \, dx &= \sum_{m=0}^{\infty} \frac{(\frac{1}{2})_m (x)^m}{m!} \int \operatorname{cosech}^{4m} x \, dx
 \end{aligned}$$

$$\begin{aligned}
&= \sum_{m=0}^{\infty} \frac{(\frac{1}{2})_m (-x)^m}{m!} \cosh x (-\sinh^2(x))^{\frac{4m+1}{2}} \operatorname{cosech}^{4m+1} x {}_2F_1 \left[\begin{matrix} \frac{1}{2}, \frac{4m+1}{2} \\ \frac{3}{2} \end{matrix} ; \cosh^2 x \right] + \text{Constant} \\
&= \cosh x (-\sinh^2(x))^{\frac{4m+1}{2}} \operatorname{cosech}^{4m+1} x F_{0;1}^{1;2} \left[\begin{matrix} \frac{1}{2} ; \frac{1}{2}, \frac{1+4m}{2} \\ - ; \frac{3}{2} \end{matrix} ; x, \cosh^2 x \right] + \text{Constant} \quad (3.6)
\end{aligned}$$

IV. CONCLUSION

In our work we have established hypergeometric form of some indefinite integrals. However, the formulae presented herein may be further developed to extend the work. Thus we can only hope that the development presented in this work will stimulate further interest and research in this important area of classical special functions. Just as the mathematical properties of the Gauss hypergeometric function are already of immense and significant utility in mathematical sciences and numerous other areas of pure and applied mathematics, the elucidation and discovery of the formula of hypergeometric functions considered herein should certainly eventually prove useful to further developments in the broad areas alluded to above.

REFERENCES RÉFÉRENCES REFERENCIAS

1. Agarwal, R. P.; *Resonance of Ramanujan's Mathematics*. Vol. I, New Age International (P) Ltd., New Delhi, 1996.
2. Agarwal, R. P.; *Resonance of Ramanujan's Mathematics*. Vol. II, New Age International (P) Ltd., New Delhi, 1996.
3. Agarwal, R. P.; *Resonance of Ramanujan's Mathematics*. Vol. III, New Age International (P) Ltd., New Delhi, 1999.
4. Berndt, B. C.; *Ramanujan's Notebooks*. Part I, Springer-Verlag, New York, 1985.
5. Berndt, B. C.; *Ramanujan's Notebooks*. Part II, Springer-Verlag, New York, 1989.
6. Berndt, B. C.; *Ramanujan's Notebooks*. Part III, Springer-Verlag, New York, 1991.
7. Berndt, B. C.; *Ramanujan's Notebooks*. Part IV, Springer-Verlag, New York, 1994.
8. Berndt, B. C.; *Ramanujan's Notebooks*. Part V, Springer-Verlag, New York, 1998.
9. Srivastava, H.M. and Daoust, M.C. ; A note on the convergence of Kampje de Fjeriet double hypergeometric series. *Math.Nachr.* 53(1972).151-159.
10. Qureshi, M.I., Salahuddin, Chaudhary, M. P. and Quraishi, K.A.; Evaluation of Certain Elliptic Type Single, Double Integrals of Ramanujan and Erdjelyi, *J. Mathematics Research.*, 2(2010), 148-156.



This page is intentionally left blank