



GLOBAL JOURNAL OF SCIENCE FRONTIER RESEARCH
BIOLOGICAL SCIENCES

Volume 13 Issue 1 Version 1.0 Year 2013

Type : Double Blind Peer Reviewed International Research Journal

Publisher: Global Journals Inc. (USA)

Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Leaching of Bifenthrin in two Different Types of Soil under Laboratory Condition

By Sachin Kumari, Ramprakash & Beena Kumari

CCS Haryana Agricultural University

Abstract - Leaching studies of bifenthrin in two different soils were carried out under laboratory conditions during October- November 2011. Commercial formulation of bifenthrin available as Talstar 10 % EC were applied at two treatment levels i.e. 25 and 50 μg as single and double dose, respectively. More than 90 per cent of bifenthrin were retained in 5 cm soil section in sandy loam and clay loam soil which shows the immobility of this pyrethroid insecticide in soil. No residues were detected in the leachate fractions of the insecticide.

Keywords : *bifenthrin. residues. persistence. soil.*

GJSFR-C Classification : *FOR Code: 300102*



Strictly as per the compliance and regulations of :



RESEARCH | DIVERSITY | ETHICS

© 2013. Sachin Kumari, Ramprakash & Beena Kumari. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License <http://creativecommons.org/licenses/by-nc/3.0/>, permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Leaching of Bifenthrin in two Different Types of Soil under Laboratory Condition

Sachin Kumari ^α, Ramprakash ^σ & Beena Kumari ^ρ

Abstract - Leaching studies of bifenthrin in two different soils were carried out under laboratory conditions during October-November 2011. Commercial formulation of bifenthrin available as Talstar 10 % EC were applied at two treatment levels i.e. 25 and 50 µg as single and double dose, respectively. More than 90 per cent of bifenthrin were retained in 5 cm soil section in sandy loam and clay loam soil which shows the immobility of this pyrethroid insecticide in soil. No residues were detected in the leachate fractions of the insecticide

Keywords : bifenthrin. residues. persistence. soil.

I. INTRODUCTION

Over use of pesticide is the contamination of soil and water sources including the aquatic system. Several scientists reported effect of pesticides in soil (Frampton et al. 2006; Coupe et al. 2005; Willian and Mueller 1994; Balasubramanya and Patil 1980). Bifenthrin (2-Methyl-3-phenyl phenyl methyl (1*S*, 3*S*)-3-(*Z*)-2-chloro-3, 3, 3-trifluoroprop-1-enyl 2, 2-dimethylcyclopropane-1-carboxylate; Fig. 1) is a third generation pyrethroid. It is used in agriculture, public health and forestry. Pesticides may have potential to bind to soil, the extent of which depends greatly on the nature of the chemical used. Therefore it has been observed that the physico-chemical nature of the soil is important for the persistence, metabolism and binding of pesticides in soil. They include sorption to soil constituents, movement of the soil in runoff, movement into the air by volatilization, movement downward in the soil by leaching, and movement upward in the soil through the soil by leaching, and movement upward in the soil through capillary forces. The main processes potentially affecting the ultimate fate of pesticides in soil are retention by soil materials (involving adsorption/desorption processes), transformation processes (biological and chemical degradation), and transport (through soil, atmosphere, surface water, or ground water) (Saltzman and Yaron 1986; van der Hoff and van Zoonen 1999).

The present study was undertaken to determine the leaching of bifenthrin in two different soil at different doses under laboratory condition.




Figure 1 : Chemical structure of bifenthrin

II. MATERIALS AND METHODS

Leaching studies were carried out in sandy loam and clay loam soil. Soil were collected from Haryana. The physico- chemical properties of the soil are presented in Table 1 and 2.

For laboratory experiment, plexi glass columns (90 cm x 5 cm i.d) fitted with perforated plexi glass sieve covered with filter paper (Whatmann No.1) at the bottom were used. Each column was filled with soil up to 60 cm height to a uniform bulk density of 1.35 (sandy loam) and 1.40 (clay loam) g cm⁻³ ± 0.1. Weighed amount of different soils 128 g (sandy loam) and 137 g (clay loam) was poured in the different columns each time with the help of a funnel (1 cm i.d) tapped gently from a fixed height and pressed uniformly by a wooden roller. The soil in each column was covered with a filter paper which was topped by a glass wool swab. Soil column so filled were installed vertically on wooden stand with their bottom resting on the fixed perforated sieve, so as to facilitate collection of effluents during leaching.

Leaching studies were carried out in triplicates. Soil (5 g) was fortified with 25 µg (single dose) and 50 µg (double dose) solution of bifenthrin. Added this fortified soil at the top of the column. Cotton swabs prewashed with acetone dried and were placed on top of soil to avoid disturbance of soil. One column packed with soil was kept as control to which no pesticide was added.

Each time 50 mm lot of leaching water (98 ml) was added with the help of a pipette, with the care that no spattering of soil took place. The leaching with the next 50 mm lot of distilled water was done after 24 hrs. After the disappearance of the standing water in the column, leaching was repeated till each leaching cycle of 200, 400 and 600 mm depth was completed. During leaching, the columns were covered with perforated polythene bags to minimize the loss of leaching water due to evaporation. During leaching 2-3 drops of toluene solution was added to each column to check microbial growth in it.

Author ^α : Department of Chemistry & Physics Department of Soil science.

Author ^σ : Department of Entomology CCS Haryana Agricultural University, Hisar 125 004, India. E-mail : beena@hau.ernet.in

Leachate oozing out of all the column were collected and processed for the presence of bifenthrin. Total three fraction of leachates were collected. Collect 8 ml (sandy loam) and 5 ml (clay loam) of each fraction. After completion of leaching, intact soil cores were taken out of the plexi glass column. Leachate samples were processed as per method of Kumari et al. (2008).

Leachate sample was taken in 250ml separatory funnel, to which 5g sodium chloride was added.

The solution was partitioned with 15% dichloromethane (DCM) in hexane thrice (30, 20, 10 ml) by shaking vigorously for 1 minute every time. The organic layers were collected and passed through 2-3 cm pad of anhydrous sodium sulphate. The organic layer was evaporated to near dryness on rotary evaporator followed by gas manifold evaporator. The residues were redissolved in n- hexane to make final volume to 2 ml for GC analysis.

Extraction and clean-up of soil sample was carried out as per method of Kumari et al. (2008). Ground, sieved and dry representative (15 g) of soil mixed with charcoal and florisil (0.3 g each) and 10 g of anhydrous sodium sulphate. The mixture was packed compactly in a glass column (60 cm × 22 mm) in between two layers of anhydrous sodium sulphate. Residues were eluted with 100 mL of hexane:acetone (9:1 v/v) at flow rate of 2–3 mL/min. The eluate was concentrated on flash evaporator and made the final volume to 2 mL for GC analysis. Estimation of micro quantities of bifenthrin residues in leachate as well as in soil samples were carried out by GC-ECD system equipped with capillary column.

III. RESULTS AND DISCUSSION

a) Sandy Loam Soil

The data on leaching potential of bifenthrin in sandy loam soil are presented in Table 3 and Fig.2. Total amount of bifenthrin recovered out of 25 and 50 µg were 92 and 95.00 per cent from single and double dose, respectively. Beyond the plough depth (15 cm) of the soil, the amount of bifenthrin could not be detected in any of the doses. None of the leachate fractions from both the treatments showed the presence of bifenthrin residues.

b) Clay Loam Soil

The data on leaching potential of bifenthrin in clay loam soil are presented in Table 4 and Fig.3. Total amount of bifenthrin recovered out of 25 and 50 µg were 96.40 and 98.50 per cent from single and double dose, respectively. Beyond 10 cm depth of soil, the amount of bifenthrin could not be detected in any of the doses. None of the leachate fractions from both the treatments showed the presence of bifenthrin residues.

Similar results were observed by Chauhan et al. (2011) for leaching behaviour of bifenthrin and lambda-

cyhalothrin in sandy loam soil. More than 90 % of bifenthrin as well as lambda-cyhalothrin were retained in 5 cm soil section, which shows their immobility in soil. Both the insecticides restricted up to ploughed soil (15 cm depth) in both the doses.

Manoj and Gajbhiye (2007) reported low mobility of bifenthrin in soil. Its residues remained within top 15 cm and more than 99 percent of residues were recovered from top 0-10 cm layer. Tariq *et al*, (2006) reported highest concentrations of lambda-cyhalothrin were found in the top 0–10 cm layer.

ACKNOWLEDGMENT

The authors wish to express their gratitude to the Head, Department of Entomology for providing research facilities.

REFERENCES RÉFÉRENCES REFERENCIAS

1. BalasubramanyaRH, PatilRB (1980) Degradation of carboxin and oxycarboxin in different soils. *Plant Soil* 57:195–201.
2. Chauhan Reena, Indu Chopra. Beena Kumari (2011). Leaching behaviour of bifenthrin and lambda-cyhalothrin in sandy loam soil. *Global Journal of Science Frontier Research*. Vol.12(6)21–24.
3. Coupe R, Welch H, Pell A, Thurman E (2005) Herbicide and degradate flux in the Yazoo River Basin. *Int J Environ Anal Chem* 85:1127–1140.
4. Frampton GK, Jnsch S, Scott-Fordsmand JJ, Römbke J, van den Brink PJ (2006) Effects of pesticides on soil invertebrates in laboratory studies: a review and analysis using species sensitivity distributions. *Environ Toxicol Chem* 25:2480–2489.
5. Kumari B, Madan VK, Kathpal TS (2008) Status of insecticide contamination of soil and water in Haryana, India. *Environ Monit Assess* 136:239–244.
6. Manoj VB and Gajbhiye VT. 2007. Adsorption – Desorption and leaching of bifenthrin in soil. *Pestic. Res. J.* 19(2):257–261.
7. Saltzman S, Yaron B (1986) *Pesticides in soils*. Van Nostrand Reinhold, New York.
8. Sharma D, Dubey JK, Patyal SK (2006) Persistence of fenazaquin in apple fruits and soil. *Pesticide Res J* 18(1):79–81.
9. Tariq MY, Afzal S and Hussain I. 2006. Degradation and persistence of cotton pesticides in sandy loame soils from Punjab. *Pakistan. Environ. Res.* 100:184–196
10. van der Hoff GR, van Zoonen P (1999) Trace analysis of pesticides by gas chromatography. *J Chromatogr A* 843:301–322.
11. WillianWT, MuellerTC (1994) Liquid chromatographic determination of norflurazon and its initial metabolite in soil. *JAOAC Int* 77:752–755.

Table 1 : Physico-chemical characteristics of sandy loam soil

Soil Property				
Texture	pH	EC _e (dSm ⁻¹)	O.C. (%)	P ₂ O ₅ (kg ha ⁻¹)
Sandy Loam	7.6	2.0	0.32	33

Table 2 : Physico-chemical characteristics of clay loam soil

Soil Property						
Texture	pH	EC _e (dSm ⁻¹)	O.C. (%)	N (kg ha ⁻¹)	P (kg ha ⁻¹)	K (kg ha ⁻¹)
Clayey loam	8.9	0.4	0.36	180	18	140

Table 3 : Leaching behaviour of bifenthrin in sandy loam soil


Soil Column Depth (cm)	Residues (μg)*			
	Single dose (25 μg) ±SD	% Retention	Double dose (50 μg) ±SD	% Retention
0-5	19.75±1.1.9	85.86	43.00±3.05	90.52
5-10	2.50±2.30	10.86	3.25±1.64	6.84
10-15	0.75±3.03	3.26	1.25±2.04	2.63

*Average residues of three replicates
Leachate contained no residues

Table 4 : Leaching behaviour of bifenthrin in clay loam soil

Soil Column Depth (cm)	Residues (μg)*			
	Single dose (25 μg) ±SD	% Retention	Double dose (50 μg) ±SD	% Retention
0-5	22.00±2.52	91.28	45.55±3.07	92.48
5-10	2.10±3.41	8.71	3.70±1.11	7.51

*Average residues of three replicates
Leachate contained no residues

**Figure 2 :** Distribution of bifenthrin at different soil depths for two doses in sandy loam soil




Figure 3 : Distribution of bifenthrin at different soil depths for two doses in clay loam soil