

GLOBAL JOURNAL

OF SCIENCE FRONTIER RESEARCH: D

Agriculture and Veterinary



Coastal Fisheries Development

Formaldehyde Treated Sesame

Highlights

Agricultural Insurance Scheme

Performance of Broiler Chicken

Discovering Thoughts, Inventing Future

VOLUME 13

ISSUE 11

VERSION 1.0



GLOBAL JOURNAL OF SCIENCE FRONTIER RESEARCH: D
AGRICULTURE & VETERINARY



GLOBAL JOURNAL OF SCIENCE FRONTIER RESEARCH: D
AGRICULTURE & VETERINARY
VOLUME 13 ISSUE 11 (VER. 1.0)

OPEN ASSOCIATION OF RESEARCH SOCIETY

© Global Journal of Science
Frontier Research .2013.

All rights reserved.

This is a special issue published in version 1.0
of "Global Journal of Science Frontier
Research." By Global Journals Inc.

All articles are open access articles distributed
under "Global Journal of Science Frontier
Research"

Reading License, which permits restricted use.
Entire contents are copyright by of "Global
Journal of Science Frontier Research" unless
otherwise noted on specific articles.

No part of this publication may be reproduced
or transmitted in any form or by any means,
electronic or mechanical, including
photocopy, recording, or any information
storage and retrieval system, without written
permission.

The opinions and statements made in this
book are those of the authors concerned.
Ultraculture has not verified and neither
confirms nor denies any of the foregoing and
no warranty or fitness is implied.

Engage with the contents herein at your own
risk.

The use of this journal, and the terms and
conditions for our providing information, is
governed by our Disclaimer, Terms and
Conditions and Privacy Policy given on our
website [http://globaljournals.us/terms-and-condition/
menu-1463/](http://globaljournals.us/terms-and-condition/menu-1463/)

By referring / using / reading / any type of
association / referencing this journal, this
signifies and you acknowledge that you have
read them and that you accept and will be
bound by the terms thereof.

All information, journals, this journal,
activities undertaken, materials, services and
our website, terms and conditions, privacy
policy, and this journal is subject to change
anytime without any prior notice.

Incorporation No.: 0423089
License No.: 42125/022010/1186
Registration No.: 430374
Import-Export Code: 1109007027
Employer Identification Number (EIN):
USA Tax ID: 98-0673427

Global Journals Inc.

(A Delaware USA Incorporation with "Good Standing"; Reg. Number: 0423089)

Sponsors: *Open Association of Research Society*
Open Scientific Standards

Publisher's Headquarters office

Global Journals Inc., Headquarters Corporate Office,
Cambridge Office Center, II Canal Park, Floor No.
5th, **Cambridge (Massachusetts)**, Pin: MA 02141
United States

USA Toll Free: +001-888-839-7392

USA Toll Free Fax: +001-888-839-7392

Offset Typesetting

Open Association of Research Society, Marsh Road,
Rainham, Essex, London RM13 8EU
United Kingdom.

Packaging & Continental Dispatching

Global Journals, India

Find a correspondence nodal officer near you

To find nodal officer of your country, please
email us at local@globaljournals.org

eContacts

Press Inquiries: press@globaljournals.org

Investor Inquiries: investers@globaljournals.org

Technical Support: technology@globaljournals.org

Media & Releases: media@globaljournals.org

Pricing (Including by Air Parcel Charges):

For Authors:

22 USD (B/W) & 50 USD (Color)

Yearly Subscription (Personal & Institutional):

200 USD (B/W) & 250 USD (Color)

EDITORIAL BOARD MEMBERS (HON.)

John A. Hamilton, "Drew" Jr.,
Ph.D., Professor, Management
Computer Science and Software
Engineering
Director, Information Assurance
Laboratory
Auburn University

Dr. Henry Hexmoor
IEEE senior member since 2004
Ph.D. Computer Science, University at
Buffalo
Department of Computer Science
Southern Illinois University at Carbondale

Dr. Osman Balci, Professor
Department of Computer Science
Virginia Tech, Virginia University
Ph.D. and M.S. Syracuse University,
Syracuse, New York
M.S. and B.S. Bogazici University,
Istanbul, Turkey

Yogita Bajpai
M.Sc. (Computer Science), FICCT
U.S.A. Email:
yogita@computerresearch.org

Dr. T. David A. Forbes
Associate Professor and Range
Nutritionist
Ph.D. Edinburgh University - Animal
Nutrition
M.S. Aberdeen University - Animal
Nutrition
B.A. University of Dublin- Zoology

Dr. Wenying Feng
Professor, Department of Computing &
Information Systems
Department of Mathematics
Trent University, Peterborough,
ON Canada K9J 7B8

Dr. Thomas Wischgoll
Computer Science and Engineering,
Wright State University, Dayton, Ohio
B.S., M.S., Ph.D.
(University of Kaiserslautern)

Dr. Abdurrahman Arslanyilmaz
Computer Science & Information Systems
Department
Youngstown State University
Ph.D., Texas A&M University
University of Missouri, Columbia
Gazi University, Turkey

Dr. Xiaohong He
Professor of International Business
University of Quinnipiac
BS, Jilin Institute of Technology; MA, MS,
PhD,. (University of Texas-Dallas)

Burcin Becerik-Gerber
University of Southern California
Ph.D. in Civil Engineering
DDes from Harvard University
M.S. from University of California, Berkeley
& Istanbul University

Dr. Bart Lambrecht

Director of Research in Accounting and Finance
Professor of Finance
Lancaster University Management School
BA (Antwerp); MPhil, MA, PhD
(Cambridge)

Dr. Carlos García Pont

Associate Professor of Marketing
IESE Business School, University of Navarra
Doctor of Philosophy (Management),
Massachusetts Institute of Technology (MIT)
Master in Business Administration, IESE,
University of Navarra
Degree in Industrial Engineering,
Universitat Politècnica de Catalunya

Dr. Fotini Labropulu

Mathematics - Luther College
University of Regina
Ph.D., M.Sc. in Mathematics
B.A. (Honors) in Mathematics
University of Windsor

Dr. Lynn Lim

Reader in Business and Marketing
Roehampton University, London
BCom, PGDip, MBA (Distinction), PhD,
FHEA

Dr. Mihaly Mezei

ASSOCIATE PROFESSOR
Department of Structural and Chemical
Biology, Mount Sinai School of Medical
Center
Ph.D., Eötvös Loránd University
Postdoctoral Training,
New York University

Dr. Söhnke M. Bartram

Department of Accounting and Finance
Lancaster University Management School
Ph.D. (WHU Koblenz)
MBA/BBA (University of Saarbrücken)

Dr. Miguel Angel Ariño

Professor of Decision Sciences
IESE Business School
Barcelona, Spain (Universidad de Navarra)
CEIBS (China Europe International Business School).
Beijing, Shanghai and Shenzhen
Ph.D. in Mathematics
University of Barcelona
BA in Mathematics (Licenciatura)
University of Barcelona

Philip G. Moscoso

Technology and Operations Management
IESE Business School, University of Navarra
Ph.D in Industrial Engineering and
Management, ETH Zurich
M.Sc. in Chemical Engineering, ETH Zurich

Dr. Sanjay Dixit, M.D.

Director, EP Laboratories, Philadelphia VA
Medical Center
Cardiovascular Medicine - Cardiac
Arrhythmia
Univ of Penn School of Medicine

Dr. Han-Xiang Deng

MD., Ph.D
Associate Professor and Research
Department Division of Neuromuscular
Medicine
Department of Neurology and Clinical
Neuroscience
Northwestern University
Feinberg School of Medicine

Dr. Pina C. Sanelli

Associate Professor of Public Health
Weill Cornell Medical College
Associate Attending Radiologist
NewYork-Presbyterian Hospital
MRI, MRA, CT, and CTA
Neuroradiology and Diagnostic
Radiology
M.D., State University of New York at
Buffalo, School of Medicine and
Biomedical Sciences

Dr. Roberto Sanchez

Associate Professor
Department of Structural and Chemical
Biology
Mount Sinai School of Medicine
Ph.D., The Rockefeller University

Dr. Wen-Yih Sun

Professor of Earth and Atmospheric
SciencesPurdue University Director
National Center for Typhoon and
Flooding Research, Taiwan
University Chair Professor
Department of Atmospheric Sciences,
National Central University, Chung-Li,
TaiwanUniversity Chair Professor
Institute of Environmental Engineering,
National Chiao Tung University, Hsin-
chu, Taiwan.Ph.D., MS The University of
Chicago, Geophysical Sciences
BS National Taiwan University,
Atmospheric Sciences
Associate Professor of Radiology

Dr. Michael R. Rudnick

M.D., FACP
Associate Professor of Medicine
Chief, Renal Electrolyte and
Hypertension Division (PMC)
Penn Medicine, University of
Pennsylvania
Presbyterian Medical Center,
Philadelphia
Nephrology and Internal Medicine
Certified by the American Board of
Internal Medicine

Dr. Bassey Benjamin Esu

B.Sc. Marketing; MBA Marketing; Ph.D
Marketing
Lecturer, Department of Marketing,
University of Calabar
Tourism Consultant, Cross River State
Tourism Development Department
Co-ordinator , Sustainable Tourism
Initiative, Calabar, Nigeria

Dr. Aziz M. Barbar, Ph.D.

IEEE Senior Member
Chairperson, Department of Computer
Science
AUST - American University of Science &
Technology
Alfred Naccash Avenue – Ashrafieh

PRESIDENT EDITOR (HON.)

Dr. George Perry, (Neuroscientist)

Dean and Professor, College of Sciences

Denham Harman Research Award (American Aging Association)

ISI Highly Cited Researcher, Iberoamerican Molecular Biology Organization

AAAS Fellow, Correspondent Member of Spanish Royal Academy of Sciences

University of Texas at San Antonio

Postdoctoral Fellow (Department of Cell Biology)

Baylor College of Medicine

Houston, Texas, United States

CHIEF AUTHOR (HON.)

Dr. R.K. Dixit

M.Sc., Ph.D., FICCT

Chief Author, India

Email: authorind@computerresearch.org

DEAN & EDITOR-IN-CHIEF (HON.)

Vivek Dubey(HON.)

MS (Industrial Engineering),

MS (Mechanical Engineering)

University of Wisconsin, FICCT

Editor-in-Chief, USA

editorusa@computerresearch.org

Sangita Dixit

M.Sc., FICCT

Dean & Chancellor (Asia Pacific)

deanind@computerresearch.org

Suyash Dixit

(B.E., Computer Science Engineering), FICCTT

President, Web Administration and

Development , CEO at IOSRD

COO at GAOR & OSS

Er. Suyog Dixit

(M. Tech), BE (HONS. in CSE), FICCT

SAP Certified Consultant

CEO at IOSRD, GAOR & OSS

Technical Dean, Global Journals Inc. (US)

Website: www.suyogdixit.com

Email: suyog@suyogdixit.com

Pritesh Rajvaidya

(MS) Computer Science Department

California State University

BE (Computer Science), FICCT

Technical Dean, USA

Email: pritesh@computerresearch.org

Luis Galárraga

J!Research Project Leader

Saarbrücken, Germany

CONTENTS OF THE VOLUME

- i. Copyright Notice
- ii. Editorial Board Members
- iii. Chief Author and Dean
- iv. Table of Contents
- v. From the Chief Editor's Desk
- vi. Research and Review Papers
 1. Effect of Formaldehyde Treated Sesame Cake Feeding on Growth Performnce of Growing Female Goats in Fodder based Basal Diet. **1-5**
 2. Value-Chain Analysis for Coastal Fisheries Development in Nigeria. **7-15**
 3. Environmental Conservation Information Needs of Farmers in Owerri West Area of Imo State, Nigeria. **17-23**
 4. Evaluation of Quality Protein Maize (QPM) and Normal Mazie for Growth Performance of Broiler Chicken in Nepal. **25-31**
 5. Effect of Heat Treated Soybean Cake Feeding on Growth Performnce of Growing Female Goats in Fodder based Basal Diet. **33-38**
 6. Analysis of Rural Cassava Farmers' Participation in the Nigeria Agricultural Insurance Scheme in Imo State, Nigeria. **39-45**
 7. Livelihood Vulnerability to Climate Change based on Agro Ecological Regions of Nepal. **47-53**
- vii. Auxiliary Memberships
- viii. Process of Submission of Research Paper
- ix. Preferred Author Guidelines
- x. Index



Effect of Formaldehyde Treated Sesame Cake Feeding on Growth Performance of Growing Female Goats in Fodder based Basal Diet

By M.R. Tiwari, D. Adhikari, D.P. Adhikari, R.P. Ghimire & S.H. Ghimire

Nepal Agricultural Research Council, Nepal

Abstract - An experiment was carried out on fifteen growing female goats (50% Barberi 6, Kiko 6 and Khari 6) at the Agriculture Research Station (Goat), Bandipur for 90 days. Female goats of an average six months age and of body weight 7.82 kg were allocated into three groups having five animals in each group by using Complete Randomized Design (CRD). For T₁ and T₂ concentrate mixture was composed by using procured feed ingredients with 16% crude protein level while T₃ was fed with commercial feed. Experimental animals of T₁ group was provided forest mixed fodder (adlib) + formaldehyde treated sesame cake included concentrate mixture @ 1.5% of body weight, T₂ group was provided forest mixed fodder (adlib) + sesame cake included concentrate mixture @ 1.5% of body weight whereas T₃ was provided forest mixed fodder (adlib) + commercial concentrate mixture @ 1.5% of body weight. Experiment revealed that higher intake of concentrate feed was recorded for T₃ (139.79 g) followed by T₂ (123.1 g) and T₁ (116.36 g) which was highly significant (P<0.001) among diet groups. Similarly, fodder intake was noted significantly higher (P<0.001) among diet groups (1425.1 g, 1422.5 g and 1321.4 g for T₂, T₁ and T₃, respectively).

Keywords : goats, bypass protein feeding.

GJSFR-D Classification : FOR Code: 111199



EFFECT OF FORMALDEHYDE TREATED SESAME CAKE FEEDING ON GROWTH PERFORMANCE OF GROWING FEMALE GOATS IN FODDER BASED BASAL DIET

Strictly as per the compliance and regulations of :



RESEARCH | DIVERSITY | ETHICS

Effect of Formaldehyde Treated Sesame Cake Feeding on Growth Performance of Growing Female Goats in Fodder based Basal Diet

M.R. Tiwari^α, D. Adhikari^σ, D.P. Adhikari^ρ, R.P. Ghimire^ω & S.H. Ghimire[¥]

Abstract - An experiment was carried out on fifteen growing female goats (50% Barberi 6, Kiko 6 and Khari 6) at the Agriculture Research Station (Goat), Bandipur for 90 days. Female goats of an average six months age and of body weight 7.82 kg were allocated into three groups having five animals in each group by using Complete Randomized Design (CRD). For T₁ and T₂ concentrate mixture was composed by using procured feed ingredients with 16% crude protein level while T₃ was fed with commercial feed. Experimental animals of T₁ group was provided forest mixed fodder (adlib) + formaldehyde treated sesame cake included concentrate mixture @ 1.5% of body weight, T₂ group was provided forest mixed fodder (adlib) + sesame cake included concentrate mixture @ 1.5% of body weight whereas T₃ was provided forest mixed fodder (adlib) + commercial concentrate mixture @ 1.5% of body weight. Experiment revealed that higher intake of concentrate feed was recorded for T₃ (139.79 g) followed by T₂ (123.1 g) and T₁ (116.36 g) which was highly significant (P<0.001) among diet groups. Similarly, fodder intake was noted significantly higher (P<0.001) among diet groups (1425.1 g, 1422.5 g and 1321.4 g for T₂, T₁ and T₃, respectively). Feed intake of different genotypes of goats was found to be non-significant whereas fodder intake was found to be significant (P<0.05) among goat breeds. In addition, feed conversion ratio per kg body weight gain was observed higher for T₃ (13.25:1) followed by T₂ (12.52:1) and T₁ (11.91:1). Likewise, daily crude protein intake was found higher for T₂ (82.59 g) followed by T₁ (81.4 g) and T₃ (79.13g) whereas crude protein expenditure per kg weight gain was found to be higher in T₃ (1.6 kg) followed by T₂ (1.52 kg) and T₁ (1.44 kg) which was none significant among diet groups. Initial body weight of T₁, T₂ and T₃ was 7.55 kg, 8.0 kg and 7.9 kg, respectively that reached 12.95 kg, 12.65 kg and 12.25 kg during 90 days of experiment for T₁, T₂ and T₃, respectively. Both initial and final body weight was non-significant among diet groups. Similarly, there was also non-significant effect of goat breed on body weight gain. Total body weight gain was recorded higher for T₁ (5.10 kg) followed by T₂ (4.9kg) and T₃ (4.45 kg) which was insignificant among diet groups. Similarly, average daily gain was also noted higher in T₁ (56.66g) followed by T₂ (54.44 g) and T₃ (49.44 g).

Keywords : goats, bypass protein feeding.

I. INTRODUCTION

Goats, important domestic animals in many part of world, have served mankind for ages. They provide substance in term of food and clothing. These hardy ruminants can exist in harsh environment in which other livestock species would perish. Goats grow and reproduce under extreme conditions from rugged mountain areas where winters are bitter cold to desert regions where it is hot and dry, and water and forage are sparse. The goat has been considered as *poor man's cow* (mini cow) of poor people because of its immense contribution in rural economy and national income. Goat products like milk and meat are not only nutritious and easily digestible but also a great source of regular income for the poor, landless and marginal farmers. Being a small sized animal it can be easily maintained by women and children (Aziz 2010). Goat population of Nepal is estimated to be 9.19 million. Of 9.19 million, goat population of western hills is 1.13 million which account 12.32% of total goat population that producing 5284 mt meat per annum (MoAD 2012).

Bypass protein is that the original amino acids in the protein meal are absorbed in the small intestine instead of converted to microbial protein in the rumen. Another benefit of feeding meals with high bypass protein is that the portion of the protein that is rumen degradable breaks down in the rumen very slowly. Formaldehyde treatment is most widely used chemical treatment for the protection of protein. Generally there is increased fecal nitrogen and decreased urinary nitrogen which indicates effectiveness of protection. The use of formaldehyde to protect dietary protein for ruminants is based on the premise that bound formaldehyde markedly reduces the solubility of the protein thereby rendering it highly resistant to microbial attack in the rumen without significantly reducing its digestibility in the small intestine.

Supplementation with other palatable materials, mainly agro-industrial by-products has been used in many developed countries for improving locally available nutrients of feed resources (Xianjun *et al.* 2012). It is well established that feed cost accounts for more than 70% of the total cost in any livestock production. Hence, it is paramount important to

Author ^α : National Animal Science Research Institute (NASRI), Khumaltar, Lalitpur. E-mail : tiwari65@yahoo.com

Authors ^{σ ρ ω ¥} : Agriculture Research Station (Goat), Bandipur, Tanahun.

incorporate locally available byproducts and raw materials into the feed of ruminant animals. Agricultural and industrial by-products that are relatively cheaper are best sources for supplementation of animals on fibrous basal feeds. Sesame cake is one of the byproducts available in Nepal at cheaper price throughout the year. The sesame oilseed cake contains DM (83-96%), CP (23-46%), ash (7.5-17%), ether extract (1.4-27%), NFE (25-32%) and crude fiber (5-12%), respectively (FAO 1990).

Growth comparison of goats fed with formaldehyde treated sesame cake is not evaluated so far in Nepal. Hence, a study was carried out to compare the growth performance of growing female goats fed with formaldehyde treated sesame cake mixed concentrate mixture at Agriculture Research Station (Goat), Bandipur, Tanahun.

II. METHODOLOGY

a) Experimental Animals

This experiment was carried out on fifteen growing female goats (50% Barberi 5, Khari 5 and Kiko

goats 5) at Agriculture Research Station (Goat), Bandipur, Tanahun from 30 March to 26 June 2013 (069/12/17 to 070/03/12). Female goats of average five months old with average body weight of 11.86 kg were allocated into three groups having six animals in each group by using Complete Randomized Design (CRD). They were drenched with Fenbendazole @ 5 mg/kg body weight against internal parasites before assigning in experiment.

b) Concentrate mixture composition

Feed ingredients maize, soybean cake, rice bran, minerals and salt were procured from Khowpa Feed Industry, Bhaktapur. For T1 and T2 concentrate mixture were composed by using procured feed ingredients with 16% crude protein level that has been presented in Table 1 while for T3 commercial compound feed was used made by Pancharatna Feed Industry, Narayangadh, Chitwan.

Table 1 : Composition of concentrate mixture

S/n	Ingredients	T 1		T2	
		Part	Crude Protein (%)	Part	Crude Protein (%)
1	Maize	50	4.4	50	4.4
2	Rice bran	18	1.58	18	1.58
3	Til cake	20	5.61	20	5.61
4	Meat com bone meal	10	4.92	10	4.92
5	Mineral mixture	1	0	1	0
6	Salt	1	0	1	0
Total		100	16.51	100	16.51

c) Formaldehyde treatment of sesame cake

Sesame cake was treated with 1-1.2g formalin (40%)/100 g crude protein (CP) as suggested by (Thomas *et al.* 1979; Hagemester *et al.* 1980). At first one part of formalin was diluted in nine part of water. That after formalin diluted solution was sprayed over cake and mixed manually for five minutes then the cake was stored in plastic bags.

d) Experimental diet of the animal

The dry matter requirement of goats was calculated based on 5 kg per 100 kg body weight. Following diets were formulated to the experimental animals (Table 2).

Table 2 : Experimental diets of the animals

Treatment	Experimental diet
1	Forest mixed fodder (adlib) + formaldehyde treated sesame cake included concentrate mixture @ 1.5% of body weight
2	Forest mixed fodder (adlib) + sesame cake included concentrate mixture @ 1.5% of body weight
3	Forest mixed fodder (adlib) + commercial concentrate mixture @ 1.5% of body weight

e) Feeding Regime

Concentrate mixture and *adlib* amount of fodder was provided to the experimental animals individually in plastic vessel. Concentrate mixture was provided once a day in the morning whereas fodder twice a day (morning

and evening). Quantity of concentrate mixture and fodder given daily to the animals was weighed daily and refusal was weighed in next morning. Experimental animal had free access to drinking water.

f) *Chemical Analysis*

The samples of feed ingredients, prepared concentrate mixture and forest mixed fodder were sent to the Animal Nutrition Division, Khumaltar, Lalitpur for proximate analysis. Representative samples were analyzed for dry matter (DM), crude protein (CP), crude fibre (CF), ether extract (EE) and total ash contents (TA). The DM was determined by oven drying at 100°C for 24 hrs. Crude protein of the samples was determined using the Kjeldahl method. Ether extract was determined using Soxhlet apparatus. Ash content was determined by ashing at 550°C in a muffle furnace for 16 hrs (AOAC 1980). Crude fibre of the samples was determined using the Van Soest method (Goering, H.K. and Van Soest 1970).

g) *Results Recording*

The trial period consisted 90 days after an adaptation period of 7 days. Total feed intake by the

goats was recorded daily for entire experimental period. The body weight gain of individual animal was measured fortnightly in the morning before feeding.

h) *Data Analysis*

Data of feed intake and body weight gain were analyzed by "*One Way Anova*" test for every measurement using computer statistical package Minitab 2003, versions 13.20.

III. RESULTS AND DISCUSSION

a) *Chemical composition of feedstuffs*

The result of chemical analysis has been given in Table 3 and crude protein content of prepared concentrate mixture was verified in laboratory that is presented in Table 4.

Table 3 : Chemical composition of different feed ingredients (% DM basis)

Ingredient	DM	OM	TA	CP	CF
Maize	87.69	97.97	2.03	7.53	2.34
Rice bran	87.85	89.5	10.5	11.52	4.83
Sesame cake	90.3	92.63	7.37	30.5	11.3
Meat come bone meal	93.22	67.2	32.8	49.93	3.43
Mixed forest fodder	39.94	90.01	9.99	11.16	NA

The calculated value of crude protein was verified with laboratory analysis, which is presented in Table 4.

Table 4 : Chemical composition of prepared concentrate mixture (% DM basis)

Particular	DM	OM	TA	CP	CF
Formaldehyde treated sesame cake included concentrate mixture	91.92	92.54	7.46	16.83	5.95
Sesame cake included concentrate mixture	91.58	92.10	7.90	16.92	5.64
Commercial feed	90.74	89.85	10.15	15.94	6.45

b) *Feed Intake*

Average daily intake of concentrate mixture and fodder by goats during experimental periods has been presented in Table 5.

Table 5 : Average feed intake of experimental animal/day/animal

Feedstuffs	Mean \pm SD		
	T1	T2	T3
Feed intake (g)	116.36 \pm 52.92	123.10 \pm 49.44	139.79 \pm 47.39
Fodder intake (g)	1422.5 \pm 389.5	1425.1 \pm 397.9	1321.6 \pm 406.3
Daily crude protein intake (g)	81.4 \pm 23.46	82.59 \pm 23.38	79.13 \pm 23.01
Total protein intake (kg)	7.33	7.43	7.12
Daily dry matter intake (g)	675.11	681.91	654.7
Total dry matter intake (DMI) (kg)	60.76	61.36	58.92
Feed conversion ratio (FCR) (kg)	11.91:1	12.52:1	13.24:1
Crude protein expenditure per kg weight gain (kg)	1.44	1.52	1.6

Higher intake of concentrate feed was recorded for T₃ (139.79 g) followed by T₂ (123.1 g) and T₁ (116.52 g) which was highly significant (P<0.001) among diet groups. Similarly, fodder intake was also noted highly significant (P<0.001) among diet groups (1425.1 g,

1422.5 g and 1321.6 g for T₂, T₁ and T₃, respectively). Highest dry matter intake per day was higher in T₂ (681.91 g) followed by T₁ (675.11 g) and T₃ (654.7 g) which resulted higher total dry matter intake (61.36 kg, 60.76 kg and 58.92 kg for T₂, T₁ and T₃, respectively).

Feed intake of different genotypes of goats was found to be non-significant among goat breeds whereas fodder intake differed significantly ($P < 0.05$). In addition, feed conversion ratio per kg body weight gain was observed higher for T_3 (13.24:1) followed by T_2 (12.52:1) and T_1 (11.91:1).

Similarly, average crude protein intake per day per animal was recorded for T_2 (82.59 g) followed by T_1 (81.4 g) and T_3 (79.13 g) which resulted higher total crude protein intake (7.43 kg, 7.33 kg and 7.12 kg for T_2 , T_1 and T_3 , respectively). Furthermore, crude protein expenditure per kg weight gain was found higher in T_2 (1.52 kg) followed by T_1 (1.44 kg) and T_3 (1.6 kg). Both average crude protein intakes per day and crude protein expenditure per kg body weight gain was none significant among diet groups.

c) Growth Performance

The growth performance of experimental goats is presented in Table 6 and Figure 1. Initial body weight of T_1 , T_2 and T_3 was 7.55 kg, 8.0 kg and 7.9 kg, respectively that reached 12.95 kg, 12.65 kg and 12.25 kg during the experimental period for T_1 , T_2 and T_3 , respectively. Both initial and final body weight was non-significant among diet groups. Similarly, there was also non-significant effect of goat breed on body weight gain. Total body weight gain was recorded higher for T_1 (5.1kg) followed by T_2 (4.9 kg) and T_3 (4.45 kg) which was non-significant among diet groups. Similarly, average daily gain was also noted higher in T_1 (56.66g) followed by T_2 (54.44 g) and T_3 (49.44 g) which also non-significant among diet groups.

Table 6 : Growth performance of goats

Parameter	Mean \pm SD		
	T1	T2	T3
Initial Body weight (kg)	7.55 \pm 0.79	8.0 \pm 1.35	7.9 \pm 2.66
Initial metabolic weight (kg)	4.55	4.75	4.71
Final Body weight (kg)	12.95 \pm 1.17	12.65 \pm 1.59	12.25 \pm 3.45
Final Metabolic weight (kg)	6.83	6.70	6.55
Total weight gain (kg)	5.1 \pm 1.46	4.90 \pm 1.15	4.45 \pm 1.20
Average daily gain (g)	56.66 \pm 16.27	54.44 \pm 12.82	49.44 \pm 13.38




Figure 1 : Body weight gain trend of goats during experiment period

IV. DISCUSSION

This study was initiated with objective to compare the growth performance of female goats of different breeds fed with formaldehyde treated and none treated sesame cake incorporated concentrate mixture,

and commercial compound feed on fodder based basal diet. Result revealed that concentrate feed and fodder intake was highly significant ($P < 0.001$) among diet groups. Both initial and final body weight was not significant among groups. Similarly, total weight gain and average daily gain also found to be non significant

among the diet groups. Furthermore, average crude protein intake and crude protein expenditure per animal per day was also not significant among diet groups. Several work has been done by different researchers to improve the bypass protein level in goats through heat treatment but feeding of formaldehyde treated sesame cake to goats is not recorded / documented.

V. CONCLUSION

Our experiment revealed that there is no significant effect of formaldehyde treatment of sesame cake on body weight gain of goats, however, feed and fodder intake differed significantly. Therefore, it can be concluded that formaldehyde treatment of sesame cake does not improve the bypass of protein in goats. Hence, sesame cake can be incorporated in concentrate mixture of goats without any treatment.

VI. ACKNOWLEDGEMENT

The authors would like to express their most sincere gratitude and appreciation to Dr. Bhoj Raj Joshi, Director, National Animal Science Research Institute, Khumaltar, Lalitpur for his encouragement, help, suggestion and support during experimental period. Similarly, work of Mrs. Bishnu Ale (Enumerator) and Mr. Tek Bahadur Ale (Junior Technician of ARS, Bandipur) deserve high appreciation without which conduction of experiment was impossible. Mr. Basanta Kumar Shrestha Technical Officer, Animal Nutrition Division, Khumaltar also deserve appreciation for his dedicated work in chemical analysis of fed ingredients and fodders. Likewise, our thanks also go to the all-technical, admin and finance staff of National Animal Science Research Institute, Khumaltar and Agriculture Research Station (Goat), Bandipur, Tanahu for their help during trial period.

REFERENCES RÉFÉRENCES REFERENCIAS

1. (1980). *Association of Official Analysis Chemists*,. Official methods of analysis, U.S.A.
2. Aziz, M.A. (2010). Present status of the world goat populations and their productivity. 45 (2), 42.
3. (1990). Guide to Food and Agriculture Organization (FAO) Documents (Website).
4. Goering, H.K. and Van Soest (1970). *Forage fibre analysis apparatus, reagents, procedures and some application*.
5. Hagemester H., W. Luppig and W. Kaufmann, W. Haresign and D. Lewis, (1980). Microbial protein synthesis and digestion in high yielding dairy cows. 67-85.
6. (2012). *Ministry of Agriculture Development, Agri-Business Promotion and Statistic Division*. Statistical information in Nepalese agriculture, Nepal.

7. Thomas, E., A. Trenkle and W. Burroughs (1979). Evaluation of protective agents applied to soybean meal and fed to cattle. 49:, 1337-1345.
8. Xianjun Yuan, Chengqun Yu, Shimojo, M & Tao Shao (2012). Improvement of Fermentation and Nutritive Quality of Straw-grass Silage by Incubation of Wet Hulless-barley Distillers Grain in Tibet. 25 (4), 479-485.



This page is intentionally left blank



GLOBAL JOURNAL OF SCIENCE FRONTIER RESEARCH
AGRICULTURE AND VETERINARY
Volume 13 Issue 11 Version 1.0 Year 2013
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Value-Chain Analysis for Coastal Fisheries Development in Nigeria

By Odebiyi O.C, George F.O.A, Odulate D.O., Agbonlahor M.U. & Olaoye O.J.

Federal University of Agriculture, Abeokuta, Ogun State, Nigeria

Abstract - This study evaluated the coastal fisheries value chain (CFVC) of Ogun Waterside Local Government Area (LGA), Nigeria. 176 fisherfolks were selected using purposive and simple random sampling techniques for interview schedule administration. Data collected were analyzed using descriptive and inferential statistical tools. Fishermen, fish processors and fish-marketers were the major actors of the CFVC and their average ages were $34.79+0.57$, $39.05+0.58$ and $36.73+0.65$ years; 91.7%, 57.8% and 70.4% of them were married, having an average household size of 9, 10 and 6 persons respectively. Actors in the CFVC were mainly uneducated (43.1%, 31.2% and 66.0% respectively). Chi-Square analysis revealed a significant association ($P < 0.05$) between the socio-economic characteristics and total revenue of the respondents. There was no significant association between the constraints faced by the fishermen ($\chi^2=0.253$, $P > 0.05$) and total revenue; although it was significant for the fish processors ($\chi^2=0.000$, $P < 0.05$) and fish-marketers ($\chi^2=0.000$, $P < 0.05$) at 95% fractile level. The study concluded that poor transportation network, high cost of fishing inputs, inadequate funding and poor storage facilities were major constraints to CFVC development and that the fish-marketers stage was the most profitable compared to the other stages in the CFVC.

Keywords : value, coastal, artisanal, chain, development.

GJSFR-D Classification : FOR Code: 830199



VALUE-CHAIN ANALYSIS FOR COASTAL FISHERIES DEVELOPMENT IN NIGERIA

Strictly as per the compliance and regulations of :



© 2013. Odebiyi O.C, George F.O.A, Odulate D.O., Agbonlahor M.U. & Olaoye O.J. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License (<http://creativecommons.org/licenses/by-nc/3.0/>), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Value-Chain Analysis for Coastal Fisheries Development in Nigeria

Odebiyi O.C^α, George F.O.A^σ, Odulate D.O.^ρ, Agbonlahor M.U.^ω & Olaoye O.J.[¥]

Abstract - This study evaluated the coastal fisheries value chain (CFVC) of Ogun Waterside Local Government Area (LGA), Nigeria. 176 fisherfolks were selected using purposive and simple random sampling techniques for interview schedule administration. Data collected were analyzed using descriptive and inferential statistical tools. Fishermen, fish processors and fish-marketers were the major actors of the CFVC and their average ages were 34.79±0.57, 39.05±0.58 and 36.73±0.65 years; 91.7%, 57.8% and 70.4% of them were married, having an average household size of 9, 10 and 6 persons respectively. Actors in the CFVC were mainly uneducated (43.1%, 31.2% and 66.0% respectively). Chi-Square analysis revealed a significant association ($P < 0.05$) between the socio-economic characteristics and total revenue of the respondents. There was no significant association between the constraints faced by the fishermen ($r^2=0.253$, $P > 0.05$) and total revenue; although it was significant for the fish processors ($r^2=0.000$, $P < 0.05$) and fish-marketers ($r^2=0.000$, $P < 0.05$) at 95% fractile level. The study concluded that poor transportation network, high cost of fishing inputs, inadequate funding and poor storage facilities were major constraints to CFVC development and that the fish-marketers stage was the most profitable compared to the other stages in the CFVC.

Keywords : value, coastal, artisanal, chain, development.

I. INTRODUCTION

Nigeria has become a major destination for imported seafood ever since the Government made a tariff reduction on all fishery products in 2001 from 25% to 5% (NBS, 2006). The European Union accounts for more than 70% of the Nigerian sea food supply while the US provides about 1% (Nzeka, 2003). The massive potentials of a coastline of 853km, a 200 nautical miles Exclusive Economic Zone (EEZ), over 2,658 fish farms as well as 937 Dams and Reservoirs, 365 lakes and reservoirs and 687 ponds and floodplains totaling over 13 million hectares of water bodies (Earth Trends, 2003) remain largely untapped. Despite the abundant fisheries resources and the relatively high consumption of fish in Nigeria (FDF, 2008), its domestic output of 0.85 million metric tonnes in 2010 still falls short of demand of 3.02 million metric tonnes (CBN, 2007; FDF, 2008; FDF, 2010). The fisheries sector

represent a major food source, which is invaluable for the protein they provide and the industrial products they produce. Fish is economically, socially and culturally important as a global dietary aspect of sustainable food security. Economically fish provides an important source of food and income for both men and women and fishing has an important social and cultural position in riverine communities. However A deficit of 2.17 million metric tonnes is required to meet the ever increasing demand for fish in Nigeria. This large deficit between the demand and supply of fish is augmented by massive importation of frozen fish which is a rigorous drain on the exchange earnings of the nation (FDF, 2008).

Availability of fish to the consumers at the right time, right form, right place and at the lowest possible cost requires an effective marketing system (Shamsuddoha, 2007). Marketing of fish passes through various market participants and exchange points before they reach the final consumers (Ali *et al.*, 2008). Nigeria has a great potential of fish resources whose distribution and value chain needs to be strengthened and developed to bridge the gap between demand and supply of fish in Nigeria (Amao *et al.*, 2006).

Fish is highly susceptible to deterioration without any preservation or processing measures (Okonta and Ekelemu 2005). Immediately fish dies a number of physiological and microbial deterioration sets in which invariably degrades the quality of fish (Eyo, 2001). It will become unfit for human consumption within about one day of capture, unless it is subjected to some form of processing or preservation. Even after the fish has been processed, particularly if traditional methods have been used, the fish is still subject to many forms of loss and spoilage. Hence, fish being a highly perishable substance needs to be transported to the consumer or final user in time (Ali *et al.*, 2008) to avoid post harvest spoilage through a coordinated marketing channel (fish value chain).

Based on the aforementioned, the following research questions become pertinent:

1. Who are the actors or institution that makes up the coastal fish value chain?
2. What are the socio-economic characteristics of the major actors that make up the coastal fish value chain?
3. What is the structure of the fresh and processed fish value chain?
4. What are the constraints to the identified fish value chain?

Authors $\alpha \sigma \rho$: Department of Aquaculture and Fisheries Management, Federal University of Agriculture Abeokuta, P.M.B. 2240, Abeokuta, Ogun State, Nigeria. E-mail : odebiyioc@funaab.edu.ng

Author ω : Department of Agricultural Economics, Federal University of Agriculture Abeokuta P.M.B. 2240, Abeokuta, Ogun State, Nigeria.

Author \forall : Agricultural Media Resources and Extension Centre, Federal University of Agriculture, P.M.B 2240, Abeokuta, Ogun State, Nigeria. E-mail : funksod4real@yahoo.com

a) Objectives of the Study

i. Main Objective

To analyse and develop the fish value chain in the coastal community of Ogun Waterside, Ogun State, Nigeria

ii. Specific Objectives are to

The following specific objectives were addressed in order to achieve the broad objective:

1. Identify and describe the socio-economic characteristics of all the institutions that make up fish value chain in the study area.
2. Identify and analyze the structure of the fresh and processed fish value chains
3. Determine the constraints hindering the development of the identified coastal fish value chain.

b) Justification of the Study

Hunger, malnutrition, and poor health are widespread and serious development challenges hindering economic development in Nigeria. These can only be overcome by providing the ever rising human population with better employment opportunities to improve their livelihoods (Akinrotimi *et al.*, 2007) and also adequate nutrition to reduce the nutritional deficiencies prevalent in Nigeria. The Nigerian fishery industry plays a significant role in the development of this country by providing livelihood for more than 50 million people in terms of employment, income generation and provision of principal protein to the diet (Akpaniteaku *et al.*, 2005). Each year, there are an additional 80-90million people to feed, most of them in developing countries, the most reliable source of protein for many of them is fish (Nzeka, 2003). However, irrespective of the great opportunities embedded in capture fisheries in Nigeria, a lot of the fish resources are being discarded on a daily basis due to an unorganized or uncoordinated distribution channel (Aihonsu and Shittu, 2008). Analyzing fisheries value chain provides an insight into various employment opportunities that remain untapped in the fisheries sector (Kaplinsky and Morris, 2000). Fish value chains in Nigeria are not yet developed to meet international market requirements as limited value addition is done in the industry, with the result that market for fish and fish products are limited to domestic markets (Investopedia, 2011). Over the past decade, development practitioners have increasingly shifted their attention from farming systems to targeting agricultural value chains and fisheries by extension, to improve smallholder production and participation in markets (Rota and Sperandini, 2010). This is because small-scale producers are often unable to increase production by adopting productivity-enhancing technologies unless the value chains for their products are sufficiently developed and dynamic (Kaplinsky and Morris, 2000).

c) Hypotheses

The hypotheses of the study tested in the null forms are:

- H₀1: There is no significant relationship between the socio-economic characteristics of the chain actors and their total revenue.
- H₀2: There is no significant relationship between the total revenue of the chain actors and the constraints faced.

II. MATERIALS AND METHODS

a) The Description of the Study Area

The geographical location chosen for this study is the Ogun waterside area of Ogun State, Nigeria. The proximity of the area to the Atlantic Ocean, Lagoon systems and in particular, to the good, albeit complex network of streams, rivers, and other water-bodies make the area an appropriate geographical location for this study. She is located in the eastern part of Ogun state sharing boundaries with Ondo state in the north, Lagos state in the south and Ijebu east local government in the west. About half to three quarter of the length of the local government is surrounded by water extending from Lagos state to Ondo state, this peculiar feature gave birth to the name waterside. The study area is closely associated with other maritime states of South-western Nigeria. The area comprises over 50 towns and villages with Headquarter at Abigi at 6°29'N 4°24'E / 6.483°N 4.4°E (www.wikipedia.com), while the main town in this area are Iwopin, Oni, Ibiade, Abigi, Efire, Ilushin, Makun-Omi, Ode-Omi and Lomiro, the area consists largely of Yoruba speaking people of which, the Ijebus comprise about 70 percent, with the Ikales, Ilajes, Itsekiris and Urhobos making up the remaining 30 percent. It has an area of 1,000 km² and a population of 72,935 at the 2006 census. This area is also blessed with a large expanse of fertile land (soil) rich in organic matter, well drained and deep which makes it support various crop cultivation especially plantation crops such as oil palm . The choice of the local government is by its close proximity to the Atlantic Ocean and its relative endowment with a complex network of streams, rivers, brackish water and in particular the extension of the Lagos (Lekki) Lagoon to the area. It is the only area of the state with a coastline on the Bight of Benin and also borders Lagos lagoon (Fig 1).

b) Sample Size and Sampling Techniques

A purposive sampling technique was used in the selection of four (Igbosere, JK Camp, Olosumeta and Elefon) coastal fishing communities from the 23 fishing villages of Ogun state coastal area based on their intensity of fishing activities. The second stage involves the use of simple random sampling in the selection of 30% of the fishermen and fish processors in the four selected communities who were interviewed to

give a total of 72 and 64 respondents respectively. Lastly a purposive sampling method was used in selecting 40 fish marketers at their central market arena to give a total of 176 respondents as indicated in the table 1.

i. *Analytical Procedures*

Combinations of statistical, budgetary and parametric analysis were used to analyze the obtained data. These include descriptive analysis, inferential statistics and profitability indicators.

ii. *Descriptive Statistical Tools*

Frequencies, tables, pie-charts, bar charts and percentages were used to describe the socio-economic characteristics of the respondents. The characteristics included the ages of the farmers, marital status, educational attainment, major occupation, farming experience, sex among others.

III. RESULTS

a) *The socio-economic characteristics of the fish value chain actors at the coastal area of Ogun state, Nigeria*

The socio-economic characteristics of all institutions and actors that make up the fish value chain in the study area are presented in Table 2. One hundred percent of the fishermen and fish processors were male and female respectively; while 14.2% and 85.8% of the fish-marketers were male and female respectively. Most (43.1%) of the fishermen fell within the age bracket of 41-50 years with a mean age of 43.79 ± 0.57 , 39.05 ± 0.58 and 36.73 ± 0.65 ; while 46.8% and 50.6% of the fish processors and fish-marketers were within the age range of 31-40 years respectively. However, the ages of 11.2%, 0.0% and 8.8% of the fishermen, fish processors and fish-marketers were higher than 50 years respectively.

Majority (91.7%, 57.79% and 70.4% respectively) of the fishermen, fish processors and fish-marketers were married with household sizes of 6-10 persons and a mean household size of 9, 10 and 6 persons. Most (54.2% and 52.8%) of the fishermen and fish-marketers practiced Islam, while 56.2% of the fish processors were Christians. Respondents with primary school leaving certificate were 36.1% (fishermen), 49.9% (fish processors) and 13.2% (fish-marketers) as against 31.2% of fish processors with no formal education; hence the high level of illiteracy among value chain actors in Ogun Waterside LGA.

The survey further revealed that out of the 100% fishermen, fish processors and middle men who practiced their occupation on a full time basis, 65.5% (fishermen), 40.6 (fish processors) and 0.0% (middle men) practiced farming as their minor occupation. Based on the years of experience in fish-related business, 26.41%, 74.9% and 37.4% of fishermen, fish

processors and fish-marketers respectively had between 11 and 20 years of experience. Out of the 72, 64 and 45 fishermen, fish processors and fish-marketers interviewed respectively, 72 (100.0%), 44 (68.6%) and 17 (38.4%) were not members of any cooperative societies. The structure of the identified coastal fish value chain as mapped out during the survey is presented in Fig 2.

b) *Constraints to fish value chain development*

79.2% and 62.4% of the fisherman and fish processors reported poor transportation network to their coastal communities as a very severe constraint to the fish value chain development in the coastal area of Ogun State, Nigeria while 82.2% of the fish-marketers regarded poor storage facilities as a major and very severe constraints to the fish value chain development as indicated in Table 5.

IV. DISCUSSION

This research project evaluated the fish value chain in the coastal area of Ogun State, Nigeria. The study identified 3 major marketing nodes along the coastal area, including: fishermen, fish processors and the fish-marketers. All fishermen (100%) and all fish processors (100%) in the Ogun Waterside coastal fish value were male and female respectively. This could be attributed to strenuous and tasking nature of the typical fisherman's work which the male gender could possibly handle better than the weaker female gender; but is opposed to the practice in Idaleketa, another coastal community in Ogun Waterside where only the female gender could fish. These findings were in line with the findings of Olubanjo *et al* (2007), Olawumi *et al* (2010) and Olaoye and Odebiyi (2011) where the business was gender biased. Many (43.1%, 46.8% and 50.6%) of the fisherman, fish processors and fish-marketers interviewed were between 31 and 40 years of age, a highly productive and active age when actors could undertake strenuous task. This is in line with the findings of Bello, (2000) and George *et al* (2010) that age had a positive correlation with agricultural productivity.

In rural communities, marriage is a respected and prestigious institution that bestows social status and recognition on people. The marital status of respondents as presented in Table 4 indicated that 91.7%, 57.7% and 70.4% of the respondents respectively were married, implying that most of the women involved in the fish value chain in the study area were married, suggesting that marriage is a cherished value in the study area, with a resultant increase in their household size. This corroborates with the findings of Alfred and Fagbenro, (2006) and Odulate *et al* (2011) who reported higher percentage of married women in the coastal communities of Ondo and Ogun states, Nigeria respectively. From these results, it could be inferred that Islam was mostly practiced by the

fishermen (54.2%) and fish-marketers (52.8%) than other religions; while the majority (56.2%) of fish processors were Christians. This could be attributed to the fact that fish cuts across all the religion unlike other animals like pig.

Majority (58.4% of fishermen, 80.0% of processors and 33.0% of fish-marketers) of the Actors in the Ogun Waterside coastal fish value chain had household sizes ranging between 6 and 10 persons. This relatively large household size may decrease the number of labour needed at different nodes along the coastal fish value chain. These results were similar to the findings of Fabusoro *et al.*, (2007) that average household size in Africa was about 9 persons.

Education is an important factor which can influence fish production and determine level of awareness on the rate of return on value addition in fish. Results from this study showed that 43.1%, 31.2% and 66.0% of fishermen, fish processors and fish-marketers respectively had no form of education at all; while 36.14% (Fishermen), 49.9% (Processors) and 13.2% (Fish-marketers) respectively had a maximum of primary school education. This confirmed the general opinion that most fisherfolks were illiterate or semi-illiterates; most of whom have dropped out of the formal school system (Olubanjo *et al.*, 2007; and Alfred *et al.*, 2008). The low level of education in the Ogun waterside coastal fish value chain could be explained by two reasons: firstly, fishing communities are generally isolated and marginalized, with limited or no access to education services. Available government schools are very far, at distances of at least 30 to 40 kilometers from coastal communities. Additionally, regular flooding and threats from hostile crocodiles and hippopotamus are other reasons for parents not to send their children to school. Another important factor, given that fishing is the main economic activity, is that children are often assimilated into the fishing industry at an early age by having to help their parents or guardians in fishing and fish processing. These children are born and brought up in remote fishing communities with limited or no access to education. Due to the above mentioned threats, children often abandon school if they had a chance to even attend one.

Experiences in fishing were determined by the number of years the fisherfolks had been in the business. The respondents' mean years of experience was found to be 23years, 21years and 15years for fishermen, processors and fish-marketers respectively. It is believed that this would enhance their efficiency. This finding is in line with Schumpeterian theory of economic development, which suggests that technical efficiency is influenced by technical knowledge and understanding in addition to other socio – economic environment in which the fisher folks must take decision (Kalirajan, 1990) as cited by Olaoye *et al.*, (2012)

Fish and fish related business as the major occupation in the study area is a function of the importance attached to it as a source of livelihood. All the respondents had fishing, fish processing and fish marketing respectively as their major occupation; thus are likely to commit more number of hours and efforts towards the success of these enterprises. Cooperative Society involves social participation that helps farmers to pool their resources together, have access to fisheries inputs and insights in emerging issues in the business. Membership of cooperatives therefore influences the adoption of improved fisheries technologies resulting in higher productivity and poverty alleviation. Majority of the actors (100%, 68.6% and 38.4% of the fisherman, fish processors and fish-marketers respectively) were not in any cooperative society; which may be the reason for not benefiting from any source of loan. This was a throwback to the position of Akinbile (1998) as cited by Odebiyi (2010) that cooperative groups ensure that their members derive benefits from the groups such as they could not derive individually. One of the biggest challenges faced by the seafood sector in the coastal fishing community is value addition. A deprivation of value addition facilities in the study area is a major setback to the fish value chain industry, most of the respondents regarded poor transportation network and lack of storage facilities as major constraints to the value chain development.

V. CONCLUSION

The value-chain for the fresh and processed fish in the study area is comprised of three main stakeholder groups before fish reaches the consumer. There are virtually no exports of artisanal captured fish, and so the value-chain is short and simple compared to some value chains in other countries. One of the biggest challenges faced by the seafood sector in the coastal fishing community is value addition. Lack of value addition facilities in the study area hindered the development of the fish value chain industry. Smoking was the only form of value addition at the coastal area of Ogun State, Nigeria. Lack of adequate road network hindered other means of value addition to the fish industry which could have created additional employment opportunities along the coastal zone of Ogun Waterside LGA, Nigeria.

a) Recommendation

The study recommends that there is need to develop a comprehensive and effective road networking in the study area as water is the only means of transportation which is a major constraint to the development of the coastal fish value chain. It should be noted that the development of the coastal fish value chain of Ogun waterside LGA would encourage youth involvement and participation in the value chain.

It is important to create awareness on the need for the formation of fish cooperative societies through which some of the challenges in the value chain can be successfully tackled. Credit facilities should be made available by financial institutions and agencies to improve fish value addition; for the purchase or leasing of fishing inputs (Canoes, Outboard Engine etc). Also, sophisticated storage facilities/cold rooms should be provided by government as well as cooperative societies for the storage of unsold fish.

REFERENCES RÉFÉRENCES REFERENCIAS

- Aihonsu, J.O.Y. and A. Shittu (2008). Comparative Economic Analysis of Fish Processing Technologies in two Fishing Communities in Lagos State, 315–321.
- Akinbile, L. A. (1998). Paper presented at NAERLS Workshop on Extension Communication Techniques. 11.
- Akinrotimi, O.A., Onunkwo, D.N. Cliffe, P.T. Anyanwu, P.E. and Orokotan, O.O. (2007). The role of fish in nutrition and livelihoods of families in Niger Delta, Nigeria. *International Journal of Tropical Agriculture and food systems*, 1 (4), 344-351.
- Akpaniteaku, R. C., Weimin, M. And Xinhua, Y. (2005). Evaluation of the contribution of fisheries and aquaculture to food security in developing countries. 28 (1 & 2), 28 – 32.
- Alfred, S.D.Y. and Fagbenro, O.A. (2006). Perception of tilapia farmers on information sources in the coastal region of Ondo State, Nigeria.
- Ali, E. A, H.I.M., Gaya and T.N. Jampada (2008). Economic Analysis of fresh fish marketing in Maiduguri Gamboru Market and Kachallari Alau Dam landing site of Northeastern Nigeria. 4, 23-6.
- Amao, J. O., Oluwatayo, I. B. and F. K. Osuntope (2006). Economics of Fish Demands in Lagos State, Nigeria. 19 (1), 25-30.
- Bello, M.O. (2000). "Categorization of Potential Adopters for Organic-Based Fertilizer Among Vegetable Farmers in Ojo LGA State. i.org
- (2007). Statistical Bulletin, Initiative Publishers 16, p 10.
- (2003). World Resources Institute, Environmental Information; <http://earthtrends.wri.org/>
- Eyo, A. A. (2001). Inventory of Storage life of fresh Africa fish on ice, 11.
- Fabusoro E, Lawal-Adebawale OA, Akinloye AK (2007). A study of rural livestock farmers' patronage of veterinary services for healthcare of small farm animals in Ogun State. 34 (1&2), 132- 138.
- Fagbenro, O. A., Akinbulumo, M. O and Ojo, S. O. (2006). Aquaculture in Nigeria-past Experience, present situation and future outlook (history, status and prospects). 35 (2), 23 – 26.
- (2008). Fisheries Statistics of Nigeria, Federal Department of Fisheries.
- (2010). Fisheries Statistics of Nigeria, Federal Department of Fisheries.
- George F.O.A., O. J. Olaoye, O. P. Akande and R. R. Oghobase (2010). Determinants of Aquaculture Fish Seed Production and Development in Ogun State, Nigeria. *Journal of Sustainable Development in Africa*, 12.
- <http://www.investopedia.com/terms/v/valuechain.asp>, 16, (2011).
- Kalirajan, K (1990). "On measuring economic efficiency". *Journal of Applied Econometric*, 5, 75 – 85.
- Kaplinsky, R. and Morris, M. (2000). A Handbook for Value Chain Research, International Development Research Center (IDRC), Canada.
- (2006). *Annual Abstract of Statistics of Nigeria*, National Bureau of Statistics, Nigeria.
- (2006). National Population Commission (NPC), Nigeria.
- Nzeka, U. (2003), June. Nigeria demand for sea food is high. An article accessed from www.findarticles.com/p/articles.
- Odebiyi O.C. (2010). *Impact of microfinance bank loan on aquaculture development in Ogun state, Nigeria*. A project report submitted to the Department of Aquaculture and Fisheries management, Federal University of Agriculture Abeokuta.
- Odulate D. O., George F. O. A. and Idowu, A. A. (2011). Role of Women in Fisheries in Coastal Wetland Areas of Ogun State.
- Okonta, A. A. and Ekelemu, J. K. (2005), November. A preliminary study of micro-organisms associated with fish spoilage in Asaba, Southern Nigeria. Pp. 557-560.
- Olaoye, O. J. and Odebiyi, O. C. (2011). Economic viability for the use of microfinance bank loan on aquaculture development in Ogun State, Nigeria. *International Journal of Fisheries and Aquaculture*, 3 (4), 70-77.
- Olaoye, O. J., Idowu, A. A., Omoyinmi, G. A. K., Akintayo, I. A., Odebiyi, O. C., & Fasina, A. O. (2012). Socio-Economic Analysis of Artisanal Fisher Folks in Ogun Water-Side Local Government Areas of Ogun State, Nigeria. *Global Journal of Science Frontier Research Agriculture & Biology*, 12 (4).
- Olawumi, A.T, Dipeolu, A.O and Bamiro, O.M. (2010). Economic Analysis of Homestead Fish Production in Ogun State, Nigeria. 31 (1), 13-17.
- Olubanjo, O.O., S.O. Akinleye and M.A. Balogun, (2007). Occupational characteristics, technology use and output determinants among fisher-folks in Ogun Waterside Area, Ogun State. *FAMAN (Farm Management Association of Nigeria) Journal*, 8 (2).
- Rota A and Sperandini S. (2010). Value chains, linking producers to the markets. 56.

31. Shamsuddoha, M. (2007). Supply and value chain analysis in the marketing of marine dried fish in Bangladesh and non-tariff measures (NTMs) in International trading. 1-11. <http://ideas.repec.org/p/ags/ea106/7941.html>.

Table 1 : Study Population and Location

Chain actors/ Fishing Communities	Fishermen	Fish Processors	Fish-marketers	Total
Igbose	20	22	40	52
Elefon	12	16		38
Olosumeta	23	14	40	47
JK Camp	17	12		39
Total	72	64	40	176

Source: Field survey 2012-2103




Figure 1 : A map of the study area Table 2: Socio-demographic variables of the chain actors in the fish value chain

Table 2 : Socio-economic characteristics of the coastal fisheries value chain

Variables	Value Chain Actors					
	Fishermen		Fish Processors		Fish-marketers	
	Freq	%	Freq	%	Freq	%
Gender						
Male	72	100.0	0	0.0	6	14.2
Female	0	0.0	64	100.0	39	85.8
Total	72	100.0	64	100.0	45	100.0
Age						
< 20	0	0.0	0	0.0	0	0.0
21-30	14	19.3	30	46.8	12	26.4
31-40	19	26.4	18	28.3	23	50.6
41-50	31	43.1	16	24.9	6	13.2
>50	8	11.2	0	0.0	4	8.8
Mean \pm S.E	43.79 \pm 0.57		39.05 \pm 0.58		36.73 \pm 0.65	
Total	72	100.0	64	100.0	45	100.0
Marital status						
Single	6	8.3	11	17.16	4	8.8
Married	66	91.7	37	57.79	32	70.4
Divorced	0	0.0	10	15.6	7	15.4
Widow	0	0.0	6	9.36	2	4.4
Total	72	100.0	64	100.0	45	100.0
Religion						
Christian	27	37.5	36	56.2	21	46.2
Islam	39	54.2	20	31.2	24	52.8
Traditional	6	8.34	8	12.6	0	0.0
Total	72	100.0	64	100.0	45	100.0
Household Size						
1-5	0	0.0	0	0.0	13	28.6
6-10	42	58.4	51	80.0	32	70.4
11-15	30	41.7	13	20.0	0	0.0
Mean \pm SD	9.04 \pm 0.12		10.02 \pm 0.05		6.08 \pm 0.12	
Total	72	100.0	64	100.0	45	100.0
Education Level						
No formal education	31	43.1	20	31.2	9	19.8
Primary Education	26	36.14	32	49.9	6	13.2
Secondary Education	15	20.85	12	18.72	30	66.0
Tertiary Education	0	0.0	0	0.0	0	0.0
Total	72	100.0	64	100.0	45	100.0
Major Occupation						
Fisherfolks	72	100.0	64	100.0	45	100.0
Total	72	100.0	64	100.0	45	100.0
Minor Occupation						
Trader	21	29.2	38	59.3	30	66.0
Farming	47	65.3	26	40.6	0	0.0
Driver	1	1.39	0	0.0	15	33.0
Herbalist	3	4.2	0	0.0	0	0.0
Total	72	100.0	64	100.0	45	100.0
Years of experience						
1-10	8	11.1	6	9.4	4	8.8
11-20	19	26.41	48	74.9	17	37.4
21-30	31	43.1	8	12.5	15	33.0
>30	14	19.5	2	3.1	9	19.8
Mean \pm SD	22.7 \pm 0.61		20.95 \pm 0.54		15.3 \pm 0.63	

Total	72	100.0	64	100.0	45	100.0
Cooperative Membership						
Yes	0	0.0	20	31.4	28	61.6
No	72	100.0	44	68.6	17	38.4
Total	72	100.0	64	100.0	45	100.0




Figure 2 : Map of the coastal fish value chain in Ogun Waterside LGA, Nigeria
Source : Field Survey, 2013



Hypothesis Testing

Table 3 : H₀₁ There is no significant difference between the socio-economic characteristics of the chain actors and their total revenue

Variable	Fisherman			Fish Processors			Fish-marketers		
	Value	Df	Sig.	Value	Df	Sig.	Value	Df	Sig.
Age	34.2	9	0.000*	60.44	12	0.000*	135.000	12	0.000*
Marital Status	2.751	3	0.432	55.744	12	0.000*	66.937	8	0.000*
Religion	18.852	6	0.004**	41.007	8	0.000*	30.536	4	0.000*
HHZ	3.840	3	0.279	18.051	4	0.001**	90.000	8	0.000*
YOEX	10.212	9	0.334	78.222	12	0.000*	135.000	12	0.000*
Education	8.887	6	0.180	40.000	8	0.000*	57.600	8	0.000*

Source : Field Survey, 2012-2013; *=Significance at 0.001, **= Significance at 0.005

Key : HHZ = Household size, YOEX = Years of experience

Table 4 : H₀₂: There is no significant difference between the constraints faced by the chain actors and their total revenue

Variable	Fishermen			Fish Processors			Fish-marketers		
	Value	Df	Sig.	Value	Df	Sig.	Value	Df	Sig.
Constraints	21.54	18	0.253	84.44	8	0.000*	100.64	12	0.000*

Source : Field Survey, 2012-2013; *=Significance at 0.001

Table 5 : Constraints to the coastal area fish value chain development

Variables	Fisherman			Fish Processors			Fish-marketers		
	Very Severe	Severe	Not Severe	Very Severe	Severe	Not Severe	Very Severe	Severe	Not Severe
1. Inadequate fund	90.4	6.9	2.78	56.2	31.2	21.6	60.0	40.0	0.0
2. Poor storage facilities	0.0	0.0	0.0	0.0	22.0	78.0	82.2	17.8	0.0
3. Social amenities	98.6	1.4	0.0	74.9	25.1	0.0	8.9	20.0	71.1
4. High cost of fish input	11.1	88.9	0.0	12.5	37.4	50.0	71.1	20	8.9
5. Seasonality of fish	25.0	31.9	43.1	31.2	56.2	21.6	40.0	60.0	0.0
6. Poor transportation network	79.2	19.4	1.4	62.4	37.6	0.0	8.9	20.0	71.1
7. Breakage/spoilage of fish	2.8	69.4	27.8	18.7	65.5	15.6	0.0	100.0	0.0
8. Unpredictable weather	12.5	8.3	79.2	0.0	0.0	0.0	0.0	0.0	0.0
9. Trawlers' threat	62.5	30.6	6.9	0.0	0.0	0.0	0.0	0.0	0.0
10. Pollution effluents	66.7	26.4	6.9	0.0	0.0	0.0	0.0	0.0	0.0

This page is intentionally left blank





GLOBAL JOURNAL OF SCIENCE FRONTIER RESEARCH
AGRICULTURE AND VETERINARY
Volume 13 Issue 11 Version 1.0 Year 2013
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Environmental Conservation Information Needs of Farmers in Owerri West Area of Imo State, Nigeria

By Nnadi, F.N., Chikaire, J., Echetama, J.A., Umunnakwe, P.C. & Ihenacho, R.A.

Federal University of Technology, Owerri, Nigeria

Abstract - The enormous consequences of the wanton devastation of the environment inspired this study that evaluated the environmental conservation in Owerri West Area of Imo State, Nigeria. The objectives of the study was; to determine the socio-economic characteristics of the farmers, ascertain the farmers environment conservation information needs, investigate their environmental conservation practices and identify the problem militating against sustainable environment conservation practices. The data for the study was collected from 120 randomly selected farmers from 6 communities out of the 18 existing communicates in the area of study. The data collected were analyzed with the use of descriptive statistics such as frequency counts, percentages, mean. The result showed that farmers needed information in the area of environmental disaster management and funding sources for environmental management. The perceived effects of environmental conservation were to improve the farmers' socio-economic life and reduce hazards. The results also showed that inadequate knowledge base of environmental conservation practices was a major problem militating against environmental conservation and that maturing was the most conservation practice carried out by the farmers.

Keywords : *environment, conservation, management, farmers, information.*

GJSFR-D Classification : *FOR Code: 300805*



Strictly as per the compliance and regulations of :



© 2013. Nnadi, F.N., Chikaire, J., Echetama, J.A., Umunnakwe, P.C. & Ihenacho, R.A. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License (<http://creativecommons.org/licenses/by-nc/3.0/>), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Environmental Conservation Information Needs of Farmers in Owerri West Area of Imo State, Nigeria

Nnadi, F.N.^α, Chikaire, J.^σ, Echetama, J.A.^ρ, Umunnakwe, P.C.^ω & Ihenacho, R.A.[¥]

Abstract - The enormous consequences of the wanton devastation of the environment inspired this study that evaluated the environmental conservation in Owerri West Area of Imo State, Nigeria. The objectives of the study was; to determine the socio-economic characteristics of the farmers, ascertain the farmers environment conservation information needs, investigate their environmental conservation practices and identify the problem militating against sustainable environment conservation practices. The data for the study was collected from 120 randomly selected farmers from 6 communities out of the 18 existing communicates in the area of study. The data collected were analyzed with the use of descriptive statistics such as frequency counts, percentages, mean. The result showed that farmers needed information in the area of environmental disaster management and funding sources for environmental management. The perceived effects of environmental conservation were to improve the farmers' socio-economic life and reduce hazards. The results also showed that inadequate knowledge base of environmental conservation practices was a major problem militating against environmental conservation and that maturing was the most conservation practice carried out by the farmers. It was however recommended that extension education campaigns on environmental conservation practices should be intensified and the socio-economic determinants of the farmers' information need to be considered in interventions and advocacies.

Keywords : environment, conservation, management, farmers, information.

I. INTRODUCTION

Today's farmers are under unprecedented pressure. The world's population is closing in on seven billion, and it is projected to reach nine billion by 2050. Billions of those people will be enjoying an improving standard of living, including increased consumption of more nutritious food such as milk, meat and energy. A crowded planet adds to the environmental challenges of feeding, clothing and powering the world. Water supplies will be increasingly scarce, threatened by pollution, and diverted to population centers. We can no longer set out to farm new frontiers- we must make every acre already being

farmed even more productive and prevent environmental degradation (Towery and Werblow, 2010). With shrinking resources and little margin for expansion, the stakes of environmental degradation are too high. Protecting our soils, air and water- and our forests, wetlands and grasslands- is vital to all of us in the long term. Environmental and economic sustainability are essential on every farm (Towery and Werblow).

Norman Borlaug, the legendary plant breeder and Nobel laureate who was the driving force behind the Green Revolution of the 1960s and 1970s, summed up the task when he wrote, "Over the next 50 years, the world's farmers and ranchers will be called upon to produce more food than has been produced in the past 10,000 years combined and to do so in environmentally sustainable ways" (Towery and Werblow, 2010). Increases in agricultural production are possible through modern methods. But these advances in agricultural production are possible through modern methods. But these advances will be useless unless there is enough good land for farming. If the soil on which all agricultural and all human life depends is wasted away, then the battle to free mankind from want cannot be won (ECO-ISSUES, 2010). It is estimated that agricultural production will need to increase by 60 per cent in the next two decades to feed the world. The large part of this will need to come from an intensification of agriculture on lands already being cultivated. In addition, the area of cultivated land should be expanded by 200 million hectares (Favis – Mortlock,2005). An uncertainty in these estimates is the amount of land which is being lost through degradation.

In precise term, multiple relationships exist between man, his activities and the environment, but he has a positive role to play in shaping the environment. An influence how human lives and organizes his life, but he has equally helped to modify and rebuild the environment through time. There is an ecological relationship between man and his environment as evident in the harvesting and exploitation of the environmental resources (France and Kathleens, 2002). However, the nature and the degree of hazards done to the environmental vary according to technology of the people (be it indigenous or modern technology). It depends on the prevailing physical, technological,

Authors α σ ρ ω : Department of Agricultural Extension, Federal University of Technology, Owerri.

Author ¥ : Department of Agricultural Management and Extension Technology, Imo State Polytechnic, Umuagwo, Owerri, Imo State, Nigeria. E-mail : aksymeon35@yahoo.com

cultural and political circum-stances (France and Kathleens, 2002).

The deterioration of the environment which started when man first settled into villages has been made worse by increasing human activities such as deforestation, burning etc. Natural phenomena like desertification, drought, erosion etc have not helped matters through their degrading effects on the soil fertility, the ecosystem services of the natural environment (France and Kathleens, 2002). Human has influenced his physical environment by fertilizing the poor soil, terracing the hillsides, construction roads, railways and bridges, exploiting the forest resource like timber and extracting the metallurgic and non-metallurgic iron from the environment. Through these processes, he has changed or altered the environment to suit himself.

Sustainable development is necessary because it is capable of alleviating poverty. Therefore, all human must work towards a sustainable environment. To this end, we must have development which is sustainable and built on firm ecological foundation. The attainment of environmentally sustainable development largely depends on our willingness to be genuinely committed to the use of environment through the use of innovations and substitutes of most of the earthly minerals.

However, the fact that this state (IMO STATE) is blessed with immense mineral wealth and considerable agricultural potential, varied climatic zone, continuous sunlight and abundant rainfall which supplies many crops, fibre and medicinal plants and forest products make it imperative to sustain the environment. Although impressive progress in the area of improving the environment has been made in recent years, there seem to be a considerable lack of understanding of the fundamental mechanisms, sources and classification of pollutants causing pollution of different forms (IMSG, 2001). Even with the amount of effort rendered by government and non-government organization to farmers in both rural and urban areas towards conservative measures, farmers still lack essential information in managing their environment. It is vital to create awareness to the public and farmers toward the use of environmental conservation approaches particularly to when every aspect of our lives including agricultural are seriously threatened by variability in weather climate change. The broad objective of the study is to analyze the environmental conservation information needs of farmers in Owerri-west Area of Imo State. The specific objectives of the study include, to;

1. Investigate the socio-economic characteristics of the farmers'.
2. Ascertain the farmers' environment conservation information needs.
3. Investigate their perceived effects of environmental conservation practices.

4. Investigate their environmental conservation practices.
5. Investigate the problems militating against sustainable environmental conservation practices.

II. METHODOLOGY

1. The study area was carried out in Owerri-west area of Imo State. It is located in the rain forest zone about 120 Km north of the Atlantic coast and lies on latitude 4° 14 north and 6° 15 north, longitude 6° 51 east 8° 09 east (ISADAP, 2000). It has a population of 250,000 people and an area of 295 square kilometers in 2013 (NPC, 2006). It shares boundaries with Ngor Okpala LGA in south, Owerri municipal council in the east, Mbaitolu LGA in the north and Ohaji/Egbema LGA in the west. The average annual rainfall measures up to 2550 millimetres, the relative mean temperature ranges annually between 24.5° and 25.5° and the humidity varies according to the time of the year (ISADAP, 2000). The people are land resource dependent. The farmers' produce crops like cassava, maize, yam, plantain and rear animals like goat, pigs, fish, birds and recently rabbits. Owerri-west was purposely selected because of its nearness to the researcher and the predominance of farmers in the area. The entire farmers in owerri-west constitute the sampling population. Multi-stagerandomsamplingtechnique was employed in the sampling. First, a random sampling technique was used to select 6 communities from the existing 18 communities. These include: Oforola, Obinze, Eziobodo, Ihiagwa, Amakohia- ubi and Orogwe. From each of the communities, 4 villages were sampled (2nd stage) and from each of the villages, 5 farmers' comprising male and female were randomly sampled to give a sample size of 120 respondents. Data were collected from primary and secondary sources. Primary data were obtained from farmers' in the study area through structured questionnaire to be complemented with interview schedule for farmers' who are not literate enough. Secondary data were collected from annual reports of the Imo state Agricultural Development Programme (ADP), the Imo State Ministry of petroleum and Environment, textbooks, journals, internet and previous studies of other researchers who worked on related topics. The data collected were analysed using descriptive statistical tools such as frequency tables, percentage and mean. Specifically, objectives 1, 4 and 5 which are to investigate the socio-economic characteristics of the farmers; investigate their environmental conservation practices and investigate the problems militating against sustainable environmental conservation practices were analysed with percentages. Objectives 2 and 3

which are to ascertain the farmers' environment conservation information needs; and investigate their perceived effects of environmental conservation practices were analysed with mean. The formula used was $x = \frac{\sum X}{N}$. the likert type scale of agreement had their weights added together and divided by the number of scales. $SA=5, A=4, U=3, SD=1. (5+4+3+2+1)/5 = 3$ (discriminating index for acceptance and rejection). This means that all mean value of 3 and above were accepted as important.

III. RESULTS AND DISCUSSIONS

a) Socioeconomic Characteristics of Respondents

The result in the table 1 showed that farmers within age range of 20-29 composed 7.5 percent of the sample, those between 30 and 39 represented 16.67 percent whereas 40-49 years represented 24.17 percent. The farmers between the ages of 50 and 59 and 60-69 represented 28.33 percent and 23.33 percent respectively. The mean age of the farmers was 49. This means that the farmers' are ageing. This implies that they will not readily accept agricultural innovations (Mgbada, 2010). The dominance of the old could be adduced to youth emigration to cities for which collar jobs. This result agrees with Nnadi and Amaechi, (2007) that youth migration from the rural to the Urban areas left agriculture in the hands of the old and ageing farmers. Also Mgbada, (2010) noted that Nigerian farmers are mostly old people. There sultalso showed that the percentage of male farmers in the study was 49.17 percent. The female farmers composed 50.83 percent. The dominance of female farmers in the study could be attributed to increasing engagement of men in politics and rising cases of female headed households. The result agrees with Mollet, (1990) and Mgbada, (2010) who noted that more than one third to half of the total labour contribution to agriculture is made by women. The result on the marital status distribution of the farmers shows that single farmers represented 10.83percent of the sample. The married farmers' represented 70.83 percent, while divorced and widowed represented 3.33 percent and 15 percent respectively. The dominance of married farmers in agricultural production could be to ensure household food security. The result of the distribution of the farmers by education reveals that farmers with no formal education consisted a percentage of 18.33%. The table also shows that 31.09 percent of the respondents had primary and secondary education respectively. While 20.17% had tertiary education. The result implies that the farmers are highly literate with about 81 percent having acquired one level of education or the other. This means that they can be easily convinced to accept better practices of their farming operations. It is in conformity with Nnadi and Amaechi, (2004) that a greater deal of change has

occurred within the rural communities in recent times due to the introduction of education. Their high literacy level is an asset as the farmers would be exposed to many information sources, embrace innovations and analyze farm situations objectively. It also supports the view made by Nnadi and Onuoha, (1999) that educated farmers' are good adopters of innovations. The distribution of the farmers by occupation shows that 62.5 percent of the respondents have farming as their major occupation. Civil servants and traders represented 17.5 percent of the sample respectively while artisans represented 2.5 percent. The result further reinforces agriculture as the basis of rural economies where majority are farmers. Thus, farming is a way of life by cultural dictates. This confirms the assertion of Olayide et al., (1981) that Nigeria is characterized as a nation of small scale farmers and the rural population comprises full and part time farmers. The farmers' distribution by household is represented and the result indicates that 15.83% percent of farmers have household size of 1-5 members. A total of 49.17 percent of the farmers have 4-6 persons in their household. Whereas 32.5 percent had 7-9 persons, 2.5 percent had 10 persons and above. The mean household size was 6. The size is modest and in consonance with the household population policy of Nigeria. A large family size could entail many hands in family labour supply, typical of Nigerian agriculture which requires so many hands for labour supply. As farmers and farm families are closely knit, farmers prefer using labour from their household, wives and children (Nnadi and Amaechi, 2004). However, this could imply more household expenditure for food security, social needs and economic needs provision. Thus, production capital could be converted to consumptive capital. The result of the distribution of the farmers by farm size indicates that farmers with farm size less than one hectare represented 49.17 percent of the respondent. They are followed by 29.17 percent that have a farm size of 1-3 hectares. The result also revealed that 17.5 percent of the respondents have between 4-6 hectares of farm land. while 4.16 percent have 7 hectares and above. The mean farm size is 2.16 hectares. The above situation reveals that there is skewedness in the distribution of land in the study area. The greater number (49.17 percent) have small area usually fragmented holding which supported subsistence agriculture. The result supports the assertion made by Nnadi and Amaechi, (2004), that under inheritance, the whole heirs of every family have their shares of land no matter how fragmented and small their sizes area. This lays credence to the subsistence farming characteristic in the southeast. From the result, a dominant number of the farmers indicated membership of social organization. Specifically, 64.7 percent indicated that they belong to one social organization or the other. However, on the contrary, 43 farmers represented by 35.83 percent

indicated that they do not belong to any social organization. Membership of Social organization satisfies the social needs of farmers in additions to serving as an avenue for access to information on agricultural technology. Farmers by virtue of their membership discern the obvious advantages of agricultural technology as well as clarify their misconception of technology and the adoption. However, the farmers do not belong to social organization may have been ignorant of the obvious gains through membership-the however calls for conceited extension campaigns for farmers member of co-operative society and combined.

b) Areas of Information Needs

The distribution of the farmers according to areas of information is presented in table 2 using five point likert-type scale of agreement, the discriminating index of acceptance or rejection of an item was set at 3. All items with mean value of 3.00 and above were accepted as areas of information needs. The table shows that 8 items had mean scores of information on the environment which had a mean value of 3.06, things that constitute pollution had 3.08 mean score, drainage and irrigation had 3.45, erosion had 3.33, refuse disposal methods had 3.25, pesticide application and use had 3.49, environmental disaster management had 3.93 and was ranked first while funding sources for environmental management had 3.65. Also the result showed that 3 items were not accepted as areas of information needs since their mean scores were not up to 3. They included tree planting exercises, manure and maturing and cropping systems.

c) Effects of Environmental Conservation

The distribution of farmers according to their perceived effects of environmental conservation is presented in table 3. Using the discriminating index of 3, the result shows that 7 items are effects of environmental conservation. These included reduction of ill health and the predisposing factors with a mean score of 3.10, support for life had mean score of 3.03, reduction of hazards had 3.68 mean score. Whereas the statement, favours agricultural products by increasing income had 3.74 was ranked first. Agricultural produce are free from contamination had 3.09 and makes for odour free environment had 3.39. Only one item was not accepted as an effect of environmental conservation.

d) Farmers Conservation Practices

The result in table 4 below shows that among the conservation practices mentioned, manuring had a percentage count of 82.5 and ranked 1st, avoidance of bush burning had a percentage of 75.83 and ranked 2nd, mulching with percentage of 75 ranked 3rd, crop rotation with percentage of 66.67 raked 4th, adequate refuse disposal and cautious use of agro- chemicals and pesticides with percentages of 56.67 each ranked 5th,

tree planting ranked 7th with 55.83 percent. Whereas use of cross bars ranked 8th with 55 percent, ridging across the slop and drainage with percentage o 51.67 each ranked 9th. Rotational grazing ranked 10th with 35.83 percent, contour terracing 11th with a percentage of 34.17, use o contour bunds with 32.67 percent ranked 12th while irrigation and recycling of waste with percentage of 30 and 29.17 ranked 13th and 14th respectively.

e) Problems Militating against environmental conservation

The result shows that inadequate knowledge base of environmental conservation practices represented 75 percent to rank 1st. Following thus in rank (2nd) was poor economic base and inadequate farm size with 74.17 percent each. Poor environmental conservation orientation ranked 3rd with 69.17 percent, weak government policies and programme followed in rank (4th) with 47.5 percent. Poor pricing of agricultural produce with percentage count of 40 ranked 5th. A percentage count of 39.17 and 20 ranked 7th and 8th respectively. This means that the problems militating against environmental conservation are enormous and there are implications for extension services.

IV. CONCLUSION

Environmental degradation impacts negatively on soil fertility. Farmers need information on disaster magenament, Erosion, village practices, pollution, cropping systems and other area. They practice milling, rotation, and others to import their socio economic life, reduce hazards, increase yield and promote agro-biodiversity. Among the problems failing respondents include inadequate knowledge base of environmental conservation practices, poor economic base, small farm holdings, among others. Information provision to farmers on environmental conservation should be based on identified areas of needs like disaster management, fund chemical use.

REFERENCES RÉFÉRENCES REFERENCIAS

1. (2000). Imo State Agricultural Development Agency Report, Owerri.
2. Mgbada, J. V (2010). Agricultural Extension: the human Development Perspective. 298-299.
3. Mollet, J.A (1990). Planning for Agricultural Development (2nd Ed), England.
4. Nnadi, F.N and Onuoha, E.R. (1999.) fund amentals of Agricultural Extension and Rural Sociology.
5. Nnadi, F.N. and Amaechi, E.C.C.I (2007). Rural Sociology for development Studies, Imo State.
6. Olayide, S.O, Ogunfowora, S.M., Essang, S.M. and Idachaba, F.S (1981). *Element of Rural Economics*. Ibandan.
7. National census figure, (2006).
8. Environment and climate change. (2010).

9. Favis-Mortlock, D. (2005). Soil erosion patterns, spatio temporal dynamics and connectivity. 30, 131-132.
10. Frances, B. and Kathleen, S. (2002). Understanding knowledge management and information management-the need for an empirical perspective, 8.
11. Towery, D. and Werblow, S. (2010). Facilitating Conservation farming practices and Enhancing Environmental Conservation Technology Information center, United States America.

Table 1 : Socioeconomic characteristics of respondents

Variables	Frequency	Percentage
Age		
20-29	9	7.50
30-39	20	16.67
40-49	29	24.17
50-59	34	28.33
Sex	28	23.33
Male	59	49.17
Female	61	50.83
Marital Status		
Single	13	10.83
Married	83	70.83
Divorced	4	3.33
Widowed	18	15.0
Education		
No formal Education	22	18
Primary	37	31.09
Secondary	37	31.09
Tertiary	24	20.17
Occupation		
Farming	75	62.5
Civil servant	21	17.5
Trading	21	17.5
Artisan	3	2.5
Household size		
1-3	19	38
4-6	59	49.17
7-9	39	32.5
10 and above	3	2.5
Farm size		
Less than 1	59	49.17
1-3	35	29.17
4-6	21	17.5
7-9	5	4.16
Membership of social organization		
Not Belong	43	
Belong	77	64.17
Farming Experience		
1-5	16	13.33
6-10	43	35.83
11-15	14	11.67
16 and above	47	39

Table 2 : Distribution of the farmers according to areas of information needs

Areas	SA	A	U	D	SD	Mean	Rank
a) Environmental disaster management	245	96	114	16	1	3.93	1 st
b) Funding sources for environmental management	235	72	81	46	5	3.65	2 nd
c) Pesticide application and use	165	84	120	48	2	3.49	3 rd
d) Drainage and irrigation	210	64	72	60	8	3.45	4 th
e) Erosion	190	100	48	16	33	3.33	5 th
f) Refuse disposal methods Tillage practices	95	112	156	12	15	3.25	6 th
g) Things that constitute pollution	145	48	87	80	10	3.08	7 th
h) Sources of information on the environment/practices	180	56	42	68	22	3.06	8 th
i) Tillage practices	75	92	81	72	19	2.82	9 th
j) Tree planting exercise	60	60	135	64	16	2.79	10 th
k) Cropping system	60	60	69	84	28	2.50	11 th
l) Manure and manuring	75	28	51	128	17	2.49	12 th

Source: field survey data, 2011.

Table 3 : Perceived effects of environment conservation

Effects	SA	A	U	D	SD	Mean	Rank
a) Improves the farmers socio-economic life by increasing income	230	68	126	20	5	3.74	1 st
b) Reduction of hards-injury, harm and accident.	190	108	105	36	2	3.68	2 nd
c) Favours agricultural products by increasing yield	200	100	69	58	3	3.58	3 rd
d) Makes for odour/pollution free environment	165	60	135	40	7	3.39	4 th
e) Reduction of ill health and the predisposing factors	160	80	51	64	19	3.12	5 th
f) Improves gaseous exchange	120	36	144	58	10	3.10	6 th
g) Agricultural produce are free from contamination	155	36	99	68	13	3.09	7 th
h) Support for life	160	84	54	32	33	3.03	8 th
i) Increases longevity	115	20	102	84	16	2.82	9 th

Source: field survey data

Table 4 : Farmers conservation practices

Practices	frequency	percentage	Rank
a. Manuring	99	82.5	1 st
b. Avoidance of bush burning	91	75.83	2 nd
c. Mulching	90	75	3 rd
d. Crop rotation	80	66.67	4 th
e. Adequate refuse disposal	68	56.67	5 th
f. Cautious use of agro- chemicals And pesticides	68	56.67	6 th
g. Tree planting	67	55.83	7 th
h. Use of cross bars	66	55	8 th
i. Ridging across the slope	62	51.67	9 th
j. Drainage	62	51.67	10 th
k. Rotational grazing	43	35.83	11 th
l. Contour terracing	41	34.17	12 th
m. Use of contour bunds	38	31.67	13 th
n. Irrigation	36	30	14 th
o. Recycling of waste	35	29.17	15 th

Source : Field survey data, 2011.

Table 5 : Problems militating against environmental conservation

Practices	frequency	percentage	Rank
a. Inadequate knowledge base of environmental conservation practices	90	75	1 st
b. Poor economic base	89	74.15	2 nd
c. Inadequate farm size	89	74.15	3 rd
d. Poor environmental conservation	83	69.17	4 th
e. Weak government policies and programmes			
f. Poor pricing of agricultural produce	54	40	6 th
g. Cumbersome nature of the practices	47	39.17	7 th
h. Poor linkages between and among the major stakeholders	29	24.17	8 th
i. Religious believe	24	20	9 th

Source: field survey data, 2011.

This page is intentionally left blank



Evaluation of Quality Protein Maize (QPM) and Normal Maize for Growth Performance of Broiler Chicken in Nepal

By M.R. Tiwari, D. Neopane, T.P. Paudel & U.M. Singh

National Animal Science Research Institute, Nepal

Abstract - An experiment was carried out on Cob 500 broiler chickens at Swine and Avian Research Program, Khumaltar, Lalitpur to evaluate the growth performance of broiler chicken fed with normal Vs QPM based diet and with supplementation of synthetic lysine and methionine in both maize for 54 days (16 days starter and 38 days for finisher). The experimental day old birds were procured from Valley Feed, Bajalu, Kathmandu and were allotted into four treatments with three replications having 15 birds in each replication by using Complete Randomized Design (CRD). All experimental birds were vaccinated with F₁ vaccine @ one drop / bird against Ranikhet at the first week and vaccinated with IB+ND vaccine against Gumboro at 2nd and 5th weeks of experiment. Birds of T₁ were provided normal maize based diet, T₂ normal maize based diet with supplementation of synthetic lysine and methionine, T₃ QPM based diet and T₄ QPM based diet with supplementation of synthetic lysine and methionine. Concentrate mixture feeding was done on group basis and was provided to the experimental birds of all groups once a day (morning) in adlib amount for both periods (starter – 16 days and finisher – 38 days) of the experiment. Quantity of concentrate mixture given daily to the birds in groups weighed daily and refusal was weighed in the next morning.

Keywords : QPM feeding, synthetic lysine and methionine, broiler chicken, Nepal.

GJSFR-D Classification : FOR Code: 820401, 070799



EVALUATION OF QUALITY PROTEIN MAIZE QPM AND NORMAL MAIZE FOR GROWTH PERFORMANCE OF BROILER CHICKEN IN NEPAL

Strictly as per the compliance and regulations of :



RESEARCH | DIVERSITY | ETHICS

Evaluation of Quality Protein Maize (QPM) and Normal Maize for Growth Performance of Broiler Chicken in Nepal

M.R. Tiwari^α, D. Neopane^ο, T.P. Paudel^ρ & U.M. Singh^ω

Abstract - An experiment was carried out on Cob 500 broiler chickens at Swine and Avian Research Program, Khumaltar, Lalitpur to evaluate the growth performance of broiler chicken fed with normal Vs QPM based diet and with supplementation of synthetic lysine and methionine in both maize for 54 days (16 days starter and 38 days for finisher). The experimental day old birds were procured from Valley Feed, Bajalu, Kathmandu and were allotted into four treatments with three replications having 15 birds in each replication by using Complete Randomized Design (CRD). All experimental birds were vaccinated with F1 vaccine @ one drop / bird against Ranikhet at the first week and vaccinated with IB+ND vaccine against Gumbaro at 2nd and 5th weeks of experiment. Birds of T₁ were provided normal maize based diet, T₂ normal maize based diet with supplementation of synthetic lysine and methionine, T₃ QPM based diet and T₄ QPM based diet with supplementation of synthetic lysine and methionine. Concentrate mixture feeding was done on group basis and was provided to the experimental birds of all groups once a day (morning) in adlib amount for both periods (starter – 16 days and finisher – 38 days) of the experiment. Quantity of concentrate mixture given daily to the birds in groups weighed daily and refusal was weighed in the next morning. Experiment revealed that cumulative feed intake was found to be higher in QPM based diet group (5597 g) followed by QPM based diet with supplementation of synthetic lysine and methionine group (5403 g) and normal maize based diet with supplementation with synthetic lysine and methionine group (5377 g). The least feed consumption was recorded for normal maize based diet group (4937 g) which was none significant among diet groups. Similarly, total weight gain was found to be higher for QPM based diet with supplementation of synthetic lysine and methionine group (2180 g) followed by normal maize based diet with supplementation with synthetic lysine and methionine group (2141 g) and QPM based diet group (1996 g). The least weight gain was note in normal maize based diet group (1854 g) which was also none significant among diet groups.

Keywords : QPM feeding, synthetic lysine and methionine, broiler chicken, Nepal.

I. INTRODUCTION

Broiler is raised specifically for meat production. Typical broilers have white feathers and yellowish skin. Most commercial broilers bred for meat reach slaughter weight at between 5 to 7 weeks of age,

although slower growing strains reach slaughter weight at approximately 14 weeks of age. Broiler chickens are most efficient converters of feed into animal protein in comparison to other domestic birds. The broiler farming need less capital, less space and give quick and early returns because of shorter generation intervals. In Nepal, there is high demand of broiler meat in the markets due to low price and remunerative value in comparison to other type of meat poultry industry. There are 39.5 million chickens in the Nepal (MoAC 2010/11). More than 60% of the population belongs to indigenous and rest is exotic origin.

Maize has become the world's chief animal feed. It provides more feed than any other grain. It is outstanding being high in energy, low in fibre and easily digested by most livestock species. Maize has remained a critical feed ingredient in monogastric diets particularly poultry. About 70% of maize produced worldwide is utilized in livestock feed. The normal maize varieties are low in lysine and tryptophan contents (NRC 1988 and FAO 2004).

QPM is a nutritionally superior cereal grain that possesses a higher proportion of two key amino acids, lysine and tryptophan, than is found in normal maize. QPM has amply demonstrated its superior performance as feed for monogastric animals, especially pigs and chickens. Chickens and pigs convert QPM feed into weight gain more efficiently than regular maize feed (Jarkin *et al.* 1970; Maner 1975 and Asche *et al.* 1985). The percentage of lysine content in QPM varied between 0.33 and 0.54 with an average of 0.38. This was 46 percent higher than normal maize, and QPM contained 66 percent more tryptophan (0.08%) than normal maize. These two amino acids allow the body to manufacture complete proteins (Palit and Babu 2003). Therefore, utilization of QPM can correct this deficiency and may be advantageous in the diets of monogastric animals (Hai *et al.* 2010).

Several research works conducted by different researchers around the world reported that broiler chicken fed with QPM was much better over normal maize fed broiler; however, feeding value of QPM has not been evaluated in poultry feed in Nepal. Therefore, an attempt was made to evaluate the feeding value of normal maize and QPM with or without supplementation of synthetic lysine and methionine on broiler chickens at

Authors ^α ^ω : National Animal Science Research Institute, Khumaltar, Lalitpur. E-mail : tiwari65@yahoo.com

Authors ^ο ^ρ : Swine and Avian Research Program, Khumaltar, Lalitpur.

Swine and Avian Research Program, Khumaltar, Lalitpur.

Ranikhet at the first week and vaccinated with IB+ND vaccine against Gumbo at 2nd and 5th weeks of experiment.

II. MATERIALS AND METHODS

a) Experimental Animals

The experiment was carried out on Cob 500 broiler chickens at Swine and Avian Research Program, Khumaltar, Lalitpur from 14 March to 6 May 2013 (069/12/1 to 070/1/23 BS) for 54 days (16 days starter and 38 days finisher). The experimental day old birds were procured from Valley Feed, Bajalu, Kathmandu and were allotted into four treatments with three replications having 15 birds in each replication by using Complete Randomized Design (CRD). All experimental birds were vaccinated with F1 vaccine @ one drop /bird against

b) Diet Composition

Feed ingredients such as normal maize, soybean cake, mustard cake, rice bran, mineral mixture, methionine, lysine, oil, DCP and salt were procured from "Champadevi Feed Industries", Chapagau, Lalitpur while QPM was bought from National Maize Research Program, Rampur, Chitawan and wheat grain from Agriculture Botany Division, Khumaltar, Lalitpur. Four types of concentrate mixture were composed for experimental birds; starter (22% CP) and finisher (18% CP) which are presented in Table 1, 2, 3 and 4.

Table 1 : Starter ration for 1-15 days for T₁ and T₃ group

Ingredient	Part	CP	ME Kcal	Methionine	Lysine	Ca	P
Normal Maize / QPM	46	3.68	1351.02	0.041	0.07	0.12	0.11
Soybean meal	15	6.01	307.05	0.087	0.36	0.04	0.09
Rice bran	10	1.08	293.70	0.020	0.45	0.01	0.12
Meat and bone meal	13	7.21	267.96	0.486	0.12	1.34	0.66
Oil	0.15	0.00	1155.00	0.000	0.00	0.00	0
Mineral	0.65	0.00	0.00	0.000	0.00	0.00	0
Salt	0.7	0.00	0.00	0.000	0.00	0.00	0
Mustard cake	13	4.05	277.68	0.089	0.02	0.00	0.09
Wheat grain	0	0.00	0.00	0.000	0.00	0.00	0
DCP	1.5	0.00	0.00	0.000	0.00	0.35	0.25
Total	100	22.03	3652.41	0.72	1.02	1.86	0.66

Table 2 : Starter ration for 1 - 15 days for T₂ and T₄ group

Ingredient	Part	CP	ME Kcal	Methionine	Lysine	Ca	P
Normal Maize/ QPM	46	3.68	1351.02	0.04	0.07	0.12	0.11
Soybean meal	15	6.01	307.05	0.09	0.36	0.04	0.09
Rice bran	10	1.08	293.70	0.02	0.45	0.01	0.12
Meat and bone meal	13	7.21	267.96	0.49	0.12	1.33	0.66
Oil	0.15	0.00	1155.00	0.00	0.00	0.00	0
Mineral	0.65	0.00	0.00	0.00	0.00	0.00	0
Salt	0.5	0.00	0.00	0.00	0.00	0.00	0
Mustard cake	13	4.05	277.68	0.09	0.02	0.00	0.09
Wheat grain	0	0.00	0.00	0.00	0.00	0.00	0
Lysine	0.1	0.00	0.00	0.00	0.10	0.00	0
Methionine	0.1	0.00	0.00	0.10	0.00	0.00	0
DCP	1.5	0.00	0.00	0.00	0.00	0.35	0.25
Total	100	22.03	3652.41	0.82	1.12	1.85	0.66

Table 3 : Finisher ration for 16 - 54 days for T₁ and T₃ group

Ingredient	Part	CP	ME Kcal	Methionine	Lysine	Ca	P
Normal maize / QPM	40	3.20	1175	0.04	0.08	0.10	0.10
Soybean meal	10	4.01	205	0.06	0.24	0.03	0.06
Rice bran	10	1.08	294	0.02	0.45	0.01	0.12
Meat and bone meal	9	4.99	186	0.34	0.08	0.93	0.46
Oil	0.15	0.00	1155	0.00	0.00	0.00	0.00
Mineral	1.15	0.00	0	0.00	0.00	0.00	0.00
Salt	1.2	0.00	0	0.00	0.00	0.00	0.00

Mustard cake	11	3.43	235	0.08	0.21	0.00	0.08
Wheat grain	16	2.10	59	1.09	0.51	0.02	0.04
DCP	1.5	0	0	0	0.00	0.35	0.25
Total	100	18.80	3307	1.62	1.56	1.44	1.10

Table 4 : Finisher ration for 16 - 54 days for T₂ and T₄ group

Ingredient	Part	CP	ME Kcal	Methionine	Lysine	Ca	P
Normal maize / QPM	40	3.20	1175	0.04	0.08	0.10	0.10
Soybean meal	10	4.01	205	0.06	0.24	0.03	0.06
Rice bran	10	1.08	294	0.02	0.45	0.01	0.12
Meat and bone meal	9	4.99	186	0.34	0.08	0.93	0.46
Oil	0.15	0.00	1155	0.00	0.00	0.00	0
Mineral	1.15	0.00	0	0.00	0.00	0.00	0
Salt	1	0.00	0	0.00	0.00	0.00	0
Mustard cake	11	3.43	235	0.08	0.21	0.00	0.08
Wheat grain	16	2.10	59	1.09	0.51	0.02	0.04
Lysine	0.1	0.00	0	0.00	0.10	0.00	0
Tryptophan	0.1	0.00	0	0.10	0.00	0.00	0
DCP	1.5	0.00	0	0.00	0.00	0.35	0.25
Total	100	18.80	3307	1.72	1.66	1.44	1.10

c) *Experimental Diet*

Following diets were formulated to the experimental broiler chickens (Table 5)

Table 5 : Experimental diets of the broiler

Treatment	Experimental diets
1	Normal maize included concentrate mixture
2	Normal maize included concentrate mixture + synthetic lysine and methionine
3	QPM included concentrate mixture
4	QPM included concentrate mixture + synthetic lysine and methionine

d) *Feeding Regime*

Concentrate mixture feeding was done on group basis and was provided to the experimental birds of all groups once a day (morning) in *adlib* amount for both periods (starter – 16 days and finisher – 38 days) of the experiment. Quantity of concentrate mixture given daily to the birds in groups weighed daily and refusal was weighed in the next morning. Drinking water was provided in adequate amount.

e) *Chemical Analysis*

The samples of feed ingredients were sent to the Animal Nutrition Division, Khumaltar, Lalitpur for proximate analysis. Representative samples from offered concentrate mixture were analyzed for Dry Matter (DM), Crude Protein (CP), Crude Fibre (CF), Ether Extract (EE) and Ash contents (TA). The DM was determined by oven drying at 100°C for 24 hrs. Crude protein of the samples was determined using the Kjeldahl method. Ether extract was determined using Soxhlet apparatus. Ash content was determined by ashing at 550°C in a muffle furnace for 16 hrs (AOAC 1980). Crude Ether of the samples was determined using the Van Soest method (Goering, H.K. and Van Soest 1970). Similarly, samples of normal maize and QPM were sent to Food Research Division, Khumaltar, Lalitpur for tryptophan and lysine content analysis.

Tryptophan was analyzed as suggested by Hornandez H. and Bates L.S. (1969) and Lysine as suggested by Doll H. and Koie B. (1975).

f) *Data Measurement*

The trial period consisted for 54 days (16 days starter and 38 days finisher). Total feed intake by the experimental birds in the group was recorded daily for both experimental periods. The body weight gain was measured in group basis (replication-wise) in seven days interval in the morning before feeding.

g) *Data Analysis*

Data of feed intake and body weight gain were analyzed by "*One way Anova*" test for every measurement using statistical package Minitab 2003, versions 13.20

III. RESULTS AND DISCUSSION

a) *Chemical Composition of feed ingredients*

The result of chemical analysis and amino acids content are given in Table 6.

Table 6 : Chemical composition of feed ingredients (% DM basis)

Ingredients	DM	OM	TA	CP	CF	EE
Normal maize	87.69	97.97	2.03	9.0	2.34	4.48
QPM	89.36	97.62	2.38	9.0	6.26	5.12
Rice bran	87.85	89.5	10.5	10.0	4.83	5.1
Mustard cake	87.27	90.5	9.5	35.0	11.23	NA
Soybean cake	86.87	92.63	7.37	46.29	9.38	0.7
Wheat grain	93.0	91.3	8.7	14.0	8.45	NA
Meat cum bone meal	93.22	67.2	32.8	49.93	3.43	NA

Comparison of nutrient content of QPM and normal maize is given in Table 7

Table 7 : Comparison of the nutritional composition of QPM and Normal maize (on dry basis)

Nutrient	Ortega, <i>et al.</i> 1986		Osei <i>et al.</i> 1999		Our analysis 2012	
	Normal maize	QPM	Normal maize	QPM	Normal maize	QPM
Crude Protein, %	9.8	9.8	8.92	9.11	9	9
Ether Extract, %	NA	NA	4.48	5.12	4.48	5.12
Crude Fibre, %	NA	NA	1.93	2.14	2.34	6.26
Organic matter	NA	NA	98.10	98.40	97.97	97.62
Ash, %	NA	NA	1.90	1.60	2.03	2.38
Lysine, %	0.27	0.43	0.24	0.32	0.19	0.28
Tryptophan, %	0.06	0.10	0.06	0.08	0.04	0.07

b) Feed Intake

The average feed intake of experimental birds has been presented in Table 8.

Average feed intakes of experimental birds were observed 17.05 g in 7 days which reached 5328 g by the end of experiment (54 days) which was none significant among diet groups. Feed intake of QPM based diet group (T₃) was higher than that of normal maize based diet group (T₁) from beginning to the end of experiment which was highly significant (P<0.001) in 22 days and 29 days (P<0.001) of experiment while rest

of period was none significant. Supplementation of synthetic lysine and methionine in QPM based diet group (T₄) performed higher feed intake in 36 days (124.4 g) and 43 days (151.96 g) of experiment only whereas in other periods normal maize based diet with supplementation of synthetic lysine and methionine groups (T₂) had more feed intake. The cumulative feed intake was found to be higher in T₃ (5597 g) followed by T₄ (5403 g) and T₂ (5377 g). The least feed consumption was recorded for T₁ group (4937 g) which was none significant among diet groups.

Table 8 : Feed intake of experimental birds, g (Mean ± SD)

TRT	Days								Cumulative feed intake
	7	15	22	29	36	43	50	54	
1	16.23±0.17	36.03±1.64	64.69±0.57	90.88±2.02	107.21±10.23	132.73±9.02	148.30±17.91	153.18±5.81	4937±0.3
2	17.38±1.19	39.50±0.74	77.72±1.22	110.95±4.1	118.66±6.46	121.6±24.65	163.55±29.9	166.53±14.92	5377±0.52
3	17.28±0.83	37.45±0.60	70.40±0.67	107.07±3.43	121.05±11.43	144.73±2.22	177.87±4.34	173.29±4.19	5597±0.11
4	17.32±0.55	37.71±1.81	70.90±2.09	103.75±2.38	126.4±6.88	151.96±6.24	158.25±21.20	148.46±14.32	5403±0.28
Mean	17.05±0.83	37.67±1.07	70.93±4.94	103.16±8.3	118.33±10.61	137.76±16.72	161.99±20.74	160.37±13.99	5328±0.38
P- value	P>0.05	P>0.05	P<0.001	P<0.001	P>0.05	P>0.05	P>0.05	P>0.05	P>0.05

c) Growth Performance

The growth trend of experimental birds is presented in Table 9 and figure 1

Table 9 showed that initial body weight of experimental birds was 46 g, 45 g, 46 g and 45 g for T₁, T₂, T₃ and T₄, respectively which reached 1854 g, 2141

g, 1996 g and 2180 g at the end of experiment (54 days). Both initial and final body weights were none significant among the diet groups. Body weight gain of QPM fed group (T₃) was found to be higher from beginning to the end of experiment than that of normal maize fed group (T₁), however, significant effect of QPM feeding

was observed from 22 to 50 days of experiment while 7, 15 and 54 days of feeding was none significant. Supplementation of synthetic lysine and methionine in normal maize based diet group (T₂) performed higher body weigh gain from 7 to 36 days of experiment than that of QPM based diet with supplementation of synthetic lysine and methionine (T₄). Thatfater, from 43 to 54 days of experiment QPM based diet with supplementation of synthetic lysine and methionine (T₄) performed better in body weight gain. Effect of synthetic

lysine and methionine supplementation in normal maize and QPM based diet on body weight gain of experimental birds was significant from 22 to 50 days of experiment while 7, 15 and 54 days of feeding was none significant. The total weight gain was found to be higher for T₄ (2180 g) followed by T₂ (2141 g) and T₃ (1996 g). The least weight gain was note in T₁ (1854 g) which was also none significant among diet groups.

Table 9 : Body weight gain of experimental birds, g (Mean± SD)

TRT	Days									
	0	7	15	22	29	36	43	50	54	Total weight gain
1	46±0	114±0	250±0.01	479±0.01	755±0.02	957±0.06	1228±0.07	1305±0.12	1928±0.04	1854±0.06
2	45±0	120±0	271±0	560±0.01	915±0.03	1199±0.01	1465±0.09	1944±0.12	2204±0.08	2141±0.08
3	46±0	115±0	252±0.01	499±0	811±0.01	982±0.03	1347±0.07	1883±0.28	2061±0.07	1996±0.08
4	45±0	112±0	249±0.01	528±0.03	866±0.01	1184±0.08	1547±0.11	2021±0.23	2233±0.32	2180±0.32
Mean	46±0	115±0	255±0.01	517±0.03	837±0.06	1080±0.12	1396±0.14	1788±0.34	2107±0.19	2042.75±0.14
P-Value	P>0.05	P>0.05	P>0.05	P<0.01	P<0.001	P<0.01	P<0.05	P<0.05	P>0.05	P>0.05




Figure 1 : Body weight gain trend of experimental birds

IV. DISCUSSION

The objective of this study was to evaluate the growth performance and feed intake of broiler chickens fed with normal maize and QPM with or without supplementation of synthetic lysine and methionine in diets. Experiment revealed that there was no significant

effect on total body weight gain and cumulative feed intake among normal maize and QPM diet group with or without supplementation of synthetic amino acids. It might be due to the variation in nutrient composition among the QPM cultivars.

Tyagi *et al.* (2008) did not find any significant difference with respect to body weight gain, feed

conversion and nutrient utilization efficiencies by feeding QPM diet to broiler chickens compared to those fed the normal maize diet. These variations noticed on the performance could be attributed to variation in nutrient composition among the QPM cultivars used in different studies. Bai (2002) reported that there was no significant difference between normal maize and QPM feeding to carcass percentage, abdominal fat percentage, percentage of eviscerated yield and percentage of eviscerated yield with giblets. At a given digestible lysine content, using QPM tended to increase weight gain but there was no statistical evidence to support this ($P > 0.10$). Groote (2000) conducted an experiment by feeding normal maize and QPM to broiler chicken and noted that feed intake of both groups increased from 18.7 grams/bird/day in the first week, to 179.0 g/bird/day in the sixth week and tests showed no significant differences between the two diets (at the 5% level). Total feed intake over the six weeks was 3.38 kg/bird for the broilers on either diet, again with no significant difference between the two diets ($F = 0.265$, $p = 0.994$). Weight gain over the six weeks averaged 36.17 grams/ bird/day for broilers fed regular maize and 36.19 g/bird/ day for those fed on QPM diets and no statistical differences were found. Feed conversion efficiency, calculated as the weight gain over the feed intake, was 0.33 for both diets. In another experiment, he observed that dietary substitution of normal maize with QPM did not have any effect on body weight gain and feed consumption. However, feed conversion ratio was significantly ($P < 0.01$) improved by substituting normal maize by QPM or supplementing lysine to normal maize based diet.

Nevertheless, Onimisi *et al.* (2009) reported that QPM in poultry diet improved growth performance of broilers and resulted into higher weight gains than normal maize. QPM did not result into significant changes on carcass and organ development of broilers. Amonelo and Roxas (2008) reported that broilers fed with either normal maize or QPM based diet had higher ($P < 0.05$) live weight gain compared to those fed diets supplemented with synthetic lysine. Broilers fed either the QPM based diet or those supplemented with synthetic lysine had lower ($P < 0.05$) feed consumption compared to broilers fed with normal maize diet. Broilers fed with QPM based diet had the best performance ($P < 0.05$) in terms of feed efficiency. Ose *et al.* (2002) Birds receiving QPM as the sole source of amino acids performed significantly better ($P < 0.05$) than their counterparts fed on normal maize. QPM fed birds weighed an average 708.0 g each at the end of the trials compared with 532.0 g for those on normal maize. The corresponding feed efficiencies were 4.28 and 6.55, respectively. Compared with birds on the balanced diet, however, QPM was inadequate in supporting broiler growth.

V. CONCLUSION

This experiment revealed that quality protein maize varieties that are available in Nepal are to be crossed with local varieties due to which content of essential amino acids such as lysine and tryptophan is reduced than as mentioned in literatures. Therefore, QPM could not exhibit better performance over normal maize on total weight gain and cumulative feed intake of broiler chickens.

VI. ACKNOWLEDGEMENT

The authors would like to express their most sincere gratitude and appreciation to Dr. Bhoj Raj Joshi (Director) National Animal Science Research Institute, Khumaltar, Lalitpur for his encouragement, help, suggestion and support during experimental period. Mr. Man Bahadur Shrestha (Chief) Food Research Division, Khumaltar, Lalitpur deserves due acknowledgement for his tremendous work in lysine and methionine analysis. Similarly, work of Mr. Raj Kumar Raut, Mr. Kaji Ratna Maharjan and Shyam Khatri (Technicians of Swine and Avian Research Program, Khumaltar, Lalitpur) deserve high appreciation without which conduction of experiment was impossible. Mr. Basanta Kumar Shrestha Technical Officer, Animal Nutrition Division, Khumaltar also deserve appreciation for his dedicated work in chemical analysis of fed ingredients. Likewise, our thanks also go to the all-technical, admin and finance staff of National Animal Science Research Institute, Khumaltar for their help during trial period.

REFERENCES RÉFÉRENCES REFERENCIAS

1. Asche, G.L., A.J Lewis, E.R. Peo, and J.D. Greenshaw (1985). The nutritional value of normal and high lysine corns for weanling and growing-finishing swine when fed at four lysine levels. *Journal of Animal Science*, 60, 1412-1428.
2. (1980). *Association of Official Analysis Chemists*, Official methods of analysis, U.S.A .
3. Amonelo M. O. and D. B. Roxas (2008). Growth performance of broilers fed quality protein maize (qpm) based diet. 34 (1), 11-22.
4. Bai, Xue-Feng (2002). Nutritional evaluation and utilization of quality protein maize Zhong Dan 9409 in broiler feed.
5. Doll, H. and B. Koie (1975). Evaluation of high lysine maize mutants. 55-59.
6. (2004). *Protein Sources for the Animal Feeds Industry*. FAO Expert Consultation Workshop Report.
7. Goering, H.K. and Van Soest (1970). Forage fibre analysis apparatus, reagents, procedures and some application, ARS, USDA.
8. Groote H. D., T. Nyanamba and R. Wahome (2000). Quality protein maize for the feed industry in Kenya. www.researchgate.net.

9. Hornadez, H. and L.S. Bates (1969). A modified method for rapid tryptophan analysis of maize.
10. Hai, G., Q.Y. Diao and S.H. Zhang (2010). Nutritional evaluation and utilization of quality protein maize in animal feed.
11. Jarkin, R., C. Albertazzi and R. Bressani (1970). Value of Opaque-2 corn protein for chicks. *Journal of Agricultural and Food Chemistry*, 18, 268-272.
12. Maner, J.H. (1975). Quality protein maize in swine production. In High Quality Protein Maize. 51-82.
13. (2010/11). *Ministry of Agriculture and Cooperatives, Singha Durbar, Kathmandu, Nepal NRC*. Statistical Information on Nepalese Agriculture., USA.
14. Ortega, E. I., E. Villegas and S. K. Vasal (1986). A comparative study of protein changes in normal and quality protein maize during tortilla making. 63 (5), 446-451.
15. Osei, S. A., H. K. Dei and A. K. Tuah (1999). Evaluation of quality protein maize as a feed ingredient for layer pullet. *Journal of Animal*., 181-189.
16. Onimisi, P.A., J.J. Omenge, I.I. Dafwang and G.S. Bawa (2009). Replacement Value of Normal Maize with Quality Protein Maize (Obatampa) in Broiler Diets. *Pakistan Journal of Nutrition*, 8 (2), 112-115.
17. Osei S. A., C. C. Atuahene, D. B. Okai, A. Donkoh and A. K. Tuah (2002). *The nutritive value of quality protein maize in the diets of broiler chickens*. Department of Animal Science, Kumasi Ghana.
18. Palit, K.K. and S.C. Babu (2003). Food systems for improved human nutrition: linking agriculture, nutrition and productivity.
19. Tyagi, P. K., A. K. Shrivastav, A. B. Mandal, Promod K. Tyagi, A.V. Elangovan and C. Deo (2008). The apparent metabolizable energy and feeding value of quality protein maize for broiler chicken. 43, 169-174.





This page is intentionally left blank



Effect of Heat Treated Soybean Cake Feeding on Growth Performnce of Growing Female Goats in Fodder based Basal Diet

By M.R. Tiwari, D. Adhikary, D.P. Adhikary, R.P. Ghimire & S.H. Ghimire

Nepal Agricultural Research Council, Nepal

Abstract - Growth comparison of goats fed with treated and none treated soybean cake is not evaluated so far in Nepal. Therefore, an experiment was carried out on eighteen growing female goats (50% Jamunapari 6, 50% Barberi 6 and Kiko goats 6) at the Agriculture Research Station (Goat), Bandipur for 90 days after an adaptation period of 7 days. Female goats of average five months age having body weight 11.86 kg were allocated into three groups having six animals in each group by using Complete Randomized Design (CRD). For T₁ and T₂ concentrate mixture were composed by using procured feed ingredients with 16% crude protein level while T₃ was fed with commercial feed. Experimental animals of T₁ group was provided forest mixed fodder (adlib) + treated soybean cake included concentrate mixture @ 1.5% of body weight, T₂ group was provided forest mixed fodder (adlib) + untreated soybean cake included concentrate mixture @ 1.5% of body weight whereas T₃ was provided forest mixed fodder (adlib) + commercial concentrate mixture @ 1.5% of body weight. Experiment revealed that higher intake of concentrate feed was recorded for T₁ (207.6 g) followed by T₃ (199.58 g) and T₂ (193.87 g) which was highly significant (P<0.001) among diet groups.

Keywords : goats, bypass protein feeding.

GJSFR-D Classification : FOR Code: 070799, 070204



EFFECT OF HEAT TREATED SOYBEAN CAKE FEEDING ON GROWTH PERFORMNCE OF GROWING FEMALE GOATS IN FODDER BASED BASAL DIET

Strictly as per the compliance and regulations of :



RESEARCH | DIVERSITY | ETHICS

Effect of Heat Treated Soybean Cake Feeding on Growth Performance of Growing Female Goats in Fodder based Basal Diet

M.R. Tiwari^α, D. Adhikary^σ, D.P. Adhikary^ρ, R.P. Ghimire^ω & S.H. Ghimire[¥]

Abstract - Growth comparison of goats fed with treated and none treated soybean cake is not evaluated so far in Nepal. Therefore, an experiment was carried out on eighteen growing female goats (50% Jamunapari 6, 50% Barberi 6 and Kiko goats 6) at the Agriculture Research Station (Goat), Bandipur for 90 days after an adaptation period of 7 days. Female goats of average five months age having body weight 11.86 kg were allocated into three groups having six animals in each group by using Complete Randomized Design (CRD). For T1 and T2 concentrate mixture were composed by using procured feed ingredients with 16% crude protein level while T3 was fed with commercial feed. Experimental animals of T1 group was provided forest mixed fodder (adlib) + treated soybean cake included concentrate mixture @ 1.5% of body weight, T2 group was provided forest mixed fodder (adlib) + untreated soybean cake included concentrate mixture @ 1.5% of body weight whereas T3 was provided forest mixed fodder (adlib) + commercial concentrate mixture @ 1.5% of body weight. Experiment revealed that higher intake of concentrate feed was recorded for T1 (207.6 g) followed by T3 (199.58 g) and T2 (193.87 g) which was highly significant ($P < 0.001$) among diet groups. Similarly, fodder intake was also noted significantly higher ($P < 0.001$) among diet groups (1969.5 g, 1967.6 g and 1942 g for T2, T3 and T1, respectively). Feed and fodder intake of different genotypes of goats was found to be non-significant among goat breeds. In addition, feed conversion ratio per kg body weight gain was observed higher for T3 (22.49:1) followed by T2 (17.57:1) and T1 (16.24:1). Similarly, initial body weight of T1, T2 and T3 was 12.15 kg, 11.25 kg and 12.18 kg respectively that reached 17.66 kg, 16.33 kg and 16.40 kg during 90 days of experiment for T1, T2 and T3, respectively. Both initial and final body weight was non-significant among diet groups. Similarly, there was also non-significant effect of goat breed on body weight gain. Total body weight gain was recorded higher for T1 (5.50 kg) followed by T2 (5.08 kg) and T3 (3.98 kg) which was significant ($P < 0.05$) among diet groups. Similarly, average daily gain was also noted higher in T1 (61.2g) with variation of 29-122.6g followed by T2 (56.48 g) with variation of 21-102.6 g and T3 (44.22 g) with variation of 14-101.3 g.

Keywords : goats, bypass protein feeding.

I. INTRODUCTION

Goats are important domestic animals in many parts of the world. In the developing countries, goats make a very valuable contribution,

Author α : National Animal Science Research Institute (NASRI), Khumaltar, Lalitpur. E-mail : tiwari65@yahoo.com

Authors σ ρ ω \textyen : Agriculture Research Station (Goat), Bandipur, Tanahun.

especially to the poor in the rural areas. The importance of this valuable genetic resource is underestimated and its extent of contribution to the livelihood of the poor is inadequately understood. Goats are efficient browsers and prefer eating brushy plants along with some other weedy plants found on the ranges. Nevertheless, goats are going to be more important source of livelihood for many more people in coming years and, thus, they deserve greater attention at both the micro and macro levels. Now, it is the time to consider and pay attention to the value and capacity of goats for producing food (Aziz 2010). Neopane and Pokharel (2008) reported that most of the farmers of western hills are rearing Khari goats, crossbred of Khari x Jamunapari and Khari and Barberi). Goat population of Nepal is estimated to be 9.19 million. Out of 9.19 million, goat population of western hills is 1.13 million which account 12.32% of total goat population that producing 5284 mt meat per annum (10% of total goat meat production) (MoAD 2012).

Oil seeds cakes and meals are the residues remaining after removal of the greater part of the oil from oil seeds. The residues are rich in protein and most are valuable feeds for farm animals. Most oil seeds are of tropical origin, they include groundnut, cottonseed, soya bean, mustard, til etc (Bajilieh 2002). Soybean cake has an average protein content of 40% and oil content of 20% and 2500 to 2800 kcal of metabolizable energy per kilogram. It also has a superior amino acid profile. Soybean protein has great potential as a major source of dietary protein. A by-product from the oil production (soybean cake) is used as a high-protein animal feed in many countries. Moreover, soybean protein is rich in lysine, methionine, valine, and isoleucine. The protein content of soybean tend to be 75 to 80% degraded in the rumen (Broderick *et al.* 1988; Promkot and Wanapat 2003), which restricts its inclusion in diets for high-yielding ruminants.

The protein content in diets of ruminant animals is essential for growth and production requirements. Possibility that reasonable portions of high quality protein of feedstuffs may be degraded in the rumen is occurred, which negatively affect animal utilization of the feed. In this context, there are several methods for protection of dietary protein from degradation in the rumen (EL-Shabrawy 1996). The heat treatment is

known one of the methods to increase the protection of the proteins. During the process of manufacturing oil seed meals, they are subjected to different degree of heating which partly explains differences in the degree of protection. Thorough heating of protein supplement causes denaturation of protein; it provides effective protection against microbial fermentation in the rumen. Heat treatment of protein meal at 125- 150^o C for 2-4 hours improves the bypass protein. The main benefit of "bypass" protein is that the original amino acids in the protein meal are absorbed in the small intestine instead of converted into microbial protein in the rumen, thereby providing a different balance of essential amino acids for better animal nutrition hence, production (Schroeder 1997).

Growth comparison of goats fed with treated and none treated soybean cake is not evaluated so far in Nepal. Hence, a study was carried out to compare the growth performance of growing female goats fed with heat-treated and none treated soybean meal mixed concentrate mixture at Agriculture Research Station (Goat), Bandipur, Tanahun.

II. METHODOLOGY

a) Experimental Animals

This experiment was carried out on eighteen growing female goats (50% Jamunapari 6, 50% Barberi

6 and Kiko goats 6) at Agriculture Research Station (Goat), Bandipur, Tanahun from 29 November 2012 to 25 March 2013 (069/8/14 to 069/11/14). Female goats of average five months old with average body weight of 11.86 kg were allocated into three groups having six animals in each group by using Complete Randomized Design (CRD). They were drenched with Fenbendazole @ 5 mg/kg body weight against internal parasites before assigning in experiment.

b) Concentrate Mixture Composition

Feed ingredients maize, soybean cake, rice bran, minerals and salt were procured from Khowpa Feed Industry, Bhaktapur. For T1 and T2 concentrate mixture were composed by using procured feed ingredients with 16% crude protein level that has been presented in Table 1 while for T3 commercial compound feed was used made by Pancharatna Feed Industry, Narayangadh, Chitwan.

Table 1 : Composition of concentrate mixture

S/n	Ingredients	T 1		T2	
		Part	Crude Protein (%)	Part	Crude Protein (%)
1	Maize	50	3.94	50	3.94
2	Soybean cake	28	10.7	28	10.7
3	Rice bran	20	1.77	20	1.77
4	Mineral mixture	1	0	1	0
5	Salt	1	0	1	0
Total		100	16.41	100	16.41

c) Heat treatment of soybean cake

The drying of forage is known to increase the protection of the proteins. Thorough heating of protein supplement causes denaturation of protein; it provides effective protection against microbial fermentation in the rumen. Heat treatment was done by using hot air oven at temperature 125- 150^o C for 2-4 hours as suggested by Suresh, *et al* (2009).

d) Experimental diet of the animal

The dry matter requirement of goats was calculated based on 5 kg per 100 kg body weight. Following diets were formulated to the experimental animals (Table 2).

Table 2 : Experimental diets of the animals

Treatment	Experimental diet
1	Forest mixed fodder (adlib) + treated soybean cake included concentrate mixture @ 1.5% of body weight
2	Forest mixed fodder (adlib) + untreated soybean cake included concentrate mixture @ 1.5% of body weight
3	Forest mixed fodder (adlib) + commercial concentrate mixture @ 1.5% of body weight

e) *Feeding Regime*

Concentrate mixture and *adlib* amount of fodder was provided to the experimental animals individually in plastic vessel. Concentrate mixture was provided once a day in the morning whereas fodder twice a day (morning and evening). Quantity of concentrate mixture and fodder given daily to the animals was weighed daily and refusal was weighed in next morning. Experimental animal had free access to drinking water.

f) *Chemical Analysis*

The samples of feed ingredients, prepared concentrate mixture and forest mixed fodder were sent to the Animal Nutrition Division, Khumaltar, Lalitpur for proximate analysis. Representative samples were analyzed for dry matter (DM), crude protein (CP), crude fibre (CF), ether extract (EE) and total ash contents (TA). The DM was determined by oven drying at 100°C for 24 hrs. Crude protein of the samples was determined using the Kjeldahl method. Ether extract was determined using Soxhlet apparatus. Ash content was determined by ashing at 550°C in a muffle furnace for 16 hrs (AOAC 1980). Crude fibre of the samples was determined using

the Van Soest method (Goering, H.K. and Van Soest 1970).

g) *Results Recording*

The trial period consisted 90 days after an adaptation period of 7 days. Total feed intake by the goats was recorded daily for all experimental period. The body weight gain of individual animals was measured fortnightly in the morning before feeding.

h) *Data Analysis*

Data of feed intake and body weight gain were analyzed by “*One Way Anova*” test for every measurement using computer statistical package Minitab 2003, versions 13.20.

III. RESULTS AND DISCUSSION

a) *Chemical composition of feedstuffs*

The result of chemical analysis has been given in Table 3 and crude protein content of prepared concentrate mixture was verified in laboratory that is presented in Table 4.

Table 3 : Chemical composition of different feed ingredients (percentage DM basis)

Ingredient	DM	OM	TA	CP	CF	EE
Maize	87.69	97.97	2.03	8.92	2.34	4.48
Rice bran	87.85	89.5	10.5	11.52	4.83	5.1
Soybean cake	86.87	92.63	7.37	44.29	9.38	0.7
Mixed forest fodder	39.94	90.01	9.99	11.16	NA	NA

The calculated value of crude protein was verified with laboratory analysis, which is presented in Table 4.

Table 4 : Chemical composition of prepared concentrate mixture (% DM basis)

Particular	DM	OM	TA	CP	CF
Treated soybean cake included concentrate mixture	91.92	92.54	7.46	16.83	5.95
Untreated soybean cake included concentrate mixture	91.58	92.10	7.90	16.92	5.64
Commercial feed	90.74	89.85	10.15	15.94	6.45

b) *Feed Intake*

Average daily intake of concentrate mixture and fodder by goats during experimental periods is given in Table 5. Higher intake of concentrate feed was recorded for T1 (207.6 g) followed by T3 (199.58 g) and T2 (193.87 g) which was highly significant (P<0.001) among diet groups. Similarly, fodder intake was also noted significantly higher (P<0.001) among diet groups (1969.5 g , 1967.6 g and 1942 g for T2, T3 and T1,

respectively), however, dry matter intake was observed almost similar for all treatment group (89.53 kg, 89.33 kg and 89.28 kg for T3, T1 and T2, respectively). Feed and fodder intake of different genotypes of goats was found to be non-significant among goat breeds. In addition, feed conversion ratio per kg body weight gain was observed higher for T3 (22.49:1) followed by T2 (17.57:1) and T1 (16.24:1).

Table 5 : Feed intake of experimental animals/day/animal

Feedstuffs	Mean ± SD		
	T1	T2	T3
Feed intake (g)	207.6±57.96	193.87±45.70	199.58±35.15
Fodder intake (g)	1942.3±439.1	1969.5±440.4	1967.6±443.1
Total dry matter intake (DMI) (kg)	89.33	89.28	89.53
Feed conversion ratio (FCR)	16.24:1	17.57:1	22.49:1

c) *Growth Performance*

Growth is a complex; highly integrated process involving numerous interactions among nutrients, environment, genotype, and many different hormones and receptors of these hormones in various tissues (Spencer 1985). The growth performance of experimental goats is given in Table 6 and Figure 1. Initial body weight of T1, T2 and T3 was 12.15 kg, 11.25 kg and 12.18 kg, respectively that reached 17.66 kg, 16.33 kg and 16.40 kg during 90 days of experiment for T1, T2

and T3, respectively. Both initial and final body weight was non-significant among diet groups. Similarly, there was also non-significant effect of goat breed on body weight gain. Total body weight gain was recorded higher for T1 (5.50 kg) followed by T2 (5.08 kg) and T3 (3.98 kg) which was significant ($P < 0.05$) among diet groups. Similarly, average daily gain was also noted higher in T1 (61.2g) followed by T2 (56.48 g) and T3 (44.22 g) which also significant ($P < 0.05$) among diet groups.

Table 6 : Growth performance of goats

Parameter	Mean \pm SD		
	T1	T2	T3
Initial Body weight (kg)	12.15 \pm 4.43	11.25 \pm 3.11	12.18 \pm 2.27
Initial metabolic weight (kg)	6.5	6.14	6.51
Final Body weight (kg)	17.66 \pm 4.48	16.33 \pm 3.03	16.40 \pm 2.56
Final Metabolic weight (kg)	8.61	8.12	8.14
Total weight gain (kg)	5.50 \pm 0.36	5.08 \pm 1.06	3.98 \pm 0.40
Average daily gain (g)	61.20 \pm 4.01	56.48 \pm 11.87	44.22 \pm 4.48




Figure 1 : Body weight gain trend of goats during experiment period

IV. DISCUSSION

This study was initiated with objective to compare the growth performance of female goats of different breeds fed with treated, none treated soybean cake incorporated concentrate mixture, and commercial compound feed on fodder based basal diet. Result revealed that concentrate feed and fodder intake was highly significant ($P < 0.001$) among diet groups. Both initial and final body weight was not significant among groups, however, total weight gain and average daily gain of experimental animals significantly ($P < 0.05$) differed among diet groups. Similarly, result proved that

there was not significant effect of breed in feed intake and body weight gain. This experiment revealed that heat treatment of soybean cake significantly improves the feed intake and body weight gain in comparison to without heat treatment.

Limea *et al.* (2011) tested the effects of a treated soybean cake incorporated concentrate diet on 32 Creole goats for growth, carcass fat, and fatty acid composition of muscle (supraspinatus), perirenal and intramuscular adipose tissues. Goats were fed a tropical green forage *Digitaria decumbens* ad libitum with no concentrate (G0) or 1 of 3 levels of concentrate: 140, 240 and 340 g/day, respectively. Goats were

slaughtered according to the standard procedure at the commercial body weight (22 to 24 kg). Goats fed the concentrate diets had greater average daily gain ($P < 0.001$), cold carcass weights ($P < 0.001$), perirenal ($P < 0.01$) and intramuscular ($P < 0.01$) adipose tissues weights. Feeding increased concentrate did not increase the content of any cholesterol-increasing saturated fatty acid in meat.

Sahlu *et al.* (2012) conducted an experiment on growing female angora goats with four treatment groups. Experimental animals were fed with low protein high degradability, 12% CP with conventional, solvent-extracted soybean meal; Low protein low degradability, 12% CP with expelled, heat-treated soybean meal; high protein high degradability, 19% CP with conventional, solvent-extracted soybean meal; and high protein low degradability, 19% CP with expelled, heat-treated soybean meal. They reported that average initial body weight of the goats was 22.1 kg and the final body weight was 26.2 kg with no differences ($P > 0.05$) in body weight gain between treatments. Mean DMI increased significantly ($P < 0.01$) as dietary CP level increased but was not affected significantly ($P < 0.05$) by heat treatment of the dietary protein. Their observations were in agreement with those of Roffler *et al.* (1978), Barney *et al.* (1981) and Blauwiel and Kincaid (1986). Others have reported a decrease in DMI when dietary CP was increased (Foldager and Huber 1979; Edwards *et al.* 1980; Grieve *et al.* 1980). A recent study indicated that DMI increased linearly in young growing goats as dietary CP level increased (Lu and Potchoiba 1990)

V. CONCLUSION

Our experiment revealed that there is a significant effect of heat-treated soybean cake on feed and fodder intake, and total body weight gain of goats; however, it could not be seen as mentioned in different literatures. Perhaps it might be due to winter season of experiment conduction when most of the protein and energy were spent for body maintenance. Therefore, it is suggested that this type of experiments should be continued in future also considering winter loss of weight gain of animals.

VI. ACKNOWLEDGEMENT

The authors would like to express their most sincere gratitude and appreciation to Dr. Bhoj Raj Joshi, Director, National Animal Science Research Institute, Khumaltar, Lalitpur for his encouragement, help, suggestion and support during experimental period. Similarly, work of Mrs. Bishnu Ale (Enumerator) and Mr. Tek Bahadur Ale (Junior Technician of ARS, Bandipur) deserve high appreciation without whom conduction of experiment was impossible. Mr. Basanta Kumar

Shrestha Technical Officer, Animal Nutrition Division, Khumaltar also deserve appreciation for his dedicated work in chemical analysis of fed ingredients and fodders. Likewise, our thanks also go to the all-technical, admin and finance staff of National Animal Science Research Institute, Khumaltar and Agriculture Research Station (Goat), Bandipur, Tanahu for their help during trial period.

REFERENCES RÉFÉRENCES REFERENCIAS

1. (1980). *Association of Official Analysis Chemists, Washington DC, U.S.A.* Official methods of analysis.
2. Aziz, M.A., (2001). Present status of the world goat populations and their productivity. 45 (2), 42.
3. Broderick, G.A., Wallace, R.J., Ørskov, E.R. and Hansen, L., (1988). Comparison of estimates of ruminal protein degradation by in vitro and in situ methods. *Journal of Animal Science*, 66, 1739-1745.
4. Bajjleih, N., (2002, April). Proceedings of the protein source for the animal feed Industry., 141-160.
5. Barney, D. J., D. G. Grieve, G. K. Macleod and L. G. Young, (1981). Response of cows to a reduction in dietary crude protein from 17 to 15% during early lactation. *Journal of Dairy Science*, 64:25.
6. Blauwiel, R. and R. L. Kincaid, (1986). Effect of crude protein and solubility on performance and blood constituents of dairy cows. *Journal of Dairy Science*, 89, 2091.
7. Edwards, J. S., E. E. Bartley and A. D. Dayton, (1980). Effect of dietary protein concentration on lactating cows. *Journal of Dairy Science*, 243.
8. EL-Shabrawy, H. M., (1996). Utilization of dietary protein in ruminants solubility and rumen degradability of some protein and their protection.
9. EgypFoldager, J. and J. T. Huber., (1979). Influence of protein percent and source on cows in early lactation. *Journal of Dairy Science*, 82, 954.
10. Goering, H.K. and Van Soest, (1970). Forage fibre analysis apparatus, reagents, procedures and some application.
11. Grieve, D. G., E. E. Wheeler, Y. Yu and G. K. Macleod, (1980). Effects of dry or ensiled feeds and protein percent on milk production and nitrogen utilization by lactating cows. *Journal of Dairy Science*, 63, 1282.
12. Limea, L. G. Alexandre and V. Berthelot (2011). Fatty acid composition of muscle and adipose

tissues of indigenous Caribbean goats under varying nutritional densities, *Journal of Animal Science*, 90 (2), 605-615.

13. Lu. C. D. and M. J. Potchoiba, (1990). Feed intake and weight gain of growing goats fed diets of various energy and protein levels. *Journal of Animal Science*, 68, 1751.
14. (2012). *Ministry of Agriculture Development, Agri-Business Promotion and Statistic Division*. Statistical information in Nepalese agriculture, Nepal.
15. Neopane, S.R. and P.K. Pokharel, (2008). Indigenous goat of Nepal.
16. Promkot, C. and M. Wanapat, (2003). Ruminal degradation and intestinal digestion of crude protein of tropical protein resources using nylon bag technique and three-step in vitro procedure in dairy cattle. 15, 11.
17. Roffler, R. E., L. D. Satter, A. R. Hardie, and W. J. Tyler, (1997). Influence of dietary protein concentration on milk production by dairy cattle. 12, 31- 46.
18. Suresh, K.P., R. Bhatta, S. Mondal and K.T. Sampath, (2009). Effect of bypass protein on milk yield in Indian cattle – A meta-analysis. 11, 19-26.
19. Sahlu, T., J. M. Fernandez, C., D. Lu and R. Manning, (2012). Dietary Protein Level and Ruminal Degradability for Mohair Production in Angora Goats'. *Journal of American Society of Animal Science*, 70, 1526-1533.
20. Schroeder, J.W., (1997). Corn gluten feed for dairy cattle.



GLOBAL JOURNAL OF SCIENCE FRONTIER RESEARCH
AGRICULTURE AND VETERINARY
Volume 13 Issue 11 Version 1.0 Year 2013
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Analysis of Rural Cassava Farmers' Participation in the Nigeria Agricultural Insurance Scheme in Imo State, Nigeria

By Nnadi, F.N., Nnadi, C. D., Chikaire, J., Umunnakwe, P.C. & Ihenacho, R.A.

Federal University of Technology, Owerri, Nigeria

Abstract - The importance of insurance in mitigating food insecurity necessitated this study that analyzed rural cassava farmers' participation in the Nigeria agricultural insurance scheme in Imo State, Nigeria. The study focused on the socio-economic differentials of participants and non participant; reasons for participation and determinants of participation. Data collected from 90 sampled cassava farmers using structured questionnaire and interview schedule were analyzed with the aid of percentage count, frequency tables, z-test and logit regression model at 0.05 levels of significance. The result shows socio-economic differentials in the age, education, farming experience, social organization membership, status of participants and non participants in the scheme. The reasons cited for participation included to acquire loan and continue in business even after suffering losses. While the reasons for non – participation included inadequate knowledge of the scheme and cost of insurance, the socio-economic and farm enterprise characteristics of age, education, marital status, farming status, farming experience, farm size and credit opportunity determined the farmers that participated in the scheme. It was recommended that extension education campaign be mounted for enlightenment of the scheme and consideration should be given to the farmers socio-economic and farm enterprise characteristics in designing intervention strategies and advocacy on the scheme.

GJSFR-D Classification : FOR Code: 070199



Strictly as per the compliance and regulations of :



© 2013. Nnadi, F.N., Nnadi, C. D., Chikaire, J., Umunnakwe, P.C. & Ihenacho, R.A. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License (<http://creativecommons.org/licenses/by-nc/3.0/>), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Analysis of Rural Cassava Farmers' Participation in the Nigeria Agricultural Insurance Scheme in Imo State, Nigeria

Nnadi, F.N.^α, Nnadi, C. D.^σ, Chikaire, J. ^ρ, Ummnakwe, P.C.^ω & Ihenacho, R.A.[¥]

Abstract - The importance of insurance in mitigating food insecurity necessitated this study that analyzed rural cassava farmers' participation in the Nigeria agricultural insurance scheme in Imo State, Nigeria. The study focused on the socio-economic differentials of participants and non participant; reasons for participation and determinants of participation. Data collected from 90 sampled cassava farmers using structured questionnaire and interview schedule were analyzed with the aid of percentage count, frequency tables, z-test and logit regression model at 0.05 levels of significance. The result shows socio-economic differentials in the age, education, farming experience, social organization membership, status of participants and non participants in the scheme. The reasons cited for participation included to acquire loan and continue in business even after suffering losses. While the reasons for non – participation included inadequate knowledge of the scheme and cost of insurance, the socio-economic and farm enterprise characteristics of age, education, marital status, farming status, farming experience, farm size and credit opportunity determined the farmers that participated in the scheme. It was recommended that extension education campaign be mounted for enlightenment of the scheme and consideration should be given to the farmers socio-economic and farm enterprise characteristics in designing intervention strategies and advocacy on the scheme.

1. INTRODUCTION

In spite of the impressive effort and conceited persuasion to invest in agriculture by the government, the reluctance expressed by credit institutions has been worrisome. This stems from the low confidence in the agricultural sector following unprecedented risks and uncertainties in the practice. Agriculture is bedeviled by price fluctuation, instability in input and production supplies, poor yield and post harvest losses, pests and diseases attack, inclement weather and vagaries of environmental conditions. These have individually and collectively enmeshed the rural farmers in the web of poverty. Ijere (1981) observed that large volume of investible fund is imperative to disentangle the rural

farmers from the vicious cycle of poverty. However, the nostalgia about credit disbursement to farmers is based on the skepticism on their repayment ability. Njoku and Nzenwa (1990) attributed high loan default rate to the occurrence of natural hazards. Insurance is considered as one of the most effective means of reducing the vulnerability of the poor from the impacts of disease, theft, violence, disability, fire and other hazard. Insurance protects against unexpected losses by pooling the resources of the many to compensate for the losses of the few, the more uncertain the event the more insurance becomes the most economical form of protection (Brown and Churchill 1999). Policyholders only pay the average loss suffered by the group rather than the actual costs of an individual event: insurance replaces the uncertain prospect of large losses with the certainty of making small regular affordable premium payments (Brown and Churchill, 1999). The primary function of insurance is to act as a risk transfer mechanism to provide peace of mind and protect against losses. Risk can be handled by: assumption, combination, transfer or loss prevention activities. Insurance schemes utilize the combination method by persuading a large number of individual to pool their risks into a large group to minimize overall risk (Aliero and Mukhtar, 2012). In the developed world insurance is part of society, such that some forms of cover are required by law. In developing countries the need for such a safety net is much greater particular at the poorest level where vulnerability to risk is much greater and there are fewer opportunities available to recover from a large loss (Aliero and Mukhtar, 2012).

In the light of the above, the Government of Federal Republic of Nigeria identified Agricultural insurance as a panacea to the doubt and attendant disenchantment expressed by credit institutions following the multifarious risks and uncertainties in agriculture. In 1987, the government of Nigeria formerly launched the Nigerian Agricultural Insurance Scheme (NAIS) and in 1988 incorporated the Nigeria Agricultural Insurance Company (Nnadi, *et al*, 2013). Agricultural insurance is the stabilization of income employment prices and supplies of agricultural products by means of regular and deliberate savings and accumulation of funds in small installments by many in favorable time periods to defend the participants in bad times (Mordi

Authors α ρ ω : Department of Agricultural Extension, Federal University of Technology, Owerri.

Author σ : Department of Agricultural Economics and Rural Sociology , Niger Delta University Bayelsa State Nigeria.

Author ¥ : Department of Agricultural Management and Extension Technology, Imo State Polytechnic, Umuagwo, Owerri, Imo State, Nigeria. E-mail : aksymeon35@yahoo.com

1995). Farmer's losses, indisposition and fear are allayed following the cushioning effects from the accumulated saving. Thus farmers are put back to business irrespective of misfortune suffered provided the peril is covered.

The Nigerian agricultural insurance scheme was institutionalized in 1987 to obviate problems of knowledge imperfection, risks and Uncertainties in agricultural enterprises. The scheme inter alia offers protection to the farmers from the effects of natural disaster and ensures payment of appropriate compensation, sufficient enough to keep farmers in business after suffering losses. The government on the other hand is provided with back-up and information on agricultural development in the county. The scheme covers arable crops, crops and livestock. The Nigeria agricultural insurance company (NAIC) succinctly put the objectives of the scheme as follows:

- (a) To promote agricultural production by enhancing greater confidence in adopting new and improved farming practices greater confidence in adopting new and improved farming practices and making for greater investment in the agricultural sector thereby increasing the total production.
- (b) To provide financial support to farmers in the event of losses arising from natural disasters.
- (c) To increase the flow of agricultural credit from lending institutions to the farmers.
- (d) To minimize or eliminate the need for emergency assistance provide by government during periods of agricultural disasters.

Agricultural insurance holds wonderful prospects for the transformation of the agricultural sector. By indemnifying farmers from the perils covered in the scheme (Mordi, 1995), they are put in the same closer financial pedestal to operate after suffering losses. Following the numerous advantages of the scheme, it is expected that farmers involved in the production of crops prone to high risk and uncertainties but distinguished for mitigating food insecurity, and have wide spread uses should avail themselves of the scheme. Cassava (*Manihot esculentus*) is marked with such potentials. Cassava provides about 40 percent of calories consumed in Nigeria (Nwajiuba, 1995) and about 70 percent of the daily calorie intake of more than 50million Nigerians (Ugwu et al, 1989). Whereas, Adekanya (1985) posited that cassava is the most important root crop in the tropics. Odigbo (1983) observed that the demand for cassava especially for export has increased appreciably following the Federal Government of Nigeria's cassava initiative. Cassava production is bedeviled by pests and diseases attack, fire outbreak, flooding, poor storage etc (Youdeowei et 1985; Theberge 1985).

It is however unfortunate that despite the status of cassava little or nothing is known about the farmers'

participation in Nigeria agricultural insurance scheme to boost production by forestalling risks and uncertainties. There are on empirical data on this. Information available are based on guesses and suppositions. These have given rise to a wide gap in knowledge the ensuing gap has been inhibiting the formulation of holistic policy measures. A study of cassava farmer's participation in the scheme has become as timely as it is important, not only to position the farmers strategically but to equip them with the necessary thrust essential for meeting the food security needs of the nation ensuring persistence in farming and overcoming the challenges of the present especially in the light of the global climate change.

a) *Objective of the study*

The broad objective of the study is to analyze the participation of cassava farmers in the Nigeria Agricultural Insurance Scheme with a view to making policy recommendations. The specific objectives include;

1. to analyze the socio-economic differentials of participants and non participants in the scheme.
2. to investigate reasons for the participation and non participation of the farmers in the scheme.
3. to analyze the socio-economic and farm enterprise factors that determines participation in the scheme.

II. METHODOLOGY

The study was carried out in Imo state, Nigeria with specific focus on Ohaji-Egbema Local Government area (LGA). The local Government area is one of the 27 local government areas that make up the state. Located in owerri agricultural zone of the state, the headquarters is in Egbema. It shares boundaries with Oguta L.G.A in the East, Owerri west in the West and North and Rivers State in the South. Ten communities make up the Local Government Area: Egbema, Awara Umuapu, Umuagwo Mgbirichi/ Abakuru, Ohuba Assah/ Obitti, Umuokanne, Mmahu and Abuchara. The population by 2013 census is 209,593 projected from 2009 official (FGN 2009) the people are Igbos. The area is located in the rainforest region two distinct seasons abound- rainy and dry. The mean rain fall is 200-25cm (FDLAR1985), with temperature of 26-28c and relative humidity of 80-90 percent (Ugwu and Lekwa, 1988). Agriculture features prominently in the economy. This is rain-fed. Soil fertility maintenance is mostly by natural means, bush fallow system. Crop production encompasses the cultivation of cassava, yam, maize, pineapple, banana, plantain, oil palm and various forms of vegetables. The animals reared include goat, sheep, pig, poultry and most recently grass-cutter. There are also pockets of farmers engaged in aquaculture and apiculture. Data were collected from primary and secondary sources. These

included the use of semi-structured questionnaire supplemented by interview schedule and records from Nigeria agricultural insurance company (NAIC) and agricultural development programme (ADP). The semi-structured questionnaire was validated by experts in agricultural extension and rural sociology. These were tested for reliability using test-re-test method on a group of cassava farmers in Oguta local government area of Imo state to yield a coefficient of 0.68, significant at 5 percent level. The questionnaire and interview schedule were administered between February and June, 2008 by the researchers with the assistance of the extension agents working in the ADP circles covered in the study. The cassava farmers in the L.G.A. composed the study population. They were purposively dichotomized into participants and non participants in the Nigeria agricultural insurance scheme. From the NAIC list of cassava farmers who participated in the scheme in 2007, a total of 45 participants was randomly sampled without recourse to their communities as they were few (<50). Also from the ADP list of cassava farmers, 45 non participants in the scheme were also randomly selected. Thus, a total of 90 cassava farmers; participants and non participants in the scheme made up the sample size. Percentage count and frequency tables were used to describe objectives 1 and 2. while z-test and Logit regression technique were used to analyze objectives 3 and 4 respectively. The z-test statistic expressed as:

$$z = \frac{X_1 - X_2}{\sqrt{\frac{S_1^2}{n_1} + \frac{S_2^2}{n_2}}}$$

where ;

X_1 -mean values of the socio-economic variables of participants in NAIS.

X_2 - mean values of the socio-economic variables of non participants in the scheme.

S_1^2 -variance of participants in the scheme

S_2^2 - variance of non participants in the scheme

n_1 - number of participants

n_2 - number of non participants

X_1 - X_6 - variables whose differentials were determined

X_1 - age (years)

X_2 -education (years of formal education)

X_3 - household size (number of people that feed from the same pot)

X_4 - farm size (hectares)

X_5 - farming experience (number of years of farming)

X_6 - social organization membership status none member =0, ordinary member= 1,regular attendant to meeting =2, financial member=3, committee member =4, executive member=5)

The logit regression technique was expressed as follows (pindyck and Rabinfeld, 1981):

$$P_i = c / (1 + e^{-z_i})$$

Where

P_i - probability that an individual farmer i ($i=1,2,\dots,n$) will make a particular choice)

c -constant

z_i -choice index

$$z_i = \beta_0 + \beta_1 x_{1j} + \beta_2 x_{2j} + \dots + \beta_k x_{kj}$$

where ;

$x_{ij}, j=2,\dots,k$ are the factors influencing the farmers' decision to participate or not in NAIS.

III. RESULTS AND DISCUSSION

a) Socio-economic differentials and non participants in Nigeria agricultural insurance scheme

The z-test results of the socio-economic differentials of participants and non- participants in the Nigerian agricultural insurance scheme (table1) shows that 66.67 percent of the explanatory variables investigated differed significantly between the two groups. Specifically, the variables age education farming experience and social organization membership status differed significantly between the participants and non participants. The mean age of the participants was 46.56 years while that of the non-participants was 50.24. The variance for participants was 5.39 while that of non-participants was 8.03. The z-value was -2438. This implies that the higher the age the lower the participant. Thus the participants were younger than the non-participants, and were more disposed to participating in Nigerian agricultural insurance scheme. This could be explained by their higher venturesomeness, innovativeness and more risk proneness, and insurance is a pool of risk. The result corroborates Nnadi and Akwiwu (2006 a) in which young women farmers utilized proved soil management practices more than the old.

The mean number of years of formal education by participants and non participants were 10.85 and respectively. Whereas the variance for the participants was 4.07, that of non-participants was 1.46. The z-value was 6.226. The positively significant difference implies that higher years of formal education predisposed participation in the scheme. This could be explained by better understanding of the scheme, adequate knowledge of the gain and potentialities for greater investment. of course highly educated farmers could access diverse information sources on the scheme, and better clarification for participation. These affirm the excellence of educated farmers in improved technologies adoption (Nnadi and Akwiwu, 2005a; Onu, 2005; Polson and Spencer, 1981)

Farming experience by participants in the scheme had a mean value of 17.41 while that of the non-participants was 21.03 years. With variance of 6.79 and 9.03 for participants and non participants respectively, the z-value was -2.033. The significant but inverse difference shows that non participants had more of experience but this was not an asset for participation in the scheme. Long years of farming experience presuppose increased chronological age of the farmers. The older the more risk averse and the more conservative the farmers become.

The mean value for social organization membership status; of participants in the scheme was 3.72. The variance was 1.28. The non participants had a mean of 0.39 and variance of 0.11. The z-value was 16.165. It implies that the participants had higher commitments to their social organizations and this positively impacted on their embrace of the scheme. Social organization membership besides meeting up with the farmers' social needs exposed farmers to settings where their misconceptions and distortions are clarified. The result is in consonance with Mgbada (2002) that the more active the farmers are in their social organizations, the more they are exposed to useful information about innovation and the farm size were not significantly different. These are not important variables for consideration in targeting cassava farmers for participation in the scheme.

b) Reasons for participating in Nigeria agricultural insurance scheme

Diverse reasons were given for participating in the Nigeria agricultural insurance scheme (Table 2). The whole farmer (100%) indicated that their participation was to enable them have access to loan. Agricultural credits were described as an imperative for rural transformation (Ijere, 1981). As a condition for accessing agricultural credit from Nigeria Agriculture Credit and Rural Development Bank (NACRDB), farmers must undertake insurance cover. By compulsion most prospective loan beneficiaries indemnify their farms to increase the confidence of the lending institution. On questioning the farmers for their reasons for settling for the rigors for obtaining loan from Nigeria Agriculture, Co-Operatives and Rural Development Bank (NACRDB), three-quarters of the farmers indicated that the bank render better and friendly services.

Another reason for participating in the scheme by the farmers was to continue to be in business after suffering losses (79%). The third reason on the rank was to protect the farmers from the effect of natural disaster (69.2%). The various underscore adequate understanding of the objectives of the scheme and thus laid credence to the perception of insurance as a social device to provide financial compensation for the effect of misfortune. The fourth reason for participate in the scheme was to expand investment in agriculture

(59.0%). The farmers were further interview on why they though insurance could help them expand their agricultural investment. Two thirds of the farmers noted that unindemnified losses could either push farmers to the basic or our business but indemnity portends rays for continuity and opens farmers' eyes to vistas of opportunities hitherto unexplored. To get along with other farmers rank 5th with 25.6 percent. The response does not reflect adequate understanding of the importance of the scheme. This calls concerted extension awareness campaign.

c) Reasons for non participation in the Nigeria agricultural insurance scheme

The reasons for not participating in the Nigeria agricultural insurance scheme ranged from logistics in the scheme (70.7%), 1st in the rank. The inadequate knowledge could be attributing to poor extension campaign or poor geographical spread of NAIS offices for easy access by the farmers. Logistics in the scheme could be explained by the bureaucracy in registration, subsequently verification and processing of document at the event of loss. The cost of premium ranked 2nd with 90.2 percent. The third in the rank, lack of confidence in institution had 85.4 percent. This could be attributed to unsavoury past experience. Fear of the unknown ranked 4th with 78.1 percent. This could be adduced to poor understanding of the scheme and the methods operation. Generally, the reasons for not participating in the scheme are based on ignorance and hence unfounded. Participation could be improved through education campaign.

d) Socio-economic and farm enterprise determinants of cassava farmers' participation in NAIS

The logic regression result of the socio economic and farm enterprise determinant of cassava farmers participation in NAIS (table 4) shows that seven independent explanatory variables (70%) were significant at 0.05 level. The variable included age (X_1) education (X_3) marital status (X_4) farming status (X_6) farming experience (X_7) farm size (X_9) and credit opportunity (X_{10}).

The age of the farmers (X_1) had a coefficient of 0.1847 and T value of 3.3515. The result implies that increasing the magnitude of the farmer's age increased their participation in NAIS. Specifically, each additional year to the age farmers increased the probability of their participation by about 19 percent. Age is therefore a major consideration in designing strategies to increase participation in the scheme. This could be attributed to increased maturity and experience as the farmers are bound to make better informed decisions following increased life encounters over time. More so, increased age is associated with more responsibilities, marriage, caring for children and expanded scope of dependants and insurance could become a source of respite by ensuring continuity in farm engagement, even after

suffering losses. In line with this, age was found to be positively significant to rural women's use of improved crop production technologies in Imo State (Nnadi and Akwiwu 2005b).

Education had a coefficient of 0.1192 with a t-value of 3.722. The result shows that increases number of years of normal education impacted positive on the farmers in NAIS. The result implies that each additional year of formal schooling increased the probability of participating in NAIS by about 12 percent. Education furnished facts exposed farmers to multifarious information sources, polished their reasoning and decision making processes. The result concurred with Agada and Philip (2002) in which education maize farmers participated more in NAIS in Kaduna State, Nigeria.

The status of the farmers (full time or part time) had a coefficient of 0.1604 and t-value of 2.6081. The positive but significant relationship implies that an additional improvement in farming status, by part time becoming full time farmers, increased the probability of participating in NAIS by 16 percent. Full time farming status could entail that the farm is the monolithic source of revenue to the farmer. Insurance therefore becomes a safe guard for subsistence, especially during losses. Farming experience had 0.0882 and 2.1356 as coefficient and t-value respectively. Increasing the number of years of farming experience thus increased the farmers' participation in NAIS. The result implies that each additional year of experience in farming resulted to about 9 percent increase in the probability of their participation in NAIS. Increased years of farming experience just like increased age could entail several varied encounters in farming which could influence farm decision making. The study of Nnadi and Akwiwu (2006b) also established positive significant relationship between years of farming experience of women and the number of coping strategies against economic marginalization.

The coefficient of farm size was 0.0436 while the t-value was 4.1983. The result implies that each additional hectare of land put into cassava cultivation resulted to 4 percent increase in the probability of participating in NAIS. Large farm size could entail a large farm asset base. This could also mean high level of investment; input, credit technologies, etc insurance therefore becomes an imperative option against unprecedented losses. The result of the studies of Nnadi and Akwiwu (2006c) and Nnadi and Akwiwu (2005a) affirmed the importance of farm size in farm decision making.

The coefficient of credit opportunity (0.0843) was positively significant with a t-value of 2.0611. Increased the farmers' participation in NAIS. Specifically, the result implies that additional increase in the number farmers that had access to credit offers opportunity to increase a farmer's capital base and subsequently has

investment. The finding agrees with Nnadi and Akwiwu (2006a) in which credit opportunity positively influenced rural women's adoption of proven soil management practices.

However, the variables gender, household size and social organization membership status were not significant related to the farmer's participation in NAIS. They therefore do not determine their participation and as such should be discountenanced in advocacy and designing intervention strategies.

IV. CONCLUSION

Participants in the Nigeria agricultural insurance scheme differed significantly from non participant in the scheme, in respect of age education, farming experience and social organization membership status. The reasons for participating included being able to acquire loan, to continue in business after suffering losses and expanding investment in agriculture. The reason for non participation included inadequate knowledge about NAIS, cost of insurance and lack of confidence in the institution. The farmers participation in the scheme were determined by their socio-economic and farm enterprise characteristics of age education, marital status, farming experience, farm size and credit opportunity.

Policy implication

Extension education campaign should be embarked upon enlighten non participants in the scheme on the prospect as well as sustain the interest of the participants.

- The socio-economic and farm enterprise characteristics of the farmers should be put into consideration in designing intervention strategies and advocacy for increased participation in NAIS
- Institutional reforms, land and credit should be vigorously pursued to avail more farmers of them for increased participation in the scheme.

REFERENCES RÉFÉRENCES REFERENCIAS

1. Adekanya, T.O. (1985). Some considerations for Development Technology in Agriculture Paper Presented at the 2nd national conference of farm management association Zaria. 5-7.
2. Aliero, H.M. and Muktar, S. (2012). The Prospects of Micro-Insurance in the Rural Areas of Nigeria. *European Scientific Journal*, 8 (3), 66-76.
3. Agada, J.E and Philip D (2002). "A logit analysis of the participation in the Nigerian Agriculture Insurance Scheme by maize growing farmers in Kaduna state". *Asset Journal*, 2 (1), 157-163.
4. Brown, W. and Churchill, C. (1999). Micro-Insurance: Providing Insurance to Low Income Households.
5. (1985). The reconnaissance soil survey of Imo State Nigeria.

6. Mgbada, J.U (2002). Elements of agricultural Extension Richfield and Frank law and science.
7. Mordi, O. (1995). "Elements of Insurance".
8. (1989). Nigeria Agricultural Insurance Company Operational Guidelines.
9. Njoku, J.E and Nzenwa, D (1990). "Loan repayment and its determinants under small holder direct Agricultural loan scheme in Imo State Nigerian". 28 (3), 247-254.
10. Nnadi F.N. and Akwiwu, C.D. (2006a). Adoption of Proven Soil Management Practices by Rural Woman in Imo State,. *Nigeria international journal of national and applied science*, 2 (3), 262-267.
11. Nnadi, F.N and Akwiwu, C.D (2006b). Strategies for Overcoming Economic Marginalization Among Rural women in Imo state Nigeria, *International Journal of Natural and Applied Science*, 1 (1), 10-16.
12. Nnadi F.N. and Akwiwu, C.D. (2005a). Utilization of improved cassava production technologies by urban farmers in Imo. *Nigeria international journal of natural and applied science*, 1 (1), 10-16.
13. Nnadi, F.N. and Akwiwu, Nigeria u, C.D (2005b). Rural women's response to selected crop production technologies in Imo state, Nigeria. 1 (1), 47-54.
14. Nnadi, F.N. and Akwiwu, C.D. (2005c). Adoption poultry production practices by rural women in Imo state. 1 (1), 39-44.
15. Nwajiuba, C.U. (1995). Socio- economic impact of cassava post- harvest technologies in small holders in south eastern Nigeria. 20, 10-14.
16. Nigeria national population Abuja commission. (2009). 1.
17. Nnadi, F.N., Chikaire, J., Echetama, J.A., Iheanacho, R.A., Umunnakwe, P.C. and Utazi, C.O. (2013). Agricultural Insurance: A Strategic tool for Climate Change Adaptation. *Agricultural Insurance: A Strategic tool for Climate Change Adaptation.*, 1 (1), 1-9.
18. Odigboh, E.U. (1985). Mechanization of cassava production and processing a decade of design and development. *university of Nigeria*.
19. Onu, D.O (2005). Analysis of factors influencing farmers adoption of alley farming technology under intensified agriculture in Imo State Nigeria using a qualitative choice model. 29, 177-187.
20. Pindyck, R and Rubinfeld, D. (1981). Economic models and economic forecasts.
21. Polson, R and spencer, D.S.C (1991). The technology adoption process in subsistence agriculture: the case of cassava in southwest Nigeria, 36, 65-77.
22. Theberge, R.L (1985). Common African pests and diseases of cassava, yam, sweet potato and cocoyam, Ibadan.
23. Ugwu, T.O. and lekwa, G (1988), October). Review of the land and soil resources of Imo state in relation to food production and forecasting proceedings of imo state agro statistics sub- committee seminar on food, forecasting and earth warming, 20-66.
24. Ugwu, B.O. Ajabo, O and Orkwor, G (1989). Cassava research production and utilization in Nigeria in Nweke, 15- 24.
25. Youdeowi, A Ezedinma, F.O.C. and Onazi, O.C (1986). Introduction to tropical agriculture.

Table 1 : z test result of the socio-economic differential between participants and non participants in NAIC

Explanatory variables	Means of participant	Means of non participant	Variance of participants	Variance of non participants	Z value
Age	46.53	50.24	5.39	8.03	-2.438*
Education	10.85	7.88	5.09	1.46	2.226*
Farmers size	8.41	9.59	3.05	4.19	-1.455
Farm size	4.08	3.59	1.68	1.24	1.471
Farming exp.	17.41	21.03	6.79	9.03	-2.033*
Social org. mem.	3.72	0.39	1.28	0.11	16-165*

Status

*significant z value at 0.05 levels sources survey data, 2008

Table 2 : reasons for participating in the Nigeria agricultural insurance scheme

Reasons	F	%	Rank
To continue after suffering losses	31	79.5	2 nd
To be able to acquire loan	39	100.0	1 st
To expand investment in agriculture	23	59.0	4 th
To be protected from the effect of natural disaster	27	69.2	3 ^d
To get along with other farmer	10	25.6	5 th

*multiple responses N=39

Table 3 : Reasons for non participation in NAIS

Reasons	*F	%	Rank
Inadequate knowledge of NAIS	38	92.7	1 st
Lack of confidence in the institution	35	85.4	3 rd
Logistics in the scheme	29	70.7	5 th
Cost of insurance	37	90.2	2 nd
Fear of the unknown	32	78.1	4 th

*Multiple responses N = 41

Source -fields survey data, 2008

Table 4 : logic regression result of the socio-economic and farm enterprise determinant of farmers' participation in NAIS

Explanatory variables	Logistic coefficient	t-value
Constant	-23.4016	-5.7944
Model chi square	57.0844	
Degree of freedom	49	
Number of cases	80	
Gender	0.0943	1.1542
Age (X ₁)	0.1847	3.5315*
Education (X ₂)	0.1192	3.7722*
Marital status (X ₃)	0.0349	3.2315*
Household size (X ₄)	0.0839	1.1541
Farming status (X ₅)	0.1608	2.6081*
Farming experience (X ₇)	0.0882	2.1356*
Social organization		
Membership (X ₈)	0.1903	1.0485
Farm size(X ₉)	0.0436	4.1923*
Credit opportunity (X ₁₀)	0.0843	2.0611*

*significant t-value at 0.05 level

This page is intentionally left blank



GLOBAL JOURNAL OF SCIENCE FRONTIER RESEARCH
AGRICULTURE AND VETERINARY
Volume 13 Issue 11 Version 1.0 Year 2013
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Livelihood Vulnerability to Climate Change based on Agro Ecological Regions of Nepal

By Chandra K Dhakal, P. P. Regmi, I. P. Dhakal, Binod Khanal & U. K. Bhatta

Tribhuvan University, Nepal

Abstract - This study based to assess the livestock holders' vulnerability to climate change across four agro ecological regions of Nepal. Altogether 240 households, 60 from each agro ecological region, were selected using stratified random sampling technique. Primary data were collected through household survey and Focus Group Discussion using structured and pre tested questionnaire and analyzed by descriptive statistics. Integrated vulnerability approach based on the Principle Components Analysis to create vulnerability indices to conduct a comparative analysis of vulnerability at the regional levels. The results reveals that the farmers were aware of climate change. The result shows that livestock holders from the fragile warm temperate mountainous region are more vulnerable because of greater exposure to climatic induced hazards, catastrophe, and low adaptive capacity results from highly pauper economic condition, with limited access to basic services, assets, and poor infrastructure. Based on the results, measures to prioritize and target the vulnerable livestock holders for awareness creating activities and of off farm employment opportunities is recommended to enhance their adaptive capacity.

Keywords : livelihood vulnerability, climate change, agro ecological regions, Nepal.

GJSFR-D Classification : FOR Code: 070301, 079999



Strictly as per the compliance and regulations of :



© 2013. Chandra K Dhakal, P. P. Regmi, I. P. Dhakal, Binod Khanal & U. K. Bhatta. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License (<http://creativecommons.org/licenses/by-nc/3.0/>), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Livelihood Vulnerability to Climate Change based on Agro Ecological Regions of Nepal

Chandra K Dhakal^α, P. P. Regmi^σ, I. P. Dhakal^ρ, Binod Khanal^ω & U. K. Bhatta^ξ

Abstract - This study based to assess the livestock holders' vulnerability to climate change across four agro ecological regions of Nepal. Altogether 240 households, 60 from each agro ecological region, were selected using stratified random sampling technique. Primary data were collected through household survey and Focus Group Discussion using structured and pre tested questionnaire and analyzed by descriptive statistics. Integrated vulnerability approach based on the Principle Components Analysis to create vulnerability indices to conduct a comparative analysis of vulnerability at the regional levels. The results reveals that the farmers were aware of climate change. The result shows that livestock holders from the fragile warm temperate mountainous region are more vulnerable because of greater exposure to climatic induced hazards, catastrophe, and low adaptive capacity results from highly pauper economic condition, with limited access to basic services, assets, and poor infrastructure. Based on the results, measures to prioritize and target the vulnerable livestock holders for awareness creating activities and of off farm employment opportunities is recommended to enhance their adaptive capacity.

Keywords : livelihood vulnerability, climate change, agro ecological regions, Nepal.

I. INTRODUCTION

Nepal, with a population of 26.6 million (CBS, 2011) the world's fourth most vulnerable country to climate change due to the fragile ecosystem, which is very sensitive to even slight changes in natural climate, weaker geological situation and complex topography (Maplecroft, 2011). Livestock is an integral part of the mixed farming system and socio-economical life in the country, and contributes nearly 26 percent to the total Agricultural Gross Domestic Product (MoAD, 2012). Around 87 percent of the country's total population keeps some form of livestock at home (IRIN, 2013). Although, Nepal has one of the highest ratios of livestock and poultry to humans (5.8 animals per household) in Asia, the country's livestock sector is declining (IRIN, 2013). While not definite, it would seem that livestock in Nepal is at par with livestock systems in other developing countries and is changing rapidly in response to many external and internal drivers including

Authors α σ ξ : Institute of Agricultural and Animal Science, Tribhuvan University, Rampur, Chitwan, Nepal. E-mail : dhakalck@yahoo.com
Author ρ : Agriculture and Forestry University, Chitwan, Nepal.
Author ω : Institute of Agriculture and Animal Science/ Tribhuvan University, Nepal.

climate change which is seen as a negative impact (Thornton et al., 2007).

The Intergovernmental Panel on Climate Change (IPCC, 2007) suggests that within the agricultural sector livestock are among the most climate sensitive economic areas. Studies on livestock and climate change revealed that climate change adversely affects the animal health and livestock production. Cool temperate Grassland is projected to shift northward with climate change and net primary productivity will decline (Christensen et al., 2004). The limited herbaceous production, heat stress from higher temperature, and limited water intake due to the decrease in rainfall could cause poor livestock performance and an increased incidence of animal diseases. These effects will be felt mostly by the smallholder and subsistence livestock holders' of developing countries are the most vulnerable to livelihood from the effects of climate change (Stern, 2006; Heltberg, 2009) because of lack of resources, knowledge, veterinarian extension services and research technology development (FAO, 2008). This indicates that the livestock sector is the most vulnerable to any adverse impacts of climate variability and extreme events that might result from climate change.

According to the IPCC (2007), vulnerability is defined as the degree to which a system is susceptible to, or unable to cope with, adverse effects of climate change, including climate variability and extremes. Vulnerability is a function of the character, magnitude, and rate of climate variation to which a system is exposed, its sensitivity, and its adaptive capacity. According to Raut (2006), the most vulnerable ecological and socio-economic systems are those with the greatest sensitivity to climate change and the least ability to adapt. Nepal's major natural resources, biodiversity and water, are at the forefront of climate vulnerability. Moreover, Nepal possesses less coping and adaptive capacity to address the additional impacts of climate change.

There is increasing important to understand the likely impacts of climate change on livestock production and their vulnerability. The degree of vulnerability to climate change among the different groups of people is different based upon agro ecological regions. It was necessary to identify likely hotspots that were already vulnerable and that are likely to suffer substantial

impacts as a result of climate change. In this backdrop, this study aims on the assessment of livestock vulnerability to climate change based on agro ecosystem elevation, along with other relevant biophysical and social factors and possible adaptation measures.

II. RESEARCH METHODOLOGY

a) Study sites, sampling, data collection

The Gandaki River Basin (GRB), Nepal spreads from 27.21'45" to 28.036'36" degree north longitude to 83.008'00"- 84.053'00" degree east latitude and elevation ranging from about 144 Masl to 8167 Masl (DDC, 2002). It covers the areas in the Mountain zone (Mustang, Manang, Gorakha, Rasuwa Districts), Hill zone (Myagdi, Kaski, Tanahun, Lamjung, Syangja, Parbat, Dhading, Nuwakot, Makawanpur, Baglung, Gulmi, Palpa), and the valley Terai zone (Nawalparasi, Chitwan, Kapilvastu). The average temperature of this area ranges from -9 °C in Mustang to 42.5°C in Chitwan (DADO, 2012; DLSO, 2011b). Average annual rainfall is 26.58 mms in mustang to 2500 mm in Chitwan (DADO, 2012; DLSO, 2011b). Four agro-ecological regions were selected from Chitwan, Myagdi and Mustang districts of GRB in Nepal. From each region 60 households were selected using purposive simple random sampling technique accruing the total households to be surveyed were 240 households. The primary data was collected through household survey using pretested semi structured questionnaire via face to face interview during October to December 2012.



Fig.1 : Shaded regions showing study area in Gandaki River Basin, Nepal. This research was based on four agro ecological regions namely the tropical region (below 500 meters above sea level) from Chitwan District, subtropical (500 -1000 Masl) and warm temperate regions (1000-2000 Masl) from Myagdi District and cool temperate region (2000-3000 Masl) from Mustang District

As far as Participation goes, two Focus Group Discussions (FGDs) and one Key Informant Interview (KII) were conducted to triangulate the data and to supplement the household survey. Information on the livestock holder's perception on climate change, and major effects on livestock due to changing climatic conditions were assessed through these participatory methods. The Geographical Positioning System (GPS) was used to determine the altitude and latitude of the study areas.

b) Integrated vulnerability assessment

Livestock based livelihood vulnerability to climate change in the study areas was accessed by an integrated vulnerability assessment approach. The integrated vulnerability approach is superior over other approaches and is particularly useful for policy decision making (Deressa et al., 2008). Integrated vulnerability approach comprises socioeconomic and biophysical indicators of vulnerability and classified these indicators into adaptive capacity, sensitivity, and exposure. The data on adaptive capacity and sensitivity were obtained from primary and secondary sources of data while dichotomous variables namely river flooding, landslide and drought on climate extremes captured exposure. Principal Component Analysis (PCA) was performed to compute the component score to weigh the variables to calculate the vulnerability indices. The purpose of using weights obtained from the PCA is to avoid the uncertainty of unequal weighting, given the diversity of indicators used (Deressa et al., 2008). Vulnerability was calculated as defined by IPCC (2001).

Vulnerability = Adaptive capacity - Sensitivity - Exposure.....1.1

Equation 1.1 can be expressed as follows:

$$V_k = \sum_{i=1}^n W_{ki} \cdot X_{ki} - \left(\sum_{i=1}^n W_{ki} Y_{ki} + \sum_{i=1}^n W_{ki} Z_{ki} \right) \dots 1.2$$

Where

$i = 1, 2, 3, \dots n$ Households

$k = 1, 2, 3, \text{ and } 4$, representing ecological regions

V_k = Vulnerability index for kth region

W_{ki} = Weight obtained from first principal component scores of i^{th} variable for k^{th} region

X_{ki} = Adaptive i^{th} for k^{th} region

Y_{ki} = Sensitivity i^{th} for k^{th} region

Z_{ki} = Exposure i^{th} for k^{th} region

While calculating the direction of relationship in vulnerability indicators (i.e., their sign), negative value was assigned to both exposure and sensitivity. The justification is that areas that are highly exposed to damaging climate are more sensitive to damages, assuming constant adaptive capacity (Deressa et al.,

2008). The implication is that a higher net value indicates lesser vulnerability and vice versa (Madu, 2012).

III. RESULTS AND DISCUSSION

a) Livestock Holding

Livestock is an important asset on which livelihood of majority people hinged on. Livestock

holding depicts the picture about farmers' economic condition and it also gives the idea about the total farm yard manure availability in the households. Selected household were observed to raise various kind of livestock species in different number especially Chauri, Chyangra, cattle, buffalo, pig, goat, sheep and poultry.

Table 1 : Livestock holding of respondents across the regions


	Agro Ecological Regions					χ^2
	Tropical	Subtropical	Warm Temperate	Cool Temperate	Total Mean	
LSU	54.07	22.25	8.12	18.82	25.81	
S. E.(±)	19.76	7.74	0.94	5.16	5.54	

b) Perception to climate change

Change in the weather parameters is important factor that determine the farmer's perception about climate change. Nearly all of the respondents (92%) had observed the deviation of weather parameters like variation in rainfall, temperature, humidity and snowfall (Figure 2).

The average livestock holding was 25.81 Livestock Standard Unit (LSU)¹. Livestock holding was greater in tropical (54.07 LSU) among the agro ecological zones under study while least livestock holding was found in warm temperate (8.12 LSU) (Table 1). The highest LSU in tropical may be due to large number of poultry in cluster production and commercial cow production and Chitwan is emerging as the commercial pocket area for livestock sector and contributed more in the household annual income. There was significant differences ($\chi^2 = 7.27$, $P > 0.1$) in the distribution of LSU in households across all agro ecological regions.

Similarly, study revealed that nearly 73.0% of respondents perceived the increase in temperature. According to findings, 10.8% respondent didn't observe any change in the temperature while 3.3 % realized the decrease in temperature (Table 1). The result also showed the decreased and erratic rainfall trend in both summer and winter monsoon in the recent years. More than half of the respondents opined the decreased winter monsoon. Delayed initiation of summer monsoon has noticed as in the case of winter monsoon but quite unpredictable onset of winter monsoon was experienced.



Source: Field Survey, 2012

Fig. 2 : Livestock holders' perception of climate change in the study area

¹ LSU= 1 (Cow/Bull) + 1.5 (Buffalo) + 0.4 (Goat/Sheep) + 0.6 (Swine/Pig) + 0.2 (Poultry)

Table 2 : Perception of respondents towards temperature across the regions

Temperature	Agro Ecological Regions					χ^2
	Tropical	Subtropical	Warm temperate	Cool temperate	Total	
Increased	47.00 (78.33)	43.00 (71.67)	41.00 (68.33)	45.00 (75.00)	176.00 (73.33)	10.46
Decreased	3.00 (5.00)	2.00 (3.33)	1.00 (1.67)	2.00 (3.33)	8.00 (3.33)	
Same	1.00 (1.67)	10.00 (16.67)	9.00 (15.00)	6.00 (10.00)	26.00 (10.83)	
Don't know	9.00 (15.00)	5.00 (8.33)	9.00 (15.00)	7.00 (11.67)	30.00 (12.50)	
Rainfall						
Increased	2.00 (3.33)	5.00 (8.33)	6.00 (10.00)	4.00 (6.67)	17.00 (7.08)	26.17*
Decreased	37.00 (61.67)	33.00 (55.00)	27.00 (45.00)	29.00 (48.33)	126.00 (52.50)	
Erratic	8.00 (13.33)	14.00 (23.33)	6.00 (10.00)	9.00 (15.00)	37.00 (15.42)	
Same	0.00 (0.00)	0.00 (0.00)	9.00 (15.00)	8.00 (13.33)	17.00 (7.08)	
Don't know	13.00 (21.67)	8.00 (13.33)	12.00 (20.00)	10.00 (16.67)	43.00 (17.92)	

Source : Field Survey, 2012

*Indicates significant at 10 percent level of significance. Figure in the parenthesis indicates percentage.

c) Livestock holders' vulnerability

The result of statistical description and classification of variable indicate (refer Table 3 and Appendix 1) that the socio-economic characteristics vary widely within the agro ecological regions with the highest being recorded for electricity, toilet, mobile or cell phone, and television, radio, access to road, access to market, access to credit, saving, and member of organization. The variation in the adaptive variables is remarkable. The adaptive capacity variables were categorized in to two groups, household assets; social services and facilities. Average number of household assets possessed by the sampled respondents in the tropical was highest (12.07), followed by subtropical (8.82) and least in the warm temperate (8.00). The F value indicated that the distribution was more

heterogeneous and they were significant ($P < 0.01$). There was more disparity in the ownership of assets. Analysis of access to the services and facilities indicated that sampled respondents in the tropical had more access to these services with the mean of 7.68 followed by the subtropical (7.050) and least in the warm temperate region (5.00). Similarly, there was high variation in the exposure variables like drought, flood and landslide within the agro ecological regions. The mean of drought, flood and landslide, being all of these variables Binary, indicated that cool temperate region was found more exposed followed by the warm temperate while tropical zone was least exposed to these climatic extreme events. Moreover, sensitivity variables in the study area included annual temperature variation and annual rainfall variability.

Table 3 : Summary of assets, services exposure and sensitivity variables across the regions

Variables	Agro ecological regions					F- value
	Tropical	Subtropical	Warm temperate	Cool temperate	Total	
Total assets	12.07	8.82	8.00	8.25	9.28	24.072***
Services	7.68	7.05	5.00	5.40	6.28	30.574***
Exposure	0.03	0.40	0.92	1.02	0.59	27.765***
Rainfall	170.39	129.86	129.86	33.46	117.4	130.080***
Temperature	24.35	19.25	19.25	10.98	18.52	98.932***

Source : Field Survey, 2012

*** Significant at 1 percent level of significance

Analysis of annual temperature range showed that warm temperature (Beni Bazar station) had experienced the highest range of 24.50 percent while lowest annual range of 2.60 percent was recorded in tropical (Rampur station). Analysis of annual rainfall variation showed that the coefficient of variation of 26.04 percent in the annual rainfall was highest in the warm and subtropical (Beni Bazar station) while lowest variation was found in the tropical.

The result of Principle Component Analysis (PCA) showed that nine components with the Eigen value of 1 or greater accounting for 67.85 percent of total variance. The first component had an Eigen value of 6.06 and accounted for 21.66 percent followed by the second component with an Eigen value of 2.71 and percent explanation was of 9.91. The analysis also revealed that the components scores as earlier stated only the component score of the first component were used in weighing the variables for the construction of vulnerability indices. The components score are shown in the Appendix 2.

Table 4 : Vulnerability indices across the regions

Agro Ecological Regions	Vulnerability Index
Tropical	8.11
Subtropical	5.62
Warm temperate	3.29
Cool temperate	3.62

The calculation of vulnerability indices showed that all agro ecological regions had positive values implying that they were less vulnerable. However, some regions were in better position to withstand climate change than others. Accordingly, tropical, subtropical, cool temperate and warm temperate region with the indices of 8.1, 5.62, 3.62, 3.29 respectively in the order of increasing vulnerability (Table 4). It could be concluded that warm temperate by its least indices was most vulnerable while tropical region with highest indices indicates less vulnerable to climate change. According to the GIEC report (2007), mountain ecosystems are considered extremely vulnerable to climate change.

Tropical region experienced low vulnerability to climate change because the rural households have high literacy rate, high household income, and have more access to infrastructure and technology. They are characterized by the high degree of nonfarm employment. The diversification of economic activities and access to the infrastructure technology make the household less reliant on livestock which is more sensitive to climate change. It should also be noted that although drought occasionally occurs, flood rarely occurs and landslide hasn't noticed in this region. Temperature variation and rainfall variability was least over last thirty one years and all these explain why tropical is experiencing low vulnerability. A major reason

for high vulnerability of the warm temperate region is the low level of technology and infrastructure development. There is also higher incidence of poverty in more vulnerable region. The high degree of vulnerability in the warm temperate region can in addition be explained by the higher variation in the annual rainfall and temperature variation and more frequent occurrence of occurrence of floods and landslides.

IV. CONCLUSION

Most of livestock keepers had observed the variation on weather patterns and experienced increased temperature, decreased but erratic precipitation and delayed summer monsoon. The result revealed that highly impoverished livestock keepers from the fragile mountain with limited access to basic services, wealth and assets, are most vulnerable to extreme climatic events than those with better access to services and wealth.

Policies should focus on: the investment on basic services; the inception of livestock insurance, provision of agroecology based technology package to enhance their specific adaptation potential; the strengthening of productive safety net programs through the involvement of governmental and Non Governmental organization. Tackling the problem of vulnerability to climate change among the regions in the Nepal which from the differences in a number of physical and socioeconomic factors requires climate change adaptation policies that implemented within the framework of integrated rural development specially focusing on,

- a. Creation of off farm employment opportunities.
- b. Provision of credit, training and extension exposure opportunities and climate information.
- c. Establishment of cooperatives and agricultural group in the rural areas.
- d. Tackling the climate induced hazards like floods, drought and landslide.
- e. Water harvest scheme.
- f. Rural poverty alleviation program.
- g. Income diversification.
- h. Improvement in farming practices.

V. ACKNOWLEDGEMENT

This research was supported by the Feed the Future Innovation Lab collaborative Research Adopting Livestock Systems to Climate Change formerly known as Livestock Climate change, Collaborative Research Support Program, Colorado State University, USA. The author would like to acknowledge the funding agency which has been a valuable learning experience for all of us. The author equally grateful to the farmers of Gandaki River Basin, Nepal for providing invaluable information and shared their indigenous knowledge.

REFERENCES RÉFÉRENCES REFERENCIAS

- (2011). *Central Bureau of Statistics*, Major finding of Census, Kathmandu, Nepal.
- Christensen, L., M. B. Coughenour, J. E. Ellis, and Z. Z. Chen (2004). Vulnerability of the Asian typical steppe to grazing and climate change. *63*, 351-368.
- (2012). *District Agriculture Development Office*, Chitwan District profile, Chitwan, Nepal.
- (2002). *District Development Committee, Information and Documentation Center*. Mustang District Profile.
- Deressa, T. T., R. M. Hassan, C. Ringler T. Alemu and M. Yusuf. (2009). Determinants of farmers' choice of adaptation methods to climate change in the Nile Basin of Ethiopia. *Change* doi:10.1016/j.gloenvcha.2009.01.002.
- (2011a). *District Agriculture Livestock Office*, Myagdi District profile with annual progress report, Myagdi, Nepal.
- (2011b). *District Agriculture Livestock Office*, Livestock service annual report District profile with annual progress report, Mustang, Nepal.
- (2008). Climate-related transboundary pests and diseases including relevant aquatic species.
- Heltberg, R., P. B. Siegel and S. L. Jorgensen. (2009). Addressing human vulnerability to climate change: Toward a 'no-regrets' approach. *19*, 89–99.
- M. L. Parry, O. F. Canziani, J. P. Palutikof, P. J. Linden and CE Hanson (2007). *Impacts, adaptation and vulnerability*. Contribution of working group II to the fourth assessment report of the Intergovernmental Panel on Climate Change (IPCC), UK.
- (2001). Synthesis report. Contribution of working groups I, II and III to the fourth assessment report of the Intergovernmental Panel on Climate Change, Switzerland. 104.
- (2013, July). Analysis: Why livestock matters in Nepal. <http://www.irinnews.org/report/98463/analysis-why-livestock-matters-in-nepal>.
- Madu, I. A. (2012). Spatial vulnerability of rural households to climate change in Nigeria.
- (2011). Climate change risk atlas. UK.
- (2012). *Government of Nepal, Ministry of Agriculture Development, Agribusiness and Statistics Division*, Selected indicators of Nepalese agriculture and population, Nepal.
- Raut A. (2006). Climate change impact on Nepal. *60*, 34-46.
- Stern, N. (2006). The economics of climate change.
- Thornton, P. K. Herrero, M. Freeman, A. Okeyo, M. Rege, E. Jones, P. G. & McDermott, J. (2007). Vulnerability, climate change and livestock-Research opportunities and challenges for poverty alleviation. *Journal of Semi-Arid Tropical Agricultural Research*, 4 (1). online at <http://www.icrisat.org/Journal/specialproject.htm>

Appendix 1 : Description of variables used in vulnerability assessment through PCA

Variables	Tropical	Subtropical	Warm temperate	Cool temperate	Total	F-Value
Upper caste	0.92	0.68	0.10	0.03	0.43	107.52***
Radio	0.70	0.60	0.70	0.60	0.65	0.87
TV	0.92	0.82	0.57	0.60	0.73	9.48***
Mobile	1.00	0.93	0.87	0.83	0.91	4.07***
Cycle	0.97	0.15	0.02	0.03	0.29	233.37***
Computer	0.32	0.25	0.02	0.02	0.15	13.21***
Toilet	0.98	0.97	0.85	0.92	0.93	3.354**
Organization	0.57	0.63	0.28	0.48	0.49	5.84***
Access to credit	0.78	0.92	0.72	0.63	0.76	4.96***
Saving	0.45	0.72	0.48	0.50	0.54	3.65**
100 LSU	0.54	0.22	0.08	0.19	0.26	3.27**
Off farm income	0.15	0.07	0.02	0.15	0.10	1.50
Total income	0.32	0.29	0.12	0.28	0.25	2.59*
Economically active members	4.18	3.63	3.82	3.55	3.80	1.44
Food sufficient	0.93	0.17	0.25	0.53	0.47	44.31***
Land	0.97	0.98	0.92	0.88	0.94	2.18*
Access to road	0.95	0.98	0.75	0.73	0.85	9.03***
Access to market	0.80	0.78	0.72	0.68	0.75	0.95
Access to agrovet	0.27	0.12	0.02	0.00	0.10	11.22***
Educated family	3.37	3.43	1.90	1.78	2.62	10.98***
Quality of house	0.38	0.02	0.07	0.07	0.13	17.72***
Cooking source	0.85	0.33	0.20	0.28	0.42	28.63***

Literacy rate	0.90	0.82	0.68	0.70	0.78	1.85
Rainfall	0.17	0.26	0.26	0.18	0.49	130.08***
Temperature	0.03	0.25	0.25	0.07	0.32	98.93***
Drought	0.03	0.07	0.03	0.17	0.08	3.54**
Flood	0.00	0.15	0.30	0.33	0.20	9.86***
Landslide	0.00	0.18	0.58	0.52	0.32	28.05***

***, ** and * Indicate significant at 1, 5 and 10 percent level of significance.

Appendix 2 : Component scores of coefficient for component first through PCA

Indicators	Types of variables	Component scores
Higher caste	Adaptive capacity	0.742
Ownership of radio	Adaptive capacity	0.247
Ownership of TV	Adaptive capacity	0.471
Ownership of cell phone/telephone	Adaptive capacity	0.279
Ownership of cycle	Adaptive capacity	0.761
Ownership of Computer	Adaptive capacity	0.641
Ownership of Toilet	Adaptive capacity	0.265
Member of organization	Adaptive capacity	0.159
Access to credit	Adaptive capacity	0.128
Saving	Adaptive capacity	0.051
Livestock standard unit (100 LSU)	Adaptive capacity	0.375
Off farm income in NRs	Adaptive capacity	0.443
Total income in NRs	Adaptive capacity	0.445
Economically active family members	Adaptive capacity	0.242
Food sufficient	Adaptive capacity	0.442
Ownership of land	Adaptive capacity	0.301
Access to road	Adaptive capacity	0.341
Access to market	Adaptive capacity	0.186
Access to agrovet	Adaptive capacity	0.560
Educated family member	Adaptive capacity	0.512
Quality of house	Adaptive capacity	0.447
Improved cooking source	Adaptive capacity	0.642
Literacy rate	Adaptive capacity	0.108
Rainfall variability	Sensitivity	0.707
Temperature variation	Sensitivity	0.700
Drought	Exposure	-0.390
Flood	Exposure	-0.610
Landslide	Exposure	-0.555

Extraction method: Principal component analysis. Nine components were extracted
Rotation method: Varimax with Kaiser Normalization.

GLOBAL JOURNALS INC. (US) GUIDELINES HANDBOOK 2013

WWW.GLOBALJOURNALS.ORG

FELLOWS

FELLOW OF ASSOCIATION OF RESEARCH SOCIETY IN SCIENCE (FARSS)

- 'FARSS' title will be awarded to the person after approval of Editor-in-Chief and Editorial Board. The title 'FARSS' can be added to name in the following manner. eg. Dr. John E. Hall, Ph.D., FARSS or William Walldroff Ph. D., M.S., FARSS
- Being FARSS is a respectful honor. It authenticates your research activities. After becoming FARSS, you can use 'FARSS' title as you use your degree in suffix of your name. This will definitely will enhance and add up your name. You can use it on your Career Counseling Materials/CV/Resume/Visiting Card/Name Plate etc.
- 60% Discount will be provided to FARSS members for publishing research papers in Global Journals Inc., if our Editorial Board and Peer Reviewers accept the paper. For the life time, if you are author/co-author of any paper bill sent to you will automatically be discounted one by 60%
- FARSS will be given a renowned, secure, free professional email address with 100 GB of space eg.johnhall@globaljournals.org. You will be facilitated with Webmail, SpamAssassin, Email Forwarders, Auto-Responders, Email Delivery Route tracing, etc.
- FARSS member is eligible to become paid peer reviewer at Global Journals Inc. to earn up to 15% of realized author charges taken from author of respective paper. After reviewing 5 or more papers you can request to transfer the amount to your bank account or to your PayPal account.
- Eg. If we had taken 420 USD from author, we can send 63 USD to your account.
- FARSS member can apply for free approval, grading and certification of some of their Educational and Institutional Degrees from Global Journals Inc. (US) and Open Association of Research,Society U.S.A.
- After you are FARSS. You can send us scanned copy of all of your documents. We will verify, grade and certify them within a month. It will be based on your academic records, quality of research papers published by you, and 50 more criteria. This is beneficial for your job interviews as recruiting organization need not just rely on you for authenticity and your unknown qualities, you would have authentic ranks of all of your documents. Our scale is unique worldwide.
- FARSS member can proceed to get benefits of free research podcasting in Global Research Radio with their research documents, slides and online movies.
- After your publication anywhere in the world, you can upload you research paper with your recorded voice or you can use our professional RJs to record your paper their voice. We can also stream your conference videos and display your slides online.
- FARSS will be eligible for free application of Standardization of their Researches by Open Scientific Standards. Standardization is next step and level after publishing in a journal. A team of research and professional will work with you to take your research to its next level, which is worldwide open standardization.



- FARSS is eligible to earn from their researches: While publishing his paper with Global Journals Inc. (US), FARSS can decide whether he/she would like to publish his/her research in closed manner. When readers will buy that individual research paper for reading, 80% of its earning by Global Journals Inc. (US) will be transferred to FARSS member's bank account after certain threshold balance. There is no time limit for collection. FARSS member can decide its price and we can help in decision.

MEMBER OF ASSOCIATION OF RESEARCH SOCIETY IN SCIENCE (MARSS)

- 'MARSS' title will be awarded to the person after approval of Editor-in-Chief and Editorial Board. The title 'MARSS' can be added to name in the following manner. eg. Dr. John E. Hall, Ph.D., MARSS or William Walldroff Ph. D., M.S., MARSS
- Being MARSS is a respectful honor. It authenticates your research activities. After becoming MARSS, you can use 'MARSS' title as you use your degree in suffix of your name. This will definitely will enhance and add up your name. You can use it on your Career Counseling Materials/CV/Resume/Visiting Card/Name Plate etc.
- 40% Discount will be provided to MARSS members for publishing research papers in Global Journals Inc., if our Editorial Board and Peer Reviewers accept the paper. For the life time, if you are author/co-author of any paper bill sent to you will automatically be discounted one by 60%
- MARSS will be given a renowned, secure, free professional email address with 30 GB of space eg.johnhall@globaljournals.org. You will be facilitated with Webmail, SpamAssassin, Email Forwarders, Auto-Responders, Email Delivery Route tracing, etc.
- MARSS member is eligible to become paid peer reviewer at Global Journals Inc. to earn up to 10% of realized author charges taken from author of respective paper. After reviewing 5 or more papers you can request to transfer the amount to your bank account or to your PayPal account.
- MARSS member can apply for free approval, grading and certification of some of their Educational and Institutional Degrees from Global Journals Inc. (US) and Open Association of Research,Society U.S.A.
- MARSS is eligible to earn from their researches: While publishing his paper with Global Journals Inc. (US), MARSS can decide whether he/she would like to publish his/her research in closed manner. When readers will buy that individual research paper for reading, 40% of its earning by Global Journals Inc. (US) will be transferred to MARSS member's bank account after certain threshold balance. There is no time limit for collection. MARSS member can decide its price and we can help in decision.

AUXILIARY MEMBERSHIPS

ANNUAL MEMBER

- Annual Member will be authorized to receive e-Journal GJSFR for one year (subscription for one year).
- The member will be allotted free 1 GB Web-space along with subDomain to contribute and participate in our activities.
- A professional email address will be allotted free 500 MB email space.

PAPER PUBLICATION

- The members can publish paper once. The paper will be sent to two-peer reviewer. The paper will be published after the acceptance of peer reviewers and Editorial Board.



PROCESS OF SUBMISSION OF RESEARCH PAPER

The Area or field of specialization may or may not be of any category as mentioned in 'Scope of Journal' menu of the GlobalJournals.org website. There are 37 Research Journal categorized with Six parental Journals GJCST, GJMR, GJRE, GJMBR, GJSFR, GJHSS. For Authors should prefer the mentioned categories. There are three widely used systems UDC, DDC and LCC. The details are available as 'Knowledge Abstract' at Home page. The major advantage of this coding is that, the research work will be exposed to and shared with all over the world as we are being abstracted and indexed worldwide.

The paper should be in proper format. The format can be downloaded from first page of 'Author Guideline' Menu. The Author is expected to follow the general rules as mentioned in this menu. The paper should be written in MS-Word Format (*.DOC, *.DOCX).

The Author can submit the paper either online or offline. The authors should prefer online submission. Online Submission: There are three ways to submit your paper:

(A) (I) First, register yourself using top right corner of Home page then Login. If you are already registered, then login using your username and password.

(II) Choose corresponding Journal.

(III) Click 'Submit Manuscript'. Fill required information and Upload the paper.

(B) If you are using Internet Explorer, then Direct Submission through Homepage is also available.

(C) If these two are not convenient, and then email the paper directly to dean@globaljournals.org.

Offline Submission: Author can send the typed form of paper by Post. However, online submission should be preferred.



PREFERRED AUTHOR GUIDELINES

MANUSCRIPT STYLE INSTRUCTION (Must be strictly followed)

Page Size: 8.27" X 11"

- Left Margin: 0.65
- Right Margin: 0.65
- Top Margin: 0.75
- Bottom Margin: 0.75
- Font type of all text should be Swis 721 Lt BT.
- Paper Title should be of Font Size 24 with one Column section.
- Author Name in Font Size of 11 with one column as of Title.
- Abstract Font size of 9 Bold, "Abstract" word in Italic Bold.
- Main Text: Font size 10 with justified two columns section
- Two Column with Equal Column with of 3.38 and Gaping of .2
- First Character must be three lines Drop capped.
- Paragraph before Spacing of 1 pt and After of 0 pt.
- Line Spacing of 1 pt
- Large Images must be in One Column
- Numbering of First Main Headings (Heading 1) must be in Roman Letters, Capital Letter, and Font Size of 10.
- Numbering of Second Main Headings (Heading 2) must be in Alphabets, Italic, and Font Size of 10.

You can use your own standard format also.

Author Guidelines:

1. General,
2. Ethical Guidelines,
3. Submission of Manuscripts,
4. Manuscript's Category,
5. Structure and Format of Manuscript,
6. After Acceptance.

1. GENERAL

Before submitting your research paper, one is advised to go through the details as mentioned in following heads. It will be beneficial, while peer reviewer justify your paper for publication.

Scope

The Global Journals Inc. (US) welcome the submission of original paper, review paper, survey article relevant to the all the streams of Philosophy and knowledge. The Global Journals Inc. (US) is parental platform for Global Journal of Computer Science and Technology, Researches in Engineering, Medical Research, Science Frontier Research, Human Social Science, Management, and Business organization. The choice of specific field can be done otherwise as following in Abstracting and Indexing Page on this Website. As the all Global



Journals Inc. (US) are being abstracted and indexed (in process) by most of the reputed organizations. Topics of only narrow interest will not be accepted unless they have wider potential or consequences.

2. ETHICAL GUIDELINES

Authors should follow the ethical guidelines as mentioned below for publication of research paper and research activities.

Papers are accepted on strict understanding that the material in whole or in part has not been, nor is being, considered for publication elsewhere. If the paper once accepted by Global Journals Inc. (US) and Editorial Board, will become the copyright of the Global Journals Inc. (US).

Authorship: The authors and coauthors should have active contribution to conception design, analysis and interpretation of findings. They should critically review the contents and drafting of the paper. All should approve the final version of the paper before submission

The Global Journals Inc. (US) follows the definition of authorship set up by the Global Academy of Research and Development. According to the Global Academy of R&D authorship, criteria must be based on:

- 1) Substantial contributions to conception and acquisition of data, analysis and interpretation of the findings.
- 2) Drafting the paper and revising it critically regarding important academic content.
- 3) Final approval of the version of the paper to be published.

All authors should have been credited according to their appropriate contribution in research activity and preparing paper. Contributors who do not match the criteria as authors may be mentioned under Acknowledgement.

Acknowledgements: Contributors to the research other than authors credited should be mentioned under acknowledgement. The specifications of the source of funding for the research if appropriate can be included. Suppliers of resources may be mentioned along with address.

Appeal of Decision: The Editorial Board's decision on publication of the paper is final and cannot be appealed elsewhere.

Permissions: It is the author's responsibility to have prior permission if all or parts of earlier published illustrations are used in this paper.

Please mention proper reference and appropriate acknowledgements wherever expected.

If all or parts of previously published illustrations are used, permission must be taken from the copyright holder concerned. It is the author's responsibility to take these in writing.

Approval for reproduction/modification of any information (including figures and tables) published elsewhere must be obtained by the authors/copyright holders before submission of the manuscript. Contributors (Authors) are responsible for any copyright fee involved.

3. SUBMISSION OF MANUSCRIPTS

Manuscripts should be uploaded via this online submission page. The online submission is most efficient method for submission of papers, as it enables rapid distribution of manuscripts and consequently speeds up the review procedure. It also enables authors to know the status of their own manuscripts by emailing us. Complete instructions for submitting a paper is available below.

Manuscript submission is a systematic procedure and little preparation is required beyond having all parts of your manuscript in a given format and a computer with an Internet connection and a Web browser. Full help and instructions are provided on-screen. As an author, you will be prompted for login and manuscript details as Field of Paper and then to upload your manuscript file(s) according to the instructions.



To avoid postal delays, all transaction is preferred by e-mail. A finished manuscript submission is confirmed by e-mail immediately and your paper enters the editorial process with no postal delays. When a conclusion is made about the publication of your paper by our Editorial Board, revisions can be submitted online with the same procedure, with an occasion to view and respond to all comments.

Complete support for both authors and co-author is provided.

4. MANUSCRIPT'S CATEGORY

Based on potential and nature, the manuscript can be categorized under the following heads:

Original research paper: Such papers are reports of high-level significant original research work.

Review papers: These are concise, significant but helpful and decisive topics for young researchers.

Research articles: These are handled with small investigation and applications

Research letters: The letters are small and concise comments on previously published matters.

5. STRUCTURE AND FORMAT OF MANUSCRIPT

The recommended size of original research paper is less than seven thousand words, review papers fewer than seven thousands words also. Preparation of research paper or how to write research paper, are major hurdle, while writing manuscript. The research articles and research letters should be fewer than three thousand words, the structure original research paper; sometime review paper should be as follows:

Papers: These are reports of significant research (typically less than 7000 words equivalent, including tables, figures, references), and comprise:

- (a) Title should be relevant and commensurate with the theme of the paper.
- (b) A brief Summary, "Abstract" (less than 150 words) containing the major results and conclusions.
- (c) Up to ten keywords, that precisely identifies the paper's subject, purpose, and focus.
- (d) An Introduction, giving necessary background excluding subheadings; objectives must be clearly declared.
- (e) Resources and techniques with sufficient complete experimental details (wherever possible by reference) to permit repetition; sources of information must be given and numerical methods must be specified by reference, unless non-standard.
- (f) Results should be presented concisely, by well-designed tables and/or figures; the same data may not be used in both; suitable statistical data should be given. All data must be obtained with attention to numerical detail in the planning stage. As reproduced design has been recognized to be important to experiments for a considerable time, the Editor has decided that any paper that appears not to have adequate numerical treatments of the data will be returned un-refereed;
- (g) Discussion should cover the implications and consequences, not just recapitulating the results; conclusions should be summarizing.
- (h) Brief Acknowledgements.
- (i) References in the proper form.

Authors should very cautiously consider the preparation of papers to ensure that they communicate efficiently. Papers are much more likely to be accepted, if they are cautiously designed and laid out, contain few or no errors, are summarizing, and be conventional to the approach and instructions. They will in addition, be published with much less delays than those that require much technical and editorial correction.



The Editorial Board reserves the right to make literary corrections and to make suggestions to improve brevity.

It is vital, that authors take care in submitting a manuscript that is written in simple language and adheres to published guidelines.

Format

Language: The language of publication is UK English. Authors, for whom English is a second language, must have their manuscript efficiently edited by an English-speaking person before submission to make sure that, the English is of high excellence. It is preferable, that manuscripts should be professionally edited.

Standard Usage, Abbreviations, and Units: Spelling and hyphenation should be conventional to The Concise Oxford English Dictionary. Statistics and measurements should at all times be given in figures, e.g. 16 min, except for when the number begins a sentence. When the number does not refer to a unit of measurement it should be spelt in full unless, it is 160 or greater.

Abbreviations supposed to be used carefully. The abbreviated name or expression is supposed to be cited in full at first usage, followed by the conventional abbreviation in parentheses.

Metric SI units are supposed to generally be used excluding where they conflict with current practice or are confusing. For illustration, 1.4 l rather than $1.4 \times 10^{-3} \text{ m}^3$, or 4 mm somewhat than $4 \times 10^{-3} \text{ m}$. Chemical formula and solutions must identify the form used, e.g. anhydrous or hydrated, and the concentration must be in clearly defined units. Common species names should be followed by underlines at the first mention. For following use the generic name should be constricted to a single letter, if it is clear.

Structure

All manuscripts submitted to Global Journals Inc. (US), ought to include:

Title: The title page must carry an instructive title that reflects the content, a running title (less than 45 characters together with spaces), names of the authors and co-authors, and the place(s) wherever the work was carried out. The full postal address in addition with the e-mail address of related author must be given. Up to eleven keywords or very brief phrases have to be given to help data retrieval, mining and indexing.

Abstract, used in Original Papers and Reviews:

Optimizing Abstract for Search Engines

Many researchers searching for information online will use search engines such as Google, Yahoo or similar. By optimizing your paper for search engines, you will amplify the chance of someone finding it. This in turn will make it more likely to be viewed and/or cited in a further work. Global Journals Inc. (US) have compiled these guidelines to facilitate you to maximize the web-friendliness of the most public part of your paper.

Key Words

A major linchpin in research work for the writing research paper is the keyword search, which one will employ to find both library and Internet resources.

One must be persistent and creative in using keywords. An effective keyword search requires a strategy and planning a list of possible keywords and phrases to try.

Search engines for most searches, use Boolean searching, which is somewhat different from Internet searches. The Boolean search uses "operators," words (and, or, not, and near) that enable you to expand or narrow your affords. Tips for research paper while preparing research paper are very helpful guideline of research paper.

Choice of key words is first tool of tips to write research paper. Research paper writing is an art. A few tips for deciding as strategically as possible about keyword search:



- One should start brainstorming lists of possible keywords before even begin searching. Think about the most important concepts related to research work. Ask, "What words would a source have to include to be truly valuable in research paper?" Then consider synonyms for the important words.
- It may take the discovery of only one relevant paper to let steer in the right keyword direction because in most databases, the keywords under which a research paper is abstracted are listed with the paper.
- One should avoid outdated words.

Keywords are the key that opens a door to research work sources. Keyword searching is an art in which researcher's skills are bound to improve with experience and time.

Numerical Methods: Numerical methods used should be clear and, where appropriate, supported by references.

Acknowledgements: Please make these as concise as possible.

References

References follow the Harvard scheme of referencing. References in the text should cite the authors' names followed by the time of their publication, unless there are three or more authors when simply the first author's name is quoted followed by et al. unpublished work has to only be cited where necessary, and only in the text. Copies of references in press in other journals have to be supplied with submitted typescripts. It is necessary that all citations and references be carefully checked before submission, as mistakes or omissions will cause delays.

References to information on the World Wide Web can be given, but only if the information is available without charge to readers on an official site. Wikipedia and Similar websites are not allowed where anyone can change the information. Authors will be asked to make available electronic copies of the cited information for inclusion on the Global Journals Inc. (US) homepage at the judgment of the Editorial Board.

The Editorial Board and Global Journals Inc. (US) recommend that, citation of online-published papers and other material should be done via a DOI (digital object identifier). If an author cites anything, which does not have a DOI, they run the risk of the cited material not being noticeable.

The Editorial Board and Global Journals Inc. (US) recommend the use of a tool such as Reference Manager for reference management and formatting.

Tables, Figures and Figure Legends

Tables: Tables should be few in number, cautiously designed, uncrowned, and include only essential data. Each must have an Arabic number, e.g. Table 4, a self-explanatory caption and be on a separate sheet. Vertical lines should not be used.

Figures: Figures are supposed to be submitted as separate files. Always take in a citation in the text for each figure using Arabic numbers, e.g. Fig. 4. Artwork must be submitted online in electronic form by e-mailing them.

Preparation of Electronic Figures for Publication

Even though low quality images are sufficient for review purposes, print publication requires high quality images to prevent the final product being blurred or fuzzy. Submit (or e-mail) EPS (line art) or TIFF (halftone/photographs) files only. MS PowerPoint and Word Graphics are unsuitable for printed pictures. Do not use pixel-oriented software. Scans (TIFF only) should have a resolution of at least 350 dpi (halftone) or 700 to 1100 dpi (line drawings) in relation to the imitation size. Please give the data for figures in black and white or submit a Color Work Agreement Form. EPS files must be saved with fonts embedded (and with a TIFF preview, if possible).

For scanned images, the scanning resolution (at final image size) ought to be as follows to ensure good reproduction: line art: >650 dpi; halftones (including gel photographs) : >350 dpi; figures containing both halftone and line images: >650 dpi.



Color Charges: It is the rule of the Global Journals Inc. (US) for authors to pay the full cost for the reproduction of their color artwork. Hence, please note that, if there is color artwork in your manuscript when it is accepted for publication, we would require you to complete and return a color work agreement form before your paper can be published.

Figure Legends: Self-explanatory legends of all figures should be incorporated separately under the heading 'Legends to Figures'. In the full-text online edition of the journal, figure legends may possibly be truncated in abbreviated links to the full screen version. Therefore, the first 100 characters of any legend should notify the reader, about the key aspects of the figure.

6. AFTER ACCEPTANCE

Upon approval of a paper for publication, the manuscript will be forwarded to the dean, who is responsible for the publication of the Global Journals Inc. (US).

6.1 Proof Corrections

The corresponding author will receive an e-mail alert containing a link to a website or will be attached. A working e-mail address must therefore be provided for the related author.

Acrobat Reader will be required in order to read this file. This software can be downloaded

(Free of charge) from the following website:

www.adobe.com/products/acrobat/readstep2.html. This will facilitate the file to be opened, read on screen, and printed out in order for any corrections to be added. Further instructions will be sent with the proof.

Proofs must be returned to the dean at dean@globaljournals.org within three days of receipt.

As changes to proofs are costly, we inquire that you only correct typesetting errors. All illustrations are retained by the publisher. Please note that the authors are responsible for all statements made in their work, including changes made by the copy editor.

6.2 Early View of Global Journals Inc. (US) (Publication Prior to Print)

The Global Journals Inc. (US) are enclosed by our publishing's Early View service. Early View articles are complete full-text articles sent in advance of their publication. Early View articles are absolute and final. They have been completely reviewed, revised and edited for publication, and the authors' final corrections have been incorporated. Because they are in final form, no changes can be made after sending them. The nature of Early View articles means that they do not yet have volume, issue or page numbers, so Early View articles cannot be cited in the conventional way.

6.3 Author Services

Online production tracking is available for your article through Author Services. Author Services enables authors to track their article - once it has been accepted - through the production process to publication online and in print. Authors can check the status of their articles online and choose to receive automated e-mails at key stages of production. The authors will receive an e-mail with a unique link that enables them to register and have their article automatically added to the system. Please ensure that a complete e-mail address is provided when submitting the manuscript.

6.4 Author Material Archive Policy

Please note that if not specifically requested, publisher will dispose off hardcopy & electronic information submitted, after the two months of publication. If you require the return of any information submitted, please inform the Editorial Board or dean as soon as possible.

6.5 Offprint and Extra Copies

A PDF offprint of the online-published article will be provided free of charge to the related author, and may be distributed according to the Publisher's terms and conditions. Additional paper offprint may be ordered by emailing us at: editor@globaljournals.org .



Before start writing a good quality Computer Science Research Paper, let us first understand what is Computer Science Research Paper? So, Computer Science Research Paper is the paper which is written by professionals or scientists who are associated to Computer Science and Information Technology, or doing research study in these areas. If you are novel to this field then you can consult about this field from your supervisor or guide.

TECHNIQUES FOR WRITING A GOOD QUALITY RESEARCH PAPER:

1. Choosing the topic: In most cases, the topic is searched by the interest of author but it can be also suggested by the guides. You can have several topics and then you can judge that in which topic or subject you are finding yourself most comfortable. This can be done by asking several questions to yourself, like Will I be able to carry our search in this area? Will I find all necessary recourses to accomplish the search? Will I be able to find all information in this field area? If the answer of these types of questions will be "Yes" then you can choose that topic. In most of the cases, you may have to conduct the surveys and have to visit several places because this field is related to Computer Science and Information Technology. Also, you may have to do a lot of work to find all rise and falls regarding the various data of that subject. Sometimes, detailed information plays a vital role, instead of short information.

2. Evaluators are human: First thing to remember that evaluators are also human being. They are not only meant for rejecting a paper. They are here to evaluate your paper. So, present your Best.

3. Think Like Evaluators: If you are in a confusion or getting demotivated that your paper will be accepted by evaluators or not, then think and try to evaluate your paper like an Evaluator. Try to understand that what an evaluator wants in your research paper and automatically you will have your answer.

4. Make blueprints of paper: The outline is the plan or framework that will help you to arrange your thoughts. It will make your paper logical. But remember that all points of your outline must be related to the topic you have chosen.

5. Ask your Guides: If you are having any difficulty in your research, then do not hesitate to share your difficulty to your guide (if you have any). They will surely help you out and resolve your doubts. If you can't clarify what exactly you require for your work then ask the supervisor to help you with the alternative. He might also provide you the list of essential readings.

6. Use of computer is recommended: As you are doing research in the field of Computer Science, then this point is quite obvious.

7. Use right software: Always use good quality software packages. If you are not capable to judge good software then you can lose quality of your paper unknowingly. There are various software programs available to help you, which you can get through Internet.

8. Use the Internet for help: An excellent start for your paper can be by using the Google. It is an excellent search engine, where you can have your doubts resolved. You may also read some answers for the frequent question how to write my research paper or find model research paper. From the internet library you can download books. If you have all required books make important reading selecting and analyzing the specified information. Then put together research paper sketch out.

9. Use and get big pictures: Always use encyclopedias, Wikipedia to get pictures so that you can go into the depth.

10. Bookmarks are useful: When you read any book or magazine, you generally use bookmarks, right! It is a good habit, which helps to not to lose your continuity. You should always use bookmarks while searching on Internet also, which will make your search easier.

11. Revise what you wrote: When you write anything, always read it, summarize it and then finalize it.



12. Make all efforts: Make all efforts to mention what you are going to write in your paper. That means always have a good start. Try to mention everything in introduction, that what is the need of a particular research paper. Polish your work by good skill of writing and always give an evaluator, what he wants.

13. Have backups: When you are going to do any important thing like making research paper, you should always have backup copies of it either in your computer or in paper. This will help you to not to lose any of your important.

14. Produce good diagrams of your own: Always try to include good charts or diagrams in your paper to improve quality. Using several and unnecessary diagrams will degrade the quality of your paper by creating "hotchpotch." So always, try to make and include those diagrams, which are made by your own to improve readability and understandability of your paper.

15. Use of direct quotes: When you do research relevant to literature, history or current affairs then use of quotes become essential but if study is relevant to science then use of quotes is not preferable.

16. Use proper verb tense: Use proper verb tenses in your paper. Use past tense, to present those events that happened. Use present tense to indicate events that are going on. Use future tense to indicate future happening events. Use of improper and wrong tenses will confuse the evaluator. Avoid the sentences that are incomplete.

17. Never use online paper: If you are getting any paper on Internet, then never use it as your research paper because it might be possible that evaluator has already seen it or maybe it is outdated version.

18. Pick a good study spot: To do your research studies always try to pick a spot, which is quiet. Every spot is not for studies. Spot that suits you choose it and proceed further.

19. Know what you know: Always try to know, what you know by making objectives. Else, you will be confused and cannot achieve your target.

20. Use good quality grammar: Always use a good quality grammar and use words that will throw positive impact on evaluator. Use of good quality grammar does not mean to use tough words, that for each word the evaluator has to go through dictionary. Do not start sentence with a conjunction. Do not fragment sentences. Eliminate one-word sentences. Ignore passive voice. Do not ever use a big word when a diminutive one would suffice. Verbs have to be in agreement with their subjects. Prepositions are not expressions to finish sentences with. It is incorrect to ever divide an infinitive. Avoid clichés like the disease. Also, always shun irritating alliteration. Use language that is simple and straight forward. put together a neat summary.

21. Arrangement of information: Each section of the main body should start with an opening sentence and there should be a changeover at the end of the section. Give only valid and powerful arguments to your topic. You may also maintain your arguments with records.

22. Never start in last minute: Always start at right time and give enough time to research work. Leaving everything to the last minute will degrade your paper and spoil your work.

23. Multitasking in research is not good: Doing several things at the same time proves bad habit in case of research activity. Research is an area, where everything has a particular time slot. Divide your research work in parts and do particular part in particular time slot.

24. Never copy others' work: Never copy others' work and give it your name because if evaluator has seen it anywhere you will be in trouble.

25. Take proper rest and food: No matter how many hours you spend for your research activity, if you are not taking care of your health then all your efforts will be in vain. For a quality research, study is must, and this can be done by taking proper rest and food.

26. Go for seminars: Attend seminars if the topic is relevant to your research area. Utilize all your resources.



27. Refresh your mind after intervals: Try to give rest to your mind by listening to soft music or by sleeping in intervals. This will also improve your memory.

28. Make colleagues: Always try to make colleagues. No matter how sharper or intelligent you are, if you make colleagues you can have several ideas, which will be helpful for your research.

29. Think technically: Always think technically. If anything happens, then search its reasons, its benefits, and demerits.

30. Think and then print: When you will go to print your paper, notice that tables are not be split, headings are not detached from their descriptions, and page sequence is maintained.

31. Adding unnecessary information: Do not add unnecessary information, like, I have used MS Excel to draw graph. Do not add irrelevant and inappropriate material. These all will create superfluous. Foreign terminology and phrases are not apropos. One should NEVER take a broad view. Analogy in script is like feathers on a snake. Not at all use a large word when a very small one would be sufficient. Use words properly, regardless of how others use them. Remove quotations. Puns are for kids, not grunt readers. Amplification is a billion times of inferior quality than sarcasm.

32. Never oversimplify everything: To add material in your research paper, never go for oversimplification. This will definitely irritate the evaluator. Be more or less specific. Also too, by no means, ever use rhythmic redundancies. Contractions aren't essential and shouldn't be there used. Comparisons are as terrible as clichés. Give up ampersands and abbreviations, and so on. Remove commas, that are, not necessary. Parenthetical words however should be together with this in commas. Understatement is all the time the complete best way to put onward earth-shaking thoughts. Give a detailed literary review.

33. Report concluded results: Use concluded results. From raw data, filter the results and then conclude your studies based on measurements and observations taken. Significant figures and appropriate number of decimal places should be used. Parenthetical remarks are prohibitive. Proofread carefully at final stage. In the end give outline to your arguments. Spot out perspectives of further study of this subject. Justify your conclusion by at the bottom of them with sufficient justifications and examples.

34. After conclusion: Once you have concluded your research, the next most important step is to present your findings. Presentation is extremely important as it is the definite medium through which your research is going to be in print to the rest of the crowd. Care should be taken to categorize your thoughts well and present them in a logical and neat manner. A good quality research paper format is essential because it serves to highlight your research paper and bring to light all necessary aspects in your research.

INFORMAL GUIDELINES OF RESEARCH PAPER WRITING

Key points to remember:

- Submit all work in its final form.
- Write your paper in the form, which is presented in the guidelines using the template.
- Please note the criterion for grading the final paper by peer-reviewers.

Final Points:

A purpose of organizing a research paper is to let people to interpret your effort selectively. The journal requires the following sections, submitted in the order listed, each section to start on a new page.

The introduction will be compiled from reference matter and will reflect the design processes or outline of basis that direct you to make study. As you will carry out the process of study, the method and process section will be constructed as like that. The result segment will show related statistics in nearly sequential order and will direct the reviewers next to the similar intellectual paths throughout the data that you took to carry out your study. The discussion section will provide understanding of the data and projections as to the implication of the results. The use of good quality references all through the paper will give the effort trustworthiness by representing an alertness of prior workings.



Writing a research paper is not an easy job no matter how trouble-free the actual research or concept. Practice, excellent preparation, and controlled record keeping are the only means to make straightforward the progression.

General style:

Specific editorial column necessities for compliance of a manuscript will always take over from directions in these general guidelines.

To make a paper clear

- Adhere to recommended page limits

Mistakes to evade

- Insertion a title at the foot of a page with the subsequent text on the next page
- Separating a table/chart or figure - impound each figure/table to a single page
- Submitting a manuscript with pages out of sequence

In every sections of your document

- Use standard writing style including articles ("a", "the," etc.)
- Keep on paying attention on the research topic of the paper
- Use paragraphs to split each significant point (excluding for the abstract)
- Align the primary line of each section
- Present your points in sound order
- Use present tense to report well accepted
- Use past tense to describe specific results
- Shun familiar wording, don't address the reviewer directly, and don't use slang, slang language, or superlatives
- Shun use of extra pictures - include only those figures essential to presenting results

Title Page:

Choose a revealing title. It should be short. It should not have non-standard acronyms or abbreviations. It should not exceed two printed lines. It should include the name(s) and address (es) of all authors.



Abstract:

The summary should be two hundred words or less. It should briefly and clearly explain the key findings reported in the manuscript-- must have precise statistics. It should not have abnormal acronyms or abbreviations. It should be logical in itself. Shun citing references at this point.

An abstract is a brief distinct paragraph summary of finished work or work in development. In a minute or less a reviewer can be taught the foundation behind the study, common approach to the problem, relevant results, and significant conclusions or new questions.

Write your summary when your paper is completed because how can you write the summary of anything which is not yet written? Wealth of terminology is very essential in abstract. Yet, use comprehensive sentences and do not let go readability for briefness. You can maintain it succinct by phrasing sentences so that they provide more than lone rationale. The author can at this moment go straight to shortening the outcome. Sum up the study, with the subsequent elements in any summary. Try to maintain the initial two items to no more than one ruling each.

- Reason of the study - theory, overall issue, purpose
- Fundamental goal
- To the point depiction of the research
- Consequences, including definite statistics - if the consequences are quantitative in nature, account quantitative data; results of any numerical analysis should be reported
- Significant conclusions or questions that track from the research(es)

Approach:

- Single section, and succinct
- As a outline of job done, it is always written in past tense
- A conceptual should situate on its own, and not submit to any other part of the paper such as a form or table
- Center on shortening results - bound background information to a verdict or two, if completely necessary
- What you account in an conceptual must be regular with what you reported in the manuscript
- Exact spelling, clearness of sentences and phrases, and appropriate reporting of quantities (proper units, important statistics) are just as significant in an abstract as they are anywhere else

Introduction:

The **Introduction** should "introduce" the manuscript. The reviewer should be presented with sufficient background information to be capable to comprehend and calculate the purpose of your study without having to submit to other works. The basis for the study should be offered. Give most important references but shun difficult to make a comprehensive appraisal of the topic. In the introduction, describe the problem visibly. If the problem is not acknowledged in a logical, reasonable way, the reviewer will have no attention in your result. Speak in common terms about techniques used to explain the problem, if needed, but do not present any particulars about the protocols here. Following approach can create a valuable beginning:

- Explain the value (significance) of the study
- Shield the model - why did you employ this particular system or method? What is its compensation? You strength remark on its appropriateness from a abstract point of vision as well as point out sensible reasons for using it.
- Present a justification. Status your particular theory (es) or aim(s), and describe the logic that led you to choose them.
- Very for a short time explain the tentative propose and how it skilled the declared objectives.

Approach:

- Use past tense except for when referring to recognized facts. After all, the manuscript will be submitted after the entire job is done.
- Sort out your thoughts; manufacture one key point with every section. If you make the four points listed above, you will need a least of four paragraphs.



- Present surroundings information only as desirable in order hold up a situation. The reviewer does not desire to read the whole thing you know about a topic.
- Shape the theory/purpose specifically - do not take a broad view.
- As always, give awareness to spelling, simplicity and correctness of sentences and phrases.

Procedures (Methods and Materials):

This part is supposed to be the easiest to carve if you have good skills. A sound written Procedures segment allows a capable scientist to replacement your results. Present precise information about your supplies. The suppliers and clarity of reagents can be helpful bits of information. Present methods in sequential order but linked methodologies can be grouped as a segment. Be concise when relating the protocols. Attempt for the least amount of information that would permit another capable scientist to spare your outcome but be cautious that vital information is integrated. The use of subheadings is suggested and ought to be synchronized with the results section. When a technique is used that has been well described in another object, mention the specific item describing a way but draw the basic principle while stating the situation. The purpose is to text all particular resources and broad procedures, so that another person may use some or all of the methods in one more study or referee the scientific value of your work. It is not to be a step by step report of the whole thing you did, nor is a methods section a set of orders.

Materials:

- Explain materials individually only if the study is so complex that it saves liberty this way.
- Embrace particular materials, and any tools or provisions that are not frequently found in laboratories.
- Do not take in frequently found.
- If use of a definite type of tools.
- Materials may be reported in a part section or else they may be recognized along with your measures.

Methods:

- Report the method (not particulars of each process that engaged the same methodology)
- Describe the method entirely
- To be succinct, present methods under headings dedicated to specific dealings or groups of measures
- Simplify - details how procedures were completed not how they were exclusively performed on a particular day.
- If well known procedures were used, account the procedure by name, possibly with reference, and that's all.

Approach:

- It is embarrassed or not possible to use vigorous voice when documenting methods with no using first person, which would focus the reviewer's interest on the researcher rather than the job. As a result when script up the methods most authors use third person passive voice.
- Use standard style in this and in every other part of the paper - avoid familiar lists, and use full sentences.

What to keep away from

- Resources and methods are not a set of information.
- Skip all descriptive information and surroundings - save it for the argument.
- Leave out information that is immaterial to a third party.

Results:

The principle of a results segment is to present and demonstrate your conclusion. Create this part a entirely objective details of the outcome, and save all understanding for the discussion.

The page length of this segment is set by the sum and types of data to be reported. Carry on to be to the point, by means of statistics and tables, if suitable, to present consequences most efficiently. You must obviously differentiate material that would usually be incorporated in a study editorial from any unprocessed data or additional appendix matter that would not be available. In fact, such matter should not be submitted at all except requested by the instructor.



Content

- Sum up your conclusion in text and demonstrate them, if suitable, with figures and tables.
- In manuscript, explain each of your consequences, point the reader to remarks that are most appropriate.
- Present a background, such as by describing the question that was addressed by creation an exacting study.
- Explain results of control experiments and comprise remarks that are not accessible in a prescribed figure or table, if appropriate.
- Examine your data, then prepare the analyzed (transformed) data in the form of a figure (graph), table, or in manuscript form.

What to stay away from

- Do not discuss or infer your outcome, report surroundings information, or try to explain anything.
- Not at all, take in raw data or intermediate calculations in a research manuscript.
- Do not present the similar data more than once.
- Manuscript should complement any figures or tables, not duplicate the identical information.
- Never confuse figures with tables - there is a difference.

Approach

- As forever, use past tense when you submit to your results, and put the whole thing in a reasonable order.
- Put figures and tables, appropriately numbered, in order at the end of the report
- If you desire, you may place your figures and tables properly within the text of your results part.

Figures and tables

- If you put figures and tables at the end of the details, make certain that they are visibly distinguished from any attach appendix materials, such as raw facts
- Despite of position, each figure must be numbered one after the other and complete with subtitle
- In spite of position, each table must be titled, numbered one after the other and complete with heading
- All figure and table must be adequately complete that it could situate on its own, divide from text

Discussion:

The Discussion is expected the trickiest segment to write and describe. A lot of papers submitted for journal are discarded based on problems with the Discussion. There is no head of state for how long a argument should be. Position your understanding of the outcome visibly to lead the reviewer through your conclusions, and then finish the paper with a summing up of the implication of the study. The purpose here is to offer an understanding of your results and hold up for all of your conclusions, using facts from your research and generally accepted information, if suitable. The implication of result should be visibly described. Infer your data in the conversation in suitable depth. This means that when you clarify an observable fact you must explain mechanisms that may account for the observation. If your results vary from your prospect, make clear why that may have happened. If your results agree, then explain the theory that the proof supported. It is never suitable to just state that the data approved with prospect, and let it drop at that.

- Make a decision if each premise is supported, discarded, or if you cannot make a conclusion with assurance. Do not just dismiss a study or part of a study as "uncertain."
- Research papers are not acknowledged if the work is imperfect. Draw what conclusions you can based upon the results that you have, and take care of the study as a finished work
- You may propose future guidelines, such as how the experiment might be personalized to accomplish a new idea.
- Give details all of your remarks as much as possible, focus on mechanisms.
- Make a decision if the tentative design sufficiently addressed the theory, and whether or not it was correctly restricted.
- Try to present substitute explanations if sensible alternatives be present.
- One research will not counter an overall question, so maintain the large picture in mind, where do you go next? The best studies unlock new avenues of study. What questions remain?
- Recommendations for detailed papers will offer supplementary suggestions.

Approach:

- When you refer to information, differentiate data generated by your own studies from available information
- Submit to work done by specific persons (including you) in past tense.
- Submit to generally acknowledged facts and main beliefs in present tense.



ADMINISTRATION RULES LISTED BEFORE
SUBMITTING YOUR RESEARCH PAPER TO GLOBAL JOURNALS INC. (US)

Please carefully note down following rules and regulation before submitting your Research Paper to Global Journals Inc. (US):

Segment Draft and Final Research Paper: You have to strictly follow the template of research paper. If it is not done your paper may get rejected.

- The **major constraint** is that you must independently make all content, tables, graphs, and facts that are offered in the paper. You must write each part of the paper wholly on your own. The Peer-reviewers need to identify your own perceptives of the concepts in your own terms. NEVER extract straight from any foundation, and never rephrase someone else's analysis.
- Do not give permission to anyone else to "PROOFREAD" your manuscript.
- **Methods to avoid Plagiarism is applied by us on every paper, if found guilty, you will be blacklisted by all of our collaborated research groups, your institution will be informed for this and strict legal actions will be taken immediately.)**
- To guard yourself and others from possible illegal use please do not permit anyone right to use to your paper and files.



CRITERION FOR GRADING A RESEARCH PAPER (COMPILATION)
BY GLOBAL JOURNALS INC. (US)

Please note that following table is only a Grading of "Paper Compilation" and not on "Performed/Stated Research" whose grading solely depends on Individual Assigned Peer Reviewer and Editorial Board Member. These can be available only on request and after decision of Paper. This report will be the property of Global Journals Inc. (US).

Topics	Grades		
	A-B	C-D	E-F
<i>Abstract</i>	Clear and concise with appropriate content, Correct format. 200 words or below	Unclear summary and no specific data, Incorrect form Above 200 words	No specific data with ambiguous information Above 250 words
<i>Introduction</i>	Containing all background details with clear goal and appropriate details, flow specification, no grammar and spelling mistake, well organized sentence and paragraph, reference cited	Unclear and confusing data, appropriate format, grammar and spelling errors with unorganized matter	Out of place depth and content, hazy format
<i>Methods and Procedures</i>	Clear and to the point with well arranged paragraph, precision and accuracy of facts and figures, well organized subheads	Difficult to comprehend with embarrassed text, too much explanation but completed	Incorrect and unorganized structure with hazy meaning
<i>Result</i>	Well organized, Clear and specific, Correct units with precision, correct data, well structuring of paragraph, no grammar and spelling mistake	Complete and embarrassed text, difficult to comprehend	Irregular format with wrong facts and figures
<i>Discussion</i>	Well organized, meaningful specification, sound conclusion, logical and concise explanation, highly structured paragraph reference cited	Wordy, unclear conclusion, spurious	Conclusion is not cited, unorganized, difficult to comprehend
<i>References</i>	Complete and correct format, well organized	Beside the point, Incomplete	Wrong format and structuring



INDEX

A

Amaechi · 29, 35

B

Bureaucracy · 61

C

Caribbean · 53
Catastrophe · 66

D

Degradability · 51, 52
Dichotomized · 57

F

Fabusoro · 18, 20
Fenbendazole · 4, 48
Fisherfolks · 13, 18

H

Herbaceous · 66
Hornandez · 40

I

Incubation · 11
Indemnifying · 56

M

Methionine · 37, 38, 39, 40, 41, 42, 43, 46
Monogastric · 37
Monolithic · 61

N

Narayangadh · 4, 48

P

Potchoiba · 51, 53

Q

Questionnaire · 28, 54, 57, 66, 68

R

Randomized · 2, 4, 37, 39, 46, 48
Randomly · 26, 28, 57
Reconnaissance · 63

S

Schumpeterian · 18
Slaughtered · 51
Soxhlet · 6, 40, 49

T

Transboundary · 75
Tryptophan · 37, 40, 43, 44

U

Unindemnified · 59
Unsavory · 61
Urhobos · 16

V

Venturesomeness · 58
Vigorously · 63

W

Wasexpressed · 57



save our planet



Global Journal of Science Frontier Research

Visit us on the Web at www.GlobalJournals.org | www.JournalofScience.org
or email us at helpdesk@globaljournals.org

ISSN 9755896



© Global Journals