



## Preparation, Characterisation and Reactions of Stable Adducts of Tris (Pentafluorophenyl) Antimony (V) Diisothiocyanates

By Ram Nath Prasad Yadav

*Tribhuvan Uiniversity, Nepal*

**Abstract-** A series of new stable adducts of tris (pentafluorophenyl)antimony(V) diisothiocyanate of the type  $(C_6H_5)_3 Sb(NCS)_2 \cdot L$ . Where, L = dipropyl formamide (DPF), 3-methylpyridine(3-Picoline), dimethyl formamide (DMF), triphenylphosphine oxide( $Ph_3PO$ ), triphenylarsine oxide ( $Ph_3AsO$ ), dimethyl sulfoxide (DMSO), thiourea (TU), pyridine( $C_5H_5N$ ) have been synthesized by the reaction of tris (pentafluorophenyl)antimony(V) diisothiocyanate with desired ligand in anhydrous methanol. Tris(pentafluorophenyl) antimony diisothiocyanate was obtained by the metathetical reaction of tris(pentafluorophenyl) antimony (V) dichloride and potassium thiocyanate in anhydrous methanol. The molecular weight measurement and molar conductance data of the complexes revealed them to be monomeric and non-conducting in nature. Elemental analysis and IR data indicates that the complexes have hexacoordinated octahedral structure.

**Keywords:** tris(pentafluorophenyl) antimony, stable adduct, diisothiocyanate, elemental analysis, IR spectra, hexacoordinate, octahedral.

*GJSFR-B Classification : FOR Code: 039999*



*Strictly as per the compliance and regulations of :*



# Preparation, Characterisation and Reactions of Stable Adducts of Tris (Pentafluorophenyl) Antimony (V) Diisothiocyanates

Ram Nath Prasad Yadav

**Abstract-** A series of new stable adducts of tris (pentafluorophenyl)antimony(V) diisothiocyanate of the type  $(C_6F_5)_3Sb(NCS)_2.L$ . Where, L = dipropyl formamide (DPF), 3-methylpyridine(3-Picoline), dimethyl formamide (DMF), triphenylphosphine oxide( $Ph_3PO$ ), triphenylarsine oxide ( $Ph_3AsO$ ), dimethyl sulfoxide (DMSO), thiourea (TU), pyridine( $C_5H_5N$ ) have been synthesized by the reaction of tris (pentafluorophenyl)antimony(V) diisothiocyanate with desired ligand in anhydrous methanol. Tris(pentafluorophenyl) antimony diisothiocyanate was obtained by the metathetical reaction of tris(pentafluorophenyl) antimony (V) dichloride and potassium thiocyanate in anhydrous methanol. The molecular weight measurement and molar conductance data of the complexes revealed them to be monomeric and non-conducting in nature. Elemental analysis and IR data indicates that the complexes have hexacoordinated octahedral structure.

**Keywords:** tris(pentafluorophenyl) antimony, stable adduct, diisothiocyanate, elemental analysis, IR spectra, hexacoordinate, octahedral.

## I. INTRODUCTION



The Lewis acidity of pentavalent organoantimony compounds,  $R_nSbX_{5-n}$  has extensively been investigated in the last two decades by various groups of workers. The studies are not confined to  $R_2SbCl_3$  and  $RSbCl_4$  having more chlorine content but has been extended to  $R_3SbCl_2$  derivatives as well. The latter class of compounds, based on hydrocarbon ligands are not good acceptors, but the introduction of  $CF_3$  and  $C_6F_5$  groups on to the metal atom (Sb) considerably enhances the Lewis acidity as evident by the formation of hexacoordinate complexes of the type  $(CF_3)_3SbCl_2.L$  and  $(C_6F_5)_3SbCl_2.L$ . The synthesis and stereochemistry of tris(pentafluorophenyl)antimony(V) dichloride with a number of ligands viz. Dimethyl

formamide, diphenyl formamide, triphenylphosphine oxide, arsine oxide, pyridine, picolines, thiourea. etc. has been reported(Singhal et al. 2002). An octahedral environment around antimony has tentatively been proposed for such complexes. On the basis of analytical, and spectroscopic data, it may be noted that except for a single reference on the formation and characterisation of  $(C_6F_5)_3SbCl_2.L$ , no other study related to the synthesis of molecular adducts has been reported to date (Agarwal 1990).

In view of our interest in the chemistry of pentafluorophenylantimony (III and V) derivatives (Yadav 2012<sup>a</sup>, Yadav 2013<sup>a</sup>, Yadav 2013<sup>b</sup> and Yadav 2014) and various other aspects of fluorocarbon based organoantimony compounds including their antimicrobial and antitumour activity (Yadav 2012<sup>b</sup> and Yadav 2013<sup>c</sup>), coupled with the paucity of published data in the field, we have synthesised a series of neutral adducts of tris (pentafluorophenyl) antimony (V) diisothiocyanate,  $(C_6F_5)_3Sb(NCS)_2$ , with oxygen, nitrogen and sulphur donor Lewis bases. A few complexes of  $(C_6F_5)_3SbCl_2$  have also been synthesised for the sake of comparison. The results of this investigation are reported in this paper.

## II. RESULT AND DISCUSSION

Tris (pentafluorophenyl) antimony (v) diisothiocyanate obtained by the metathetical reaction of  $(C_6F_5)_3SbCl_2$  and potassium thiocyanate, recrystallized and dried before use, was treated with the desired ligand in equivalent molar ratio in anhydrous methanol. The reactions were carried out under anhydrous oxygen free conditions.



Where, L = DPF, 3-Picoline, DMF,  $Ph_3PO$ ,  $Ph_3AsO$ , DMSO, TU,  $C_5H_5N$

All the reactions were found to proceed smoothly under mild conditions. The completion of the reaction takes place within 3 hrs. In most of the cases products were obtained as solid after

evaporating the solvent which were crystallized with petroleum ether (40-60°C) or the mixture of diethyl ether and petroleum ether (60-80°). The complexes are soluble in common organic solvents such as chloroform, acetonitrile etc. They show monomeric constitution in freezing benzene. The complexes are

**Author:** Department of Chemistry, Tribhuvan University, Thakur Ram Multiple Campus, Birganj, Nepal. e-mail: ramnathy@rocketmail.com

stable, non-susceptible to oxygen and can be stored for several weeks without decomposition. The constancy in melting point after repeated crystallization as well as TLC run in polar solvent with a single spot excluded the presence of mixture of reactants. Elemental analysis, conductance and molecular weight data are given in table (2 & 3) and correspond well to the proposed formulation of the complexes. The observed values of molecular weight indicate their monomeric constitution while the values of molar conductance of  $10^{-3}$ M solution in acetonitrile ranges between  $20\text{--}30\text{ Ohm}^{-1}\text{ cm}^2\text{ mol}^{-1}$  at room temperature ( $30^\circ\text{C}$ ) which shows the absence of ionic species in solution.

#### a) Infrared Spectra

All the complexes, listed in Table 1, were characterised in the solid state by their infrared spectra in the region  $4000\text{--}200\text{ cm}^{-1}$ . Important IR frequencies for the complexes together with their assignments are listed in Table 4. These assignments have been made by comparing the spectra in  $4000\text{--}200\text{ cm}^{-1}$  region in the solid state of the complexes with those of the free ligands. The infrared absorptions due to pentafluorophenyl groups bonded to antimony are almost identical and do not differ significantly from those observed for other pentafluorophenyl antimony compounds reported earlier from this laboratory (Premraj et al. 1989).

#### b) Infrared Spectra of the Adducts with Oxygen Donors

The  $\nu(\text{C}=\text{O})$  modes in various amides bases appearing at  $1650 \pm 15\text{ cm}^{-1}$  undergo negative shift and are identified at  $1608 \pm 10\text{ cm}^{-1}$  in the spectra of the adducts suggesting weakening of the  $\text{C}=\text{O}$  bond and coordination through the oxygen atom of the base. On the basis of  $-\Delta\nu(\text{CO})$ , the DPF was found better donors as compared to DMF (Premraj & Mishra 1991).

An absorption of strong intensity for  $\nu(\text{S}=\text{O})$ ,  $\nu'(\text{AS}=\text{O})$  and  $\nu(\text{P}=\text{O})$  lying at  $1045$ ,  $880$  and  $1195\text{ cm}^{-1}$  respectively, in the spectra of the free ligands undergoes a distinct negative shift on complexation. The corresponding absorption in the spectra of the adducts appears at  $940$ ,  $835$  and  $1162\text{ cm}^{-1}$  suggesting coordination from oxygen atom of the base. The relative donor abilities of the ligand as apparent from the value of  $-\Delta\nu(\text{C}=\text{O})$ , follow the sequence  $\text{DMSO} > \text{Ph}_3\text{AsO} > \text{Ph}_3\text{PO}$ . On the basis of present and some previous studies a medium strong band in the region  $380\text{--}410\text{ cm}^{-1}$  is assigned to  $\nu(\text{Sb-O})$  stretching frequency (Premraj & Mishra 1991).

#### c) Infrared Spectra of the Adducts with Nitrogen Donors

The  $\nu(\text{CN})$  frequency in  $(\text{C}_6\text{F}_5)_3\text{SbCl}_2\text{Py}$  and  $(\text{C}_6\text{F}_5)_3\text{SbCl}_2\cdot 3\text{-Pic}$  is seen to decrease significantly to  $1610 \pm 5\text{ cm}^{-1}$ . In addition to this a band at  $3310 \pm 10\text{ cm}^{-1}$  assignable to  $\nu(\text{NH})$  mode in free ligand is shifted to slightly lower frequency  $3010 \pm 20\text{ cm}^{-1}$  (Premraj &

Mishra 1991). In the IR spectra of the ligand the assignment of the  $\text{Sb-N}$  bond is tentatively assigned at about  $385 \pm 5\text{ cm}^{-1}$ .

#### d) Infrared Spectra of the Adducts with Sulphur Donor

In sulphur donor ligand (TU) an absorption at  $1069\text{ cm}^{-1}$  reported to possess equal contribution from  $\nu(\text{CN})$  and  $\nu(\text{CS})$ . This remains unaffected on adduct formation and appears at  $1075\text{ cm}^{-1}$ . When coordination occurs through sulphur atom, the  $\nu(\text{CN})$  suffers a positive shift while the  $\nu(\text{CS})$  suffers an almost equal negative shift. As a consequence to this the resulting absorption remains apparently unchanged (Premraj & Mishra 1991). The positive shift of  $\nu(\text{NH})$  from  $3360\text{ cm}^{-1}$  and  $3300\text{ cm}^{-1}$  in free ligand to  $3410$  and  $3370\text{ cm}^{-1}$  in its adduct indicates absence of coordination through N-atom of the ligand and indirectly suggest  $\text{Sb} \leftarrow \text{S}$  bonding. However, on the basis of some previous observation and present studies, the  $(\text{Sb-S})$  bond is assigned at  $380\text{ cm}^{-1}$  (Premraj & Mishra 1991).

The diagnostic frequencies due to NCS group bound to antimony appear around at  $2080$ ,  $840$  and  $475\text{ cm}^{-1}$  which could be attributed to asymmetric (NCS), symmetric (C-S) and bonding mode  $\delta$  NCS, respectively. The pattern and intensity does not show any significant change reported earlier for  $(\text{C}_6\text{F}_5)_3\text{Sb}(\text{NCS})_2$  compounds (Premraj et al. 1985).  $\text{Sb-C}$  bond appears in the range  $445\text{--}465\text{ cm}^{-1}$  (Hall & Sowerby 1988 and Nunn et al. 1996).

#### e) Stereochemistry of the Neutral Molecular Adducts $(\text{Rf})_3\text{Sb}(\text{NCS})_2\cdot\text{L}$

It has been assumed that the addition of a Lewis base, L, to the central atom in a trigonal bipyramidal molecule takes place in a trigonal plane and steric and electrostatic factor play an important role in determining the position of entry of L. It is well established that the more electronegative group goes to the axial position and less electronegative on equatorial positions. Therefore, base L should settle in the equatorial position. It is also supported by a tentative assignment of  $\text{Sb-N}$  band at  $326\text{ cm}^{-1}$  appearing in all the spectra and attributed to the NCS present in the axial positions.

In view of the above idea the nucleophilic attack at the position between the two fluoro groups to produce structure (Meinema & Noltes 1976) appears to be most favourable, since Rf is less electronegative than any halogen atom directly bonded to metal.

Thus analytical, conductance measurement, molecular weight determination and IR data clearly indicates that the newly synthesized complexes have hexacoordination environment around antimony with octahedral configuration as has been suggested for  $\text{R}_2\text{SbCl}_3\cdot\text{L}$  complexes (Premraj & Mishra 1991). It is generally accepted that the tris(pentafluorophenyl)-antimony(V) diisothiocyanate have a geometry of a

trigonal bipyramidal with two halogen atoms occupying apical positions. In adduct formation as indicated, antimony atom increases its coordination number to six, for our hexacoordinate complexes. A tentative assignments of octahedral structure may be represented as below.




Figure : Suggested structure of  $(R_f)_3Sb(NCS)_2L$

$R_f = C_6F_5$  and  $L = DPF, 3\text{-Picoline, DMF, } Ph_3PO, Ph_3AsO, DMSO, TU, C_5H_5N$

### III. EXPERIMENTAL

Tris (pentafluorophenyl) antimony (v) dichloride was prepared by passing chlorine gas into the solution of tris(pentafluorophenyl) antimony in pet-ether and tris (pentafluorophenyl) antimony(V) diisothiocyanate was prepared by the metathesis of tris(pentafluorophenyl)-antimony(V) dichloride and potassium thiocyanate. All the ligands were of reagent grade and used without further purification. All solvents were purified and dried by standard procedures (Vogel 1971).

The molar conductance of 10 — 3M solutions was determined at 25°C with a PR-9500 Philips conductivity assembly. Molecular weights were determined cryoscopically in benzene using a Beckmann thermometer of  $\pm 0.01^\circ C$  accuracy. The stoichiometry of the compounds was established by elemental analysis. Percentage of C, H and N of the compounds was obtained on a semi-microscale (using elemental analyser Carlo Eaba 1106, Thomas CH and Coleman analyser).

Details of the typical representative experiments are described below, the analytical data, molar conductance and IR assignments are summarised in Table 1- 4.

#### a) Reaction of $(C_6F_5)_3Sb(NCS)_2$ with DMF Ligand (5)

In an oxygen free atmosphere a solution of tris(pentafluorophenyl) antimony(V) diisothiocyanate (0.7388 g, 0.5 mmol) in methanol (25 ml) and DMF (0.0731 g, 0.5 mmol) in the same solvent (25 ml) were stirred together at 80°C for 3 h. After that it was filtered off. The filtrate on concentration in vacuo yielded a white crystalline solid and was recrystallised from petroleum ether (40-60°C) to afford tris(pentafluorophenyl) antimony diisothiocyanate dimethyl formamide adduct  $(C_6F_5)_3Sb(NCS)_2.DMF$ . M.P.: 197°C, Yield: 0.536 g, (66%).

#### b) Reaction of $(C_6F_5)_3Sb(NCS)_2$ with $Ph_3PO$ Ligand (6)

A solution of tris(pentafluorophenyl)antimony(V) diisothiocyanate (0.7388 g, 0.5 mmol) in methanol (25 ml) and a solution of the same solvent (25 ml) of triphenyl phosphine oxide (0.278 g, 0.5 mmol) were stirred together at 80°C for 3 h under nitrogen. It was filtered off and the filtrate on concentration in vacuo afforded a light brown solid and was recrystallised from solvent ether to give tris(pentafluorophenyl)- antimony-diisothiocyanate triphenylphosphine oxide adduct. M.P.: 200°C Yield: 0.569g, (56%).

Table 1 : Preparation and Properties of Stable Adducts of Tris (Pentafluorophenyl)Antimony(V) Diisothiocyanates

| S. No. | Complex<br>$(R_f)_3Sb(NCS)_2.L$          | Molar<br>Ratio | Ligand (g)<br>Solvent (ml)               | $(R_f)_3Sb(NCS)_2(g)$<br>Solvent (ml) | M.P.<br>(°C) | Colour      | Recrystallisation<br>solvents |
|--------|--|----------------|--|---------------------------------------|--------------|-------------|-------------------------------|
| 1.     | $R_f = C_6F_5$<br>$(R_f)_3Sb(NCS)_2.DPF$ | 1:1            | DPF (0.129)<br>MeOH (25)                 | 0.7388<br>MeOH (25)                   | 195          | White       | Petroleum ether<br>(40-60°C)  |
| 2.     | $(R_f)_3Sb(NCS)_2.\alpha\text{-Pic}$     | 1:1            | $\alpha\text{-Pic}$ (0.073)<br>MeOH (25) | 0.7388<br>MeOH (25)                   | 204          | Light brown | Petroleum ether<br>(40-60°C)  |
| 3.     | $(R_f)_3Sb(NCS)_2.\beta\text{-Pic}$      | 1:1            | $\beta\text{-Pic}$ (0.073)<br>MeOH (25)  | 0.7388<br>MeOH (25)                   | 170          | Light brown | Petroleum ether<br>(40-60°C)  |
| 4.     | $(R_f)_3Sb(NCS)_2.\gamma\text{-Pic}$     | 1:1            | $\gamma\text{-Pic}$ (0.073)<br>MeOH (25) | 0.7388<br>MeOH (25)                   | 188          | Light brown | Petroleum ether<br>(40-60°C)  |
| 5.     | $(R_f)_3Sb(NCS)_2.DMF$                   | 1:1            | DMF (0.073)<br>MeOH (25)                 | 0.7388<br>MeOH (25)                   | 197          | White       | Petroleum ether<br>(40-60°C)  |
| 6.     | $(R_f)_3Sb(NCS)_2.Ph_3PO$                | 1:1            | $Ph_3PO$ (0.278)<br>MeOH (25)            | 0.7388<br>MeOH (25)                   | 200          | Light brown | Petroleum ether<br>(40-60°C)  |
| 7.     | $(R_f)_3Sb(NCS)_2.Ph_3AsO$               | 1:1            | $Ph_3AsO$ (0.322)<br>MeOH (25)           | 0.7388<br>MeOH (25)                   | 165          | White       | Petroleum ether<br>(40-60°C)  |

|     |   |     |                              |                     |     |             |                              |
|-----|---|-----|------------------------------|---------------------|-----|-------------|------------------------------|
| 8.  | (R <sub>f</sub> ) <sub>3</sub> Sb(NCS) <sub>2</sub> .DMSO | 1:1 | DMSO (0.078)<br>MeOH (25)    | 0.7388<br>MeOH (25) | 180 | White       | Petroleum ether<br>(40-60°C) |
| 9.  | (R <sub>f</sub> ) <sub>3</sub> Sb(NCS) <sub>2</sub> .TU   | 1:1 | TU (0.076)<br>MeOH (25)      | 0.7388<br>MeOH (25) | 183 | Off white   | Petroleum ether<br>(40-60°C) |
| 10. | (R <sub>f</sub> ) <sub>3</sub> Sb(NCS) <sub>2</sub> .Py.  | 1:1 | Py. (0.079)<br>Methanol (25) | 0.7388<br>MeOH (25) | 207 | Light brown | Petroleum ether<br>(40-60°C) |

Table 2 : Elemental Analysis of Stable Adducts of Tris (Pentafluorophenyl) Antimony (V) Diisothiocyanates

| S.No. | Complex  | Empirical formula   | Found (Calcd) (%) |             |             |
|-------|--|---|-------------------|-------------|-------------|
|       |  |   | C                 | H           | N           |
| 1.    | (C <sub>6</sub> F <sub>5</sub> ) <sub>3</sub> Sb(NCS) <sub>2</sub> ·(C <sub>3</sub> H <sub>7</sub> ) <sub>2</sub> HCON | C <sub>27</sub> H <sub>15</sub> F <sub>15</sub> N <sub>3</sub> OS <sub>2</sub> Sb   | 37.34 (37.39)     | 1.73 (1.79) | 4.84 (4.88) |
| 2.    | (C <sub>6</sub> F <sub>5</sub> ) <sub>3</sub> Sb(NCS) <sub>2</sub> ·C <sub>6</sub> H <sub>7</sub> N                    | C <sub>26</sub> H <sub>7</sub> F <sub>15</sub> N <sub>3</sub> S <sub>2</sub> Sb     | 37.51 (37.60)     | 0.84 (0.90) | 5.05 (5.08) |
| 3.    | (C <sub>6</sub> F <sub>5</sub> ) <sub>3</sub> Sb(NCS) <sub>2</sub> ·C <sub>6</sub> H <sub>7</sub> N                    | C <sub>26</sub> H <sub>7</sub> F <sub>15</sub> N <sub>3</sub> S <sub>2</sub> Sb     | 37.51 (37.60)     | 0.84 (0.90) | 5.05 (5.08) |
| 4.    | (C <sub>6</sub> F <sub>5</sub> ) <sub>3</sub> Sb(NCS) <sub>2</sub> ·C <sub>6</sub> H <sub>7</sub> N                    | C <sub>26</sub> H <sub>7</sub> F <sub>15</sub> N <sub>3</sub> S <sub>2</sub> Sb     | 37.51 (37.60)     | 0.84 (0.90) | 5.05 (5.08) |
| 5.    | (C <sub>6</sub> F <sub>5</sub> ) <sub>3</sub> Sb(NCS) <sub>2</sub> ·HCON(CH <sub>3</sub> ) <sub>2</sub>                | C <sub>23</sub> H <sub>7</sub> F <sub>15</sub> N <sub>3</sub> OS <sub>2</sub> Sb    | 34.00 (34.15)     | 0.86 (0.88) | 5.17 (5.20) |
| 6.    | (C <sub>6</sub> F <sub>5</sub> ) <sub>3</sub> Sb(NCS) <sub>2</sub> ·(C <sub>6</sub> H <sub>5</sub> ) <sub>3</sub> PO   | C <sub>38</sub> H <sub>15</sub> F <sub>15</sub> N <sub>2</sub> OS <sub>2</sub> Sb   | 44.85 (44.92)     | 1.48 (1.52) | 2.75 (2.78) |
| 7.    | (C <sub>6</sub> F <sub>5</sub> ) <sub>3</sub> Sb(NCS) <sub>2</sub> ·(C <sub>6</sub> H <sub>5</sub> )AsO                | C <sub>38</sub> H <sub>15</sub> F <sub>15</sub> N <sub>2</sub> OS <sub>2</sub> AsSb | 42.99 (43.02)     | 1.41 (1.45) | 2.64 (2.70) |
| 8.    | (C <sub>6</sub> F <sub>5</sub> ) <sub>3</sub> Sb(NCS) <sub>2</sub> ·(CH <sub>3</sub> ) <sub>2</sub> SO                 | C <sub>22</sub> H <sub>6</sub> F <sub>15</sub> N <sub>2</sub> OS <sub>3</sub> Sb    | 32.32 (32.37)     | 0.73 (0.78) | 3.43 (3.46) |
| 9.    | (C <sub>6</sub> F <sub>5</sub> ) <sub>3</sub> Sb(NCS) <sub>2</sub> ·NH <sub>2</sub> CSNH <sub>2</sub>                  | C <sub>21</sub> H <sub>4</sub> F <sub>15</sub> N <sub>4</sub> S <sub>3</sub> Sb     | 30.93 (30.98)     | 0.49 (0.54) | 6.87 (6.89) |
| 10.   | (C <sub>6</sub> F <sub>5</sub> ) <sub>3</sub> Sb (NCS) <sub>2</sub> ·C <sub>5</sub> H <sub>5</sub> N                   | C <sub>25</sub> H <sub>5</sub> F <sub>15</sub> N <sub>3</sub> S <sub>2</sub> Sb     | 36.69 (36.72)     | 0.61 (0.68) | 5.14 (5.22) |

Table 3 : Molecular weight, conductance measurement and yield of stable adducts of Tris(pentafluorophenyl)antimony(V) diisothiocyanates

| S. No. | Complex   | Molar conductance<br>(Ohm <sup>-1</sup> cm <sup>2</sup> mol <sup>-1</sup> )<br>acetonitrile | Molecular weight in<br>nitrobenzene Found<br>(calcd). | Yield |    |
|--------|---|---|---|-------|----|
|        |   |   |   | g     | %  |
| 1.     | (C <sub>6</sub> F <sub>5</sub> ) <sub>3</sub> Sb(NCS) <sub>2</sub> ·HCON(C <sub>3</sub> H <sub>7</sub> ) <sub>2</sub> | 20.6  | 865.70 (867.75)                                       | 0.555 | 64 |
| 2.     | (C <sub>6</sub> F <sub>5</sub> ) <sub>3</sub> Sb(NCS) <sub>2</sub> ·α-C <sub>6</sub> H <sub>7</sub> N                 | 22.2  | 825.75 (831.75)                                       | 0.574 | 69 |
| 3.     | (C <sub>6</sub> F <sub>5</sub> ) <sub>3</sub> Sb(NCS) <sub>2</sub> ·β-C <sub>6</sub> H <sub>7</sub> N                 | 22.3  | 825.75 (831.75)                                       | 0.582 | 70 |
| 4.     | (C <sub>6</sub> F <sub>5</sub> ) <sub>3</sub> Sb(NCS) <sub>2</sub> ·γ-C <sub>6</sub> H <sub>7</sub> N                 | 22.7  | 825.75 (831.75)                                       | 0.590 | 71 |
| 5.     | (C <sub>6</sub> F <sub>5</sub> ) <sub>3</sub> Sb(NCS) <sub>2</sub> ·HCON(CH <sub>3</sub> ) <sub>2</sub>               | 28.9  | 805.10 (811.75)                                       | 0.536 | 66 |
| 6.     | (C <sub>6</sub> F <sub>5</sub> ) <sub>3</sub> Sb(NCS) <sub>2</sub> ·(C <sub>6</sub> H <sub>5</sub> ) <sub>3</sub> PO  | 25.2  | 1014.75 (1016.75)                                     | 0.569 | 56 |
| 7.     | (C <sub>6</sub> F <sub>5</sub> ) <sub>3</sub> Sb(NCS) <sub>2</sub> ·(C <sub>6</sub> H <sub>5</sub> ) <sub>3</sub> AsO | 24.4  | 1069.60 (1060.67)                                     | 0.605 | 57 |
| 8.     | (C <sub>6</sub> F <sub>5</sub> ) <sub>3</sub> Sb(NCS) <sub>2</sub> ·(CH <sub>3</sub> ) <sub>2</sub> SO                | 27.6  | 812.75 (816.75)                                       | 0.564 | 69 |
| 9.     | (C <sub>6</sub> F <sub>5</sub> ) <sub>3</sub> Sb(NCS) <sub>2</sub> ·NH <sub>2</sub> CSNH <sub>2</sub>                 | 30.8  | 813.75 (814.75)                                       | 0.578 | 71 |
| 10.    | (C <sub>6</sub> F <sub>5</sub> ) <sub>3</sub> Sb (NCS) <sub>2</sub> ·C <sub>5</sub> H <sub>5</sub> N                  | 29.9  | 815 (817.75)  | 0.589 | 72 |

Table 4 : IR Spectra for (R<sub>f</sub>)<sub>3</sub>Sb(NCS)<sub>2</sub>L (Cm<sup>-1</sup>)

| Compd. No.<br>(Adduct) | ν(Sb-C) | ν(Sb-S)/(Sb-O)/ (Sb-N) | ν(C=N)/ (S=O)/<br>ν(P-O)/(N-H)/(As-O)<br>ligand (complex) |
|------------------------|---------|------------------------|---|
| 1                      | 458 ms  | 385ms                  | 1660 (1612)   |
| 2                      | 461 ms  | 382                    | 1615  |
| 3                      | 458 ms  | 384 w                  | 1612  |
| 4                      | 445 ms  | 320 w                  | 1610  |
| 5                      | 447 ms  | 395 ms                 | 1660 (1615)   |
| 6                      | 465 ms  | 405 ms                 | 1195 (1162)   |
| 7                      | 459 ms  | 390 ms                 | 880 (835)   |
| 8                      | 450 ms  | 380 ms                 | 1045 (940)  |
| 9                      | 455 ms  | 382 w                  | 3300 (3368)   |
| 10                     | 459 ms  | 381 w                  | 1612  |

ms = medium strong, w = weak

#### IV. ACKNOWLEDGEMENT

The author is thankful to the Head, Department of chemistry, Lucknow University for providing necessary laboratory facilities; the Director, Regional

Sophisticated Instrumentation Centre, CDRI, Lucknow for microanalysis and spectral data. Thanks are also due to Dr. Premraj, Professor of Chemistry, Lucknow University, Lucknow India for his valuable suggestions.



## REFERENCES RÉFÉRENCES REFERENCIAS

1. Agarwal A. K. (1990). Studies on organo-antimony (III) and antimony(V) derivatives. *Ph. D. dissertation* University of Lucknow Lucknow India.
2. Hall, M., and Sowerby, D. B. (1988). Phenylchloroantimo (III) nates; their preparations and the crystal structure of  $\text{Me}_4\text{N}[\text{PhSbCl}_3]$ ,  $[\text{Hpy}]_2[\text{PhSbCl}_4]$  and  $\text{Me}_4\text{N}[\text{Ph}_2\text{SbCl}_2]$ . *J. Organomet. Chem.* 347, 59-70.
3. Meinema H. A. and Noltes J. G. (1976). Complexes of monochlorodiorganoantimony(V) moieties with dianionic oxo tridentate ligands. *J. Organomet. Chem.* 107, 249-255.
4. Nunn, M., Begley, M. J. and Sowerby, D. B. (1996). Organoantimony(III) and (V) halides with nitrogen donors. *Polyhedron*. 15, 3167-3174.
5. Premraj, Mishra, N. (1991). Preparation and characterization of molecular adduct of diarylantimony(V) chloride,  $\text{R}_2\text{SbCl}_3\text{L}$ . *Ind. J. Chem.*, 30A, 901-903.
6. Premraj, Agarwal, A. K. and Saxena, A. K. (1989). The preparation of pentafluorophenyl antimony(III) and antimony(V) halides and mixed halides. *J. Fluorine Chem.*, 42, 163-172.
7. Premraj, Saxena A. K., Singhal K. and Ranjan A. (1985). Synthesis and some reactions of tris(pentafluorophenyl)antimony compounds. *Polyhedron*. 4, 251-258.
8. Singhal K., Yadav R.N.P., Premraj and Agarwal A. K. (2002). On the Lewis acidity of tris(pentafluorophenyl)antimony(V) dichloride towards neutral monodentate O, N and S donor ligands. *J. Fluorine Chem.* 121, 131-134.
9. Vogel A.I. (1971). *Practical Organic Chemistry*, 5<sup>th</sup> edition, Longmans London.
10. Yadav R.N.P. (2012<sup>a</sup>). Synthesis and characterization of anionic complexes of pentafluorophenylantimony (III) chloride,  $(\text{R}_i)_n\text{SbCl}_{3-n}$ . *Nep. J. of Integrated Sciences*. 2(1), 14-19.
11. Yadav R.N.P. (2012<sup>b</sup>). Metals in oncology: An overview. *Academic Voices* 2(1), 54-58.
12. Yadav R.N.P (2013<sup>a</sup>). Synthesis, characterization and reactions of cationic complexes of arylantimony(III) chlorides  $\text{Ar}_n\text{SbCl}_{3-n}$  with perchlorate and tetrafluoroborate anions. *Int. J. Appl. Sci. Biotechnol.* 1(2), 42-48.
13. Yadav R.N.P (2013<sup>b</sup>). Preparation and characterization of complex bis(pentafluorophenyl)-antimony(V) anions of the type  $[(\text{R}_i)_2\text{SbCl}_3\text{X}]^-$ . *Int. J. Appl. Sci. Biotechnol.* 1(3), 90-96.
14. Yadav R.N.P (2013<sup>c</sup>). Insecticidal and antimicrobial activity of triarylantimony(V) derivatives. *Academic Voices*. 3(1), 40-45.
15. Yadav R.N.P. (2014). Synthesis, characterization and reactions of some pentafluorophenylantimony (III) diamides. *Int.J.Adv. Res.Biol.Sci.* 1(2), 108-114.



This page is intentionally left blank