



GLOBAL JOURNAL OF SCIENCE FRONTIER RESEARCH: H  
ENVIRONMENT & EARTH SCIENCE

Volume 14 Issue 3 Version 1.0 Year 2014

Type : Double Blind Peer Reviewed International Research Journal

Publisher: Global Journals Inc. (USA)

Online ISSN: 2249-4626 & Print ISSN: 0975-5896

## A Study on Eco-Physiology of *Spirulina* in Relation to some Environmental Parameters

By Sunita Verma, Divya Tiwari & Ajay Verma

*Christ Church Collewege, India*

**Abstract-** Physico-chemical characteristics of Surajkund water body in Varanasi (UP) were monitored. High levels of nitrogen (N), phosphorus (P), calcium (Ca), potassium (K) and sodium (Na) indicated that pond was eutrophic. However, contrast to nitrate, ammonia-nitrogen was high in summer and low in winter. The high concentrations of sodium in summer stimulated uptake and transport of CO<sub>2</sub> and HCO<sub>3</sub> at high pH. Highly significant correlation was observed among studied traits. NO<sub>3</sub> exhibited significant negative relation with all traits, though the magnitude varied from NO<sub>2</sub> to NH<sub>4</sub>. More over positive correlation observed among traits only exception was NO<sub>3</sub>. Calcium, Potassium, Sodium etc showed significant positive correlation with other traits except NO<sub>3</sub>. Positive correlation maintained by NO<sub>2</sub> with other traits where negative value with NO<sub>3</sub>. Surajkund Microcystisaeruginosa was dominant plankton in the month of December January and February. Spirulina sp. was found along with the population of Oscillatoria and Chlamydomonas in April, May and June. After 15 days of growth, protein, dry weight, carbohydrate, chlorophyll and carotenoids were estimated for Spirulina platensis.

**Keywords:** water body, physico-chemical traits, spirulina, association analysis.

**GJSFR-H Classification :** FOR Code: 760299p



*Strictly as per the compliance and regulations of :*



# A Study on Eco-Physiology of *Spirulina* in Relation to some Environmental Parameters

Sunita Verma<sup>α</sup>, Divya Tiwari<sup>σ</sup> & Ajay Verma<sup>ρ</sup>

**Abstract-** Physico-chemical characteristics of Surajkund water body in Varanasi (UP) were monitored. High levels of nitrogen (N), phosphorus (P), calcium (Ca), potassium (K) and sodium (Na) indicated that pond was eutrophic. However, contrast to nitrate, ammonia-nitrogen was high in summer and low in winter. The high concentrations of sodium in summer stimulated uptake and transport of CO<sub>2</sub> and HCO<sub>3</sub> at high pH. Highly significant correlation was observed among studied traits. NO<sub>3</sub> exhibited significant negative relation with all traits, though the magnitude varied from NO<sub>2</sub> to NH<sub>4</sub>. More over positive correlation observed among traits only exception was NO<sub>3</sub>. Calcium, Potassium, Sodium etc showed significant positive correlation with other traits except NO<sub>3</sub>. Positive correlation maintained by NO<sub>2</sub> with other traits where negative value with NO<sub>3</sub>. Surajkund *Microcystisaeruginosa* was dominant plankton in the month of December January and February. *Spirulina* sp. was found along with the population of *Oscillatoria* and *Chlamydomonas* in April, May and June. After 15 days of growth, protein, dry weight, carbohydrate, chlorophyll and carotenoids were estimated for *Spirulina platensis*.

**Keywords:** water body, physico-chemical traits, *spirulina*, association analysis.

## I. INTRODUCTION

The blue green alga *Spirulina* has attracted worldwide interest as photosynthetic planktonic organism suitable for mass cultivation<sup>[1,3,6]</sup>. Efforts are underway to evolve a simple rural technology for producing *Spirulina* biomass and to utilize it as a feed supplement for cattle and poultry. *Spirulina* has a higher protein content, higher growth rate and easy digestibility<sup>[14,9,16]</sup>. Malnutrition in developing countries has catalyzed several efforts to intensify the production of protein both from conventional agriculture and from unconventional sources. *Spirulina* because of its many favourable properties has considerable future potential in improving the protein supply to mankind<sup>[7,10,12]</sup>. In India millions of hectares of available land annually rendered unfit for farming because of increasing salinity or alkalinity. The saline alkaline lands occupy an area of about 65 lakh hectares in the country. These lands are characterized by impermeability, silty texture and loose aggregation of soil particles. These saline-alkaline lands are locally known as Usar Land. Due to canal irrigation, the area of user land increasing year after year. User lands show high pH from 8.3 – 10 or even 11. These

lands also contain high levels of sodium. Due to high alkalinity and salinity user land unfit for growth for most of the crops. The problem of soil alkalinity and salinity is due to the salt formation of Na<sup>+</sup>, K<sup>+</sup>, Ca<sup>++</sup> and Mg<sup>++</sup> ions with Cl<sup>-</sup> and SO<sub>4</sub> mainly, sometimes with NO<sub>3</sub> and CO<sub>3</sub> and to a small extent with HCO<sub>3</sub> which may be nutrients for *Spirulina*. *Spirulina* can grow at high alkalinity up to pH 8 to 11 and prefers high salt content for growth<sup>(14,15,18,19,20)</sup>

## II. MATERIALS AND METHODS

### a) Collection, isolation and purification

Water and soil samples were collected from certain alkaline sites and were analysed as per standard procedure of APHA (1989).

#### i. Growth Measurement

The growth experiments for isolated strains were conducted in 100 ml flasks. Exponential grown cultures were centrifuged washed with sterile double distilled water and re-centrifuged. The inoculum of filamentous strains was prepared by cyclo-mixing. The following methods were used for growth estimation.

#### a. Protein estimation

Protein content of algal samples was determined by following the method of Lowery et al as modified by Herbert et al.

#### b. Pigment extraction

Estimation of Chlorophyll a- Chlorophyll a and carotenoids were extracted in methanol and estimated as per the methodology of Mackinney. Chlorophyll a and carotenoids were recorded for absorbance of 663 and 480 nm.

#### c. Carbohydrate estimation

Total sugar was estimated by Phenol-sulfuric acid method using the absorbance of 480 nm, using glucose as standard.

#### d. Dry weight

For the measurement of dry weight, algal cultures of known volume were filtered on Whatman Nol filters and dried in a hot air vacuum oven at 45 C for 24 hours. Ammonia was determined as per the method of Solarzano. Dissolved Oxygen in the water sample was estimated by Winkler's method. Nitrate in water sample was estimated by Brucinesulphuric acid method.

### III. RESULTS AND DISCUSSION

Different water bodies in Varanasi such as *Durgakund*, *Surajkund*, *Laxmikund*, and *Laatbhairo* were extensively surveyed for high alkalinity (Table-3). User soil and water samples (Table-2, Figure-1) [Varanasi] were also collected for isolation of *Cyanobacteria*. All strains were made axenic using standard microbiological techniques and identified with the help of *Desikachary* (1959). All these strains were maintained in air conditioned culture room at 26±3 in presence of 75μ E light m<sup>-2</sup> Sec<sup>-1</sup> with 18 hr and 6hr light dark periods. Cultures were transferred to fresh agar slants at an intervals of 16 days. Following strains were isolated from saline-alkaline habitats and their pH limits were recorded (Table 2). In Suraj Kund, a dense surface bloom of *Microcystisaeruginosa* was observed during the month of December, January and February. In the months of April, May and June *Oscillatoria amphibia* dominated in Suraj Kund. *Spirulinasp* and *Chlamydomonassp* were found along with the population of *Oscillatoria* in the months of May, June and July. The physico-chemical characteristics of the pond water recorded in different months are given in Table 3. The pH values varied hourly and monthly also, pH was lowest in December and January and highest during May and June. Figure indicate that pH of water varied greatly from morning to evening; it was highest at 2:30 PM, followed by 6:30 PM, 10:30 PM and 6:30AM. The temperature varied greatly from 19 to 42 C. Similar to pH, temperature was also highest at 2:30 PM followed by evening and morning on

the same day. It was highest in May and June and lowest in December and January. Marked increase in dissolved oxygen content of water occurred during day time at 2:30 PM in all the months and followed the same trend like pH and temperature. Hardness, alkalinity and total alkalinity increases considerably from December to June being highest in May, June and lowest in December. The ammoniacal nitrogen was highest in May, June and lowest in December, January. In contrast to this, nitrate nitrogen was minimum in May, June and maximum in December and January. Sodium, Potassium and calcium concentrations in water samples followed similar trends, being highest in May, June and lowest in December and January. After 15 days growth, protein, dry weight, carbohydrate, chlorophyll and carotenoids were 400, 780(mg/l), 125, 6.5 and 4.92 (μg/ml) respectively for *Spirulina platensis*. The specific growth rate and generation times were 0.026 and 38.7 (h) (Table-1). Correlation analysis measures the closeness of the linear relationship between chosen variables (Table-4). Moreover the value of correlation coefficient nearer to +1 or -1, shows the probability of linear relationship between the variables x and y. Highly Significant correlation observed among studied traits. NO<sub>3</sub> exhibited significant negative relation with all traits, though the magnitude varies from NO<sub>2</sub> to NH<sub>4</sub>. More over positive correlation observed among traits only exception is NO<sub>3</sub>. Calcium, Potassium, sodium etc showed significant positive correlation with other traits except NO<sub>3</sub>. Positive correlation maintained by NO<sub>2</sub> with other traits where negative value with NO<sub>3</sub> (Table-4)

**Table 1 :** Growth, biomass, chemical composition of *S.platensis* attained at 15 days of growth

Parameters	<i>S platensis</i>
Final growth 0.0 (665m)	1.52
Specific growth rate (generation h <sub>-1</sub> )	0.026
Generation time (h)	38.70
Dry weight (mg/l)	780.00
Carbohydrate (μg/ml)	125.00
Protein (μg/ml)	400.00
Chlorophyll (μg/ml)	6.50
Cartenoids (μg/ml)	4.92

**Table 2 :** Collection site, characteristics of Cynobacteria isolated from different water bodies



Cyanobacterial	Medium	Collection site	Characteristics	pH for growth
<i>Microcystisaeruginosa</i>	Jaworski medium	Suraj Kund, Laxmi Kund, Laat Bhairo	Unicellular, bloom forming	9-11
<i>Oscillatoriaamphibia</i>	CHU-10	Suraj Kund	Non-heterocystous filamentous	8-9.5
<i>Nostoccalcicola</i>	Allen Arnon	Usar Soil, Bhadooi, Varanasi	He terocystous filamentous	8
<i>Nostoc</i> sp.	Allen Arnon	-do-	-do-	8
<i>Anabaena</i> sp.	Allen Arnon	-do-	-do-	8.2
<i>Spirulina</i> sp.	Zarrouk's medium	Suraj Kund, Laat Bhairo	Non-heterocystous filamentous	9

Table 3 : Physico-chemical characteristics of Suraj Kund water in different months

	December	January	February	March	April	May	June
Hardness (mg/l)	18.6	22.4	21	23.8	24.6	26	28
Alkalinity(mg/l)	20	21	21	38	39	42	42
Total Alkalinity(mg/l)	169	165	164	189	190	199	225
NO <sub>3</sub> (mg/l)	9.4	9.3	9.2	8.4	7.8	5.25	3.75
NH <sub>4</sub> (mg/l)	0.14	0.21	0.4	0.9	1.7	3.92	4.4
NO <sub>2</sub> (mg/l)	0.46	1.6	2.9	3	3.2	3.6	3.7
Na (ppm)	110	117	119	130	135	139	140
K(ppm)	78	89	90	110	124	125	130
Ca(ppm)	23	23.4	24.3	26	28.5	29	29.2

Table 4 : Correlation matrix for physio-chemical traits of Suraj Kund

	Hardness (mg/l)	Alkalinity (mg/l)	Total Alkalinity (mg/l)	NO <sub>3</sub> (mg/l)	NH <sub>4</sub> (mg/l)	NO <sub>2</sub> (mg/l)	Na (ppm)	K (ppm)	Ca (ppm)
Hardness (mg/l)	1.0000	0.8921	0.8993	-	0.8891	0.8552	0.9531	0.9481	0.9116
Alkalinity(mg/l)		1.0000	0.8867	0.8920	0.8264	0.8133	0.9680	0.9682	0.9503
Total Alkalinity(mg/l)			1.0000	0.8054	0.9200	0.7037	0.8799	0.8881	0.8845
NO <sub>3</sub> (mg/l)				1.0000	-	-	-	-	-0.8695
NH <sub>4</sub> (mg/l)					1.0000	0.7102	0.8528	0.8452	0.8999
NO <sub>2</sub> (mg/l)						1.0000	0.8763	0.8665	0.8590
Na (ppm)							1.0000	0.9943	0.9801
K(ppm)								1.0000	0.9866
Ca(ppm)									1.0000






Figure 1 : Monthly variations in pH, temperature and DO values

#### IV. ACKNOWLEDGEMENT

We acknowledged Dr. Manoj Kr. Saika, Associate Professor & Coordinator *Institutional Biotech Hub Dhing College, Nagaon(Assam)* and Miss. Farhana Begum JRF, IBTHub, Dhing College Nagaon (Assam) for providing necessary guidance in carrying out this research work.

#### REFERENCES RÉFÉRENCES REFERENCIAS

1. Fox, R.D. 1985. Spirulina, the alga that can end malnutrition. *The Futurist*, 19: 30-35.
2. Gabby, A.R., Tel, O.E. and Gresshoff, P.M. 1993, Mechanisms of salt tolerance in cyanobacteria. Plant Sources to the Environment. *Current Topics in Plant Molecular Biology* pp. 123-132.
3. Li, D.M. 1995. Spirulina as a health food. In *Spirulina*, pp, 2128, Beijing, China, Chinese Agro technology Publication.
4. Shimamatsu, H. 2004. Mass production of Spirulina, an edible microalga, *Hydrobiol.*, 512: 39-44.
5. IECFA (FAO/WHO Expert Committee on Food Additives). 2002. Evaluation of certain food additives. Fifty ninth report of the joint FAO/WHO Expert Committee on Food Additives. Genova, World Health Organization.
6. Muthukumar, C Muralitharan, G., Vijayakumar, R., Panneerselvam, A. and Thajuddin, N. 2007. Cyanobacterial biodiversity from different freshwater ponds of Thanjavur, Tamilnadu India *Acta Botanica Malacitana*, 32: 1-9.
7. Anuradha, V. and Vidhya, D. 2001. Impact of administration of Spirulina on the blood glucose levels of selected diabetic patients *Indian J. Nutri. & Dietetics*, 38: 40-44.
8. Ayachi, S., El-Abed, A., Medhioub, A., Brouers, M. and Marzouk, B, 2004. Influence of temperature and light intensity on fatty acid composition during the development of spirulina, *Arthrospira platensis*, *Rivista Italiana delle Sostanze Grasse*, 81: 185-190.
9. Belay, A., Yoshimichi, O., Miyakawa, K. and Shimamatsu, H. 1993. Current knowledge on potential health benefits of Spirulina. *J. Appl. Phycol.*, 5: 235-241.
10. Bhattacharya, S. and Shivaprakash, M.K. 2005. Evaluation of three Spirulina species grown under similar conditions for their growth and biochemicals. *J. Sci. Food Agric*, 85: 333-336.
11. Costa, J.A.V, Colla, L.M. and Filho, P.F.D. 2004. Improving Spirulina platensis biomass yield using a fed-batch process. *Bioresource Technol.*, 92: 237-241.
12. Laliberte, G., Olguin, E.J. and de la Noue, J. 1997. Mass cultivation and wastewater treatment using Spirulina. in A. Vonshak, ed. *Spirulina platensis (Arthrospira platensis) Physiology, Cell Biology and Biotechnology*, pp, 159-174. Basingstoke, Hants, London, Taylor and Francis.
13. Pareek, A. and Srivastava, P. 2001. Optimum photoperiod for the growth of Spirulina platensis, *J. Phytol. Res.*, 14: 219-220.
14. Vonshak, A, (ed). 1997. *Spirulina platensis (Arthrospira)*. In *Physiology, Cell Biology and Biotechnology*. Basingstoke, Hants, London, UK, Taylor and Francis.
15. Abdulqader G, Barsanti L and Tredici MR 2000. Harvest of Arthrospira platensis from Lake Kossorom (Chad) and its household usage among the Kanembu. *Journal of Applied Phycology*, 12: 493-498.
16. Anaga A and Abu GO 1996. A laboratory-scale cultivation of Chlorella and spirulina using waste effluent from fertilizer Company in Nigeria. *Biores Technol*, 58: 93-95.
17. Banerjee M and Deb M. 1996. Potential of fly ash and spirulina combination as a slow release fertilizer for rice yield. *Cientifica jaboticabal*, 24: 55-62.
18. Ciferri O. 1983. Spirulina, the edible organism. *Microbiol rev.* 47: 551-578.
19. Lodi A, Binagli L, Solisio C, Converti A and Del Borghi M. 2003. Nitrate and phosphate removal by Spirulina plantensis. *J of Ind Microb and Biotech* 30: 656-666.

19. Zeng WL, Cai ZL and Ouyang F. 2001. Growth characteristics and dynamic model of *Spirulina platensis*. Chinese J Appl Environ Biol, 7:360-364.



This page is intentionally left blank