



GLOBAL JOURNAL OF SCIENCE FRONTIER RESEARCH: B
CHEMISTRY

Volume 15 Issue 6 Version 1.0 Year 2015

Type : Double Blind Peer Reviewed International Research Journal

Publisher: Global Journals Inc. (USA)

Online ISSN: 2249-4626 & Print ISSN: 0975-5896

1-Octanol/Water Partition Coefficients of Dialkylated Methylimidazolium Halide Salts

By Marissa K. Gard, Elliot G. Ennis & John H. Summerfield

Missouri Southern State University, United States

Abstract- Dialkylated methylimidazolium halide salts are a current research topic. The 1-octanol/water partition coefficient for these salts has come to signify the tendency of a charged organic molecule to cross a biological membrane, be sequestered in soil, and indicate the lethal dose of the chemical. The purpose of this work is to add to existing data for the 1-octanol/water partition coefficient for these salts. The novelty of this work is that dialkylated methylimidazolium salts are investigated.

Keywords: *ionic liquids, partition functions.*

GJSFR-B Classification : *FOR Code: 860602*



1OCTANOLWATERPARTITIONCOEFFICIENTSOFDIALKYLATEDMETHYLIMIDAZOLIUMHALIDESALTS

Strictly as per the compliance and regulations of :



RESEARCH | DIVERSITY | ETHICS

1-Octanol/Water Partition Coefficients of Dialkylated Methylimidazolium Halide Salts

Marissa K. Gard ^α, Elliot G. Ennis ^σ & John H. Summerfield ^ρ

Abstract- Dialkylated methylimidazolium halide salts are a current research topic. The 1-octanol/water partition coefficient for these salts has come to signify the tendency of a charged organic molecule to cross a biological membrane, be sequestered in soil, and indicate the lethal dose of the chemical. The purpose of this work is to add to existing data for the 1-octanol/water partition coefficient for these salts. The novelty of this work is that dialkylated methylimidazolium salts are investigated.

Keywords: ionic liquids, partition functions.

I. INTRODUCTION

Green chemistry is founded on guidelines that are designed to prevent and reduce the waste associated with the production of chemicals. Any modifications to existing processes as well as the creation of new processes that lead to more efficient and recyclable solvents, safer reactions, and renewable feedstocks are successes.[1]

Ionic liquids or molten salts are molecules composed wholly of ions and occur in the liquid state. An inorganic salts such as NaCl is an ionic liquid above its melting point. Due to their high melting point many inorganic salts are of little use in daily life. Instead, chemists have turned to imidazolium-based ionic liquids. These organic salts are current candidates for

green solvents. They can be highly soluble, can have high ionic conductivity, are generally inflammable, have high thermal and chemical stability, and relatively low vapor pressures compared to cyclohexane, benzene, and acetone. Typically the imidazolium salts under investigation are alkylmethylimidazolium compounds. The novelty of this work is that dialkylated methylimidazolium salts are investigated.[2]


Trisubstituted imidazolium salts are a current research topic. They have been investigated as a solvent, an electrolyte solution additive, a reaction initiator, a product precursor, a molecular aggregator, and a composite material.[3]

The 1-octanol/water partition coefficient (K_{ow}) for these salts has come to signify the tendency of a charged organic molecule to cross a biological membrane,[4] and as an estimate to the salt's environmental impact before widespread use. The purpose of this work is to add to existing data for the 1-octanol/water partition coefficient for these salts.[5, 6]

II. MATERIALS AND METHODS

a) Synthesis

The synthesis is shown below (Scheme 1). It is outlined in the literature.[7]



Scheme 1 : Two R groups are added to the 3-methyl-1H-imidazol-3-ium ion

The chemicals used in the syntheses, the CAS numbers, sources and grades are below. The proton NMR was performed on a Bruker AV-III 300 MHz spectrometer. The solvent was DMSO and the standard was $\text{Si}(\text{CH}_3)_4$. The peak assignments are below.

1-methyl-3-octyl-1H-imidazol-3-ium bromide

3-methyl-1H-imidazol-3-ium (1-methylimidazole 616-47-7, Aldrich, 99%, 25.00 mL), 1-bromooctane (111-

83-1, Aldrich, 99%, 71.00 mL), and toluene (108-88-3, Aldrich, 99.8%, 40.00 mL) were mixed. The solution was refluxed for two days. The resulting mixture was frozen and the excess toluene was decanted. The product was thawed and washed with ethyl acetate (141-78-6, Aldrich, 99.8%). The solution was evaporated to dryness. The product was frozen and the excess reagents were decanted. δH (300 MHz; DMSO; $\text{Si}(\text{CH}_3)_4$) 8.90 (1 H, s, 2-H), 7.91 (1 H, d, 5-H), 7.78 (1 H, d, 4-H), 5.00 (2 H, t NH), 3.75 (3 H, s, NCH_3), 2.00 (2 H, qt, CH_2), 1.30 (10 H, s, CH_2) 0.91 (3 H, br, CH_3) The x-ray photoelectric spectroscopy, FTIR, and ^1H NMR markers are also in the literature.[8, 9].

Author α ρ : Department of Chemistry and Physical Sciences, Missouri Southern State University, 3950 Newman Rd. Joplin. e-mail: summerfield-j@mssu.edu.

1,2-dimethyl-3-octyl-1H-imidazol-3-ium iodide

To a stirred solution of 1-methyl-3-octyl-1H-imidazol-3-ium bromide (10.00 g) in acetonitrile (75-05-8, Aldrich, 99.8%, 175.00 mL) was added NaH (7646-69-7, Aldrich, 60% in mineral oil, 2.00 g). After allowing the mixture to stir four hours, iodomethane (74-88-4, Aldrich, 99%, 6.90 mL) was added and the reaction was stirred for 12 hours. The solution was filtered to remove any precipitated NaI. The resulting oil as washed with ethyl acetate (141-78-6, Aldrich, 99.8%, 3x75.00 mL) and the residual volatiles were then removed under vacuum. The residual nonvolatiles were removed by freezing and decantation. δ H(300 MHz; DMSO; Si(CH₃)₄) 7.91 (1 H, d, 5-H), 7.78 (1 H, d, 4-H), 5.00 (2 H, t NH), 3.75 (3 H, s, NCH₃), 2.85 (3 H, s, 2-CH₃) 2.00 (2 H, qt, CH₂), 1.30 (10 H, s, CH₂) 0.91 (3 H, br, CH₃) The ¹H NMR markers are in the literature.[10, 11]

Workers find the IUPAC nomenclature wearisome. This paper will represent the 1-methylimidazolium fragment as mim. The molecule 1-methyl-3-octyl-1H-imidazol-3-ium bromide is shortened to [omim]Br. Similarly, 1,2-dimethyl-3-octyl-1H-imidazol-3-ium iodide becomes [momim]I. The two letters hx will refer to the hexyl group.

[bomim]Cl

2-butyl-1-methyl-3-octyl-1H-imidazol-3-ium chloride was prepared in the same manner as [momim]I with the exception that 1-chlorobutane (109-69-3, Aldrich, 99.5%, 11.6 mL) was used instead of iodomethane. δ H(300 MHz; DMSO; Si(CH₃)₄) 7.91 (1 H, d, 5-H), 7.78 (1 H, d, 4-H), 5.00 (2 H, t NH), 3.75 (3 H, s, NCH₃), 2.50 (2 H, t, CH₂), 2.00 (2 H, qt, CH₂), 1.61 (2 H, qt, CH₂) 1.30 (12 H, s, CH₂) 0.91 (6 H, br, CH₃)

[pomim]Cl

1-methyl-3-octyl-2-pentyl-1H-imidazol-3-ium chloride was prepared in the same manner as [momim]I with the exception that 1-chloropentane (543-59-5, Aldrich, 99%, 13.4 mL) was used instead of iodomethane. δ H(300 MHz; DMSO; Si(CH₃)₄) 7.91 (1 H, d, 5-H), 7.78 (1 H, d, 4-H), 5.00 (2 H, t NH), 3.75 (3 H, s, NCH₃), 2.50 (2 H, t, CH₂), 2.00 (2 H, qt, CH₂), 1.61 (2 H, qt, CH₂) 1.30 (14 H, s, CH₂) 0.91 (6 H, br, CH₃)

[oomim]Cl

1-methyl-2, 3-dioctyl-1H-imidazol-3-ium chloride was prepared in the same manner as [momim]I with the exception that chlorooctane (111-85-3, Aldrich, 99%, 18.9 mL) was used instead of iodomethane. δ H(300 MHz; DMSO; Si(CH₃)₄) 7.91 (1 H, d, 5-H), 7.78 (1 H, d, 4-H), 5.00 (2 H, t NH), 3.75 (3 H, s, NCH₃), 2.50 (2 H, t, CH₂), 2.00 (2 H, qt, CH₂), 1.61 (2 H, qt, CH₂) 1.30 (20 H, s, CH₂) 0.91 (6 H, br, CH₃)

[domim]Cl

2-decyl-3-methyl-2-octyl-1H-imidazol-3-ium chloride was prepared in the same manner as [momim]I with the

with the exception that chlorodecane (1002-69-3, Aldrich, 98%, 22.6 mL) was used instead of iodomethane. δ H(300 MHz; DMSO; Si(CH₃)₄) 7.91 (1 H, d, 5-H), 7.78 (1 H, d, 4-H), 5.00 (2 H, t NH), 3.75 (3 H, s, NCH₃), 2.50 (2 H, t, CH₂), 2.00 (2 H, qt, CH₂), 1.61 (2 H, qt, CH₂) 1.30 (22 H, s, CH₂) 0.91 (6 H, br, CH₃)

[bbmim]Cl

3-methyl-1H-imidazol-3-ium (1-methylimidazole 616-47-7, Aldrich, 99%, 25.00 mL), 1-chlorobutane (109-69-3 Aldrich, 99.5%, 43.00 mL), and toluene (108-88-3, Aldrich, 99.8%, 40.00 mL) were mixed. The solution was refluxed for two days. The resulting mixture was frozen and the excess toluene was decanted. The product was thawed and washed with ethyl acetate (141-78-6, Aldrich, 99.8%). The solution was evaporated to dryness. The product was frozen and the excess reagents were decanted. To a stirred solution of 1-methyl-3-octyl-1H-imidazol-3-ium bromide (10.00 g) in acetonitrile (75-05-8, Aldrich, 99.8%, 150.00 mL) was added NaH (7646-69-7, Aldrich, 60% in mineral oil, 2.67 g) After allowing the mixture to stir four hours, 1-chlorobutane (109-69-3 Aldrich, 99.5%, 17.8 mL) was added and the reaction was stirred for 12 hours. The solution was filtered to remove any precipitated NaCl. The resulting oil as washed with ethyl acetate (141-78-6, Aldrich, 99.8%, 3x75.00 mL) and the residual volatiles were then removed under vacuum. The residual nonvolatiles were removed by freezing and decantation. δ H(300 MHz; DMSO; Si(CH₃)₄) 7.91 (1 H, d, 5-H), 7.78 (1 H, d, 4-H), 5.00 (2 H, t NH), 3.75 (3 H, s, NCH₃), 2.50 (2 H, t, CH₂), 2.00 (2 H, qt, CH₂), 1.61 (2 H, qt, CH₂) 1.30 (4 H, sx, CH₂) 0.91 (6 H, t, CH₃)

[pbmim]Cl

2-butyl-1-methyl-3-pentyl-1H-imidazol-3-ium chloride was prepared in the same manner as [bbmim]Cl with the exception that 1-chloropentane (543-59-9, Aldrich, 99%, 20.7 mL) was used instead of 1-chlorobutane. δ H(300 MHz; DMSO; Si(CH₃)₄) 7.91 (1 H, d, 5-H), 7.78 (1 H, d, 4-H), 5.00 (2 H, t NH), 3.75 (3 H, s, NCH₃), 2.50 (2 H, t, CH₂), 2.00 (2 H, qt, CH₂), 1.61 (2 H, qt, CH₂) 1.30 (6 H, sx, CH₂) 0.91 (6 H, t, CH₃)

[hxbmim]Cl

2-butyl-3-hexyl-1-methyl-1H-imidazol-3-ium chloride was prepared in the same manner as [bbmim]Cl with the exception that 1-chlorohexane (544-10-5, Aldrich, 99%, 23.5 mL) was used instead of 1-chlorobutane. δ H(300 MHz; DMSO; Si(CH₃)₄) 7.91 (1 H, d, 5-H), 7.78 (1 H, d, 4-H), 5.00 (2 H, t NH), 3.75 (3 H, s, NCH₃), 2.50 (2 H, t, CH₂), 2.00 (2 H, qt, CH₂), 1.61 (2 H, qt, CH₂) 1.30 (8 H, br, CH₂) 0.91 (6 H, t, CH₃)

[dbmim]Cl

2-butyl-3-decyl-1-methyl-1H-imidazol-3-ium chloride was prepared in the same manner as [bbmim]Cl with the exception that 1-chlorodecane (1002-69-3,

Aldrich, 98%, 34.8 mL) was used instead of 1-chlorobutane. δ H(300 MHz; DMSO; Si(CH₃)₄) 7.91 (1 H, d, 5-H), 7.78 (1 H, d, 4-H), 5.00 (2 H, t NH), 3.75 (3 H, s, NCH₃), 2.50 (2 H, t, CH₂), 2.00 (2 H, qt, CH₂), 1.61 (2 H, qt, CH₂) 1.30 (16 H, br, CH₂) 0.91 (6 H, t, CH₃)

The UV/Vis spectra were produced on a Beckman DU-640B UV/Vis spectrophotometer. The salt, its phase, the λ_{\max} , and the corresponding absorbance are shown in the Results and Discussion section. The maximum absorbance is in the neighborhood of 265.0 nm, which indicates the imidazolium ring.[12]

The compound [oomim]Cl was excluded. The spectrum was simply noise. This may indicate a very similar blank and sample. The difference between the two---the UV/Vis spectrum---is noise.[13]

The compound [momim]I shows negative absorbance. Negative absorbance has no physical meaning except that the blank absorbs more light than the [momim]I.

This work finds the 1-octanol/water partition coefficient by the slow-stirring method.[14–18] The salt was added to a solution of 1-octanol saturated with water. Each solution was stirred for 45 days at 21°C. This was done slowly to prevent emulsification.

For this work initially 5.0 μ g of the organic salt was added to 5.0 mL of a saturated 1-octanol/water solution. When an emulsion formed, the solution was stirred for six hours and left to rest overnight. This removed the emulsion. Serial dilutions were done to keep the absorption values between –0.150 and 1.000. The aqueous layer was drawn off and compared to a saturated solution of 1-octanol. One extraction was done for each compound. The K_{OW} value was then calculated using

$$K_{OW} = \frac{(A_i \square df_i - A_f \square df_f) \square V_{\text{water}}}{A_f \square df_f \square V_{\text{octanol}}} \quad (1)$$

where A_i and A_f are the absorbances of the aqueous layer before and after extraction. V_{water} and V_{octanol} are the volumes of the water and 1-octanol phases. The symbols df_i and df_f are the dilution factors used to find A_i and A_f .[19,20]

III. RESULTS AND DISCUSSION

The UV/Vis data, the experimental log K_{OW} values, the calculated log K_{OW} values, and the absolute difference between them are shown below in Table 1.

Compound in water phase	λ_{\max} in nm	Absorbance at λ_{\max}	Exp. Log K_{OW}	Calc. Log K_{OW}	$ \Delta \text{Log } K_{OW} $
[momim]I	248.0	-0.1459	-0.62	-0.8850	0.27
[bomim]Cl	269.0	0.4209	1.1	0.5020	0.6
[pomim]Cl	265.0	0.21450	0.98	1.031	0.051
[domim]Cl	263.0	0.1249	5.1	3.676	1.4
[bbmim]Cl	264.0	0.6850	-1.3	-1.614	0.31
[pbmim]Cl	264.0	0.6413	-0.93	-1.085	0.16
[hxbmim]Cl	259.0	0.1866	-0.31	-0.5560	0.25
[dbmim]Cl	232.0	0.2346	1.9	1.560	0.3

Table 1: UV/Vis data, experimental log K_{OW} values, calculated log K_{OW} values, and the absolute difference between them. The calculated values were determined by the group contribution method rather than the linear free energy relation scheme.[5, 21]

Using the cLog P program[22] the calculated log K_{OW} values in Table 1 were found. The expectation is 85% of the calculated log K_{OW} values will be in error by less than 0.5, 10% by 0.5 to 1.0, and 5% by more than 1.0.[23] For this work, the values are 75%, 13%, and 13% respectively. The clog P program is best suited for molecules with typical fragments.[24] The charged five-membered ring and the long hydrocarbon chains may be misrepresented in this group additive method or the small sample size may explain the discrepancies.

The toxicity of organic solvents in animal tissue, in soil, and in water is an ongoing concern. In this work

we examine three toxicity indicators that are linked to the 1-octanol/ water partition coefficients. Positive value log K_{OW} values indicate the octanol phase is favored. These compounds are candidates for tissue storage. That is, candidates for the calculated of a bioconcentration factor (BCF). The soil sorption coefficient (K_{OC}) is the ratio of the mass of a chemical absorbed per weight of carbon in a soil to the concentration of the chemical in water. The lethal concentration of a chemical is defined as the lethal dose for half the population (LC_{50}).[25]

Two BCF models are shown in Figure 1. The lower dashed line is $\log(\text{BCF}) = 0.85 \log(K_{OW}) - 0.70$. The upper dashed line is $\log(\text{BCF}) = 0.79 \log(K_{OW}) - 0.40$. [25] The $\log(\text{BCF})$ for benzene from these models is 1.3 and 1.1, respectively.[2]




Figure 1 : Log of the tissue bio concentration for the positive $\log(K_{OW})$ salts. The lower dashed line is $\log(BCF) = 0.85 \log(K_{OW}) - 0.70$. The upper dashed line is $\log(BCF) = 0.79 \log(K_{OW}) - 0.40$.

Three K_{OC} models are shown below in Figure 2. The lower solid line is $\log(K_{OC}) = 0.989 \log(K_{OW}) - 0.364$. The middle line is $\log(K_{OC}) = 679 \log(K_{OW}) + 0.663$. The

top dashed line is $\log(K_{OC}) = 0.544 \log(K_{OW}) + 1.377$. [25] The $\log(K_{OC})$ for benzene for these three models are 1.7, 2.1, and 2.5, respectively. [2]




Figure 2 : Log of soil sorption coefficients for all the salts. The lower solid line is $\log(K_{OC}) = 0.989 \log(K_{OW}) - 0.364$. The middle line is $\log(K_{OC}) = 679 \log(K_{OW}) + 0.663$. The top dashed line is $\log(K_{OC}) = 0.544 \log(K_{OW}) + 1.377$

Figure 3 shows the log of the inverse lethal concentrations ($1/LC_{50}$) for each salt. The top line is $\log(1/LC_{50}) = 0.629 \log(K_{OW}) - 0.489$. The middle dotted line is $\log(1/LC_{50}) = 0.854 \log(K_{OW}) - 1.74$. The solid line is $\log(1/LC_{50}) = 0.89 \log(K_{OW}) - 1.75$. [25] The $\log(1/LC_{50})$ for benzene from these models are 0.85, 0.079, 0.15, respectively. [2]




Figure 3 : The log of the inverse lethal concentrations ($1/LC_{50}$) for each salt. The top line is $\log(1/LC_{50}) = 0.629 \log(K_{ow}) - 0.489$. The middle dotted line is $\log(1/LC_{50}) = 0.854 \log(K_{ow}) - 1.74$. The solid line is $\log(1/LC_{50}) = 0.89 \log(K_{ow}) - 1.75$

IV. CONCLUSIONS

Green chemistry has taken a winding road in its search for benign solvents. Some, such as CO_2 , have become firmly established.[26] Others, such as imidazolium salts, continue to be under investigation.[27–32] A physical quantity that continues to be of interest is the 1-octanol/water partition coefficients. One reason is their direct link to the environmental impact of the salts.[33, 34]

Dialkylated methylimidazolium salts have low vapor pressure compared to typical solvents such as benzene. This work shows they can be synthesized to be water soluble or lipophilic. This work also shows that every imidazolium salt is not “green”. The salts with octyl and/or decyl substituents have comparable toxicity to benzene. It has recently been shown that cholinium alkanoates may provide a set of molecules that are more ecologically friendly.[35]

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

REFERENCES RÉFÉRENCES REFERENCIAS

1. United States Environmental Protection Agency. <http://epa.gov/>
2. L. Ropel, L.S. Belvèze, S. Aki et al., “1-octanol-water Partition Coefficients of Imidazolium-based Ionic Liquids”, *Green Chem.*, vol. 2005, no. 2, pp. 83-90, 2005.
3. F. Hernat and H. Yu, *J. Liq. Chrom. and Rel. Tech.*, “Ion Chromatography of Tri-Substituted Imidazolium Ionic Liquid Cations: Method and Application for Octanol–Water Partition Coefficient (K_{ow}) Determination”, vol.38, no. 13, pp. 1267-1272, 2015.
4. V. Levin, “Relationship of 1-octanol/ Water Partition Coefficient and Molecular Weight to Rat Brain Capillary Permeability,” *J. Med. Chem.*, vol. 23, no. 6, pp.692-684, 1980.
5. C. Cho, C. Jungnickel, J. Arning et al., “Ionic Liquids: Predictions of Physicochemical Properties with Experimental and/or DFT-calculated LFER Parameters to Understand Molecular Interactions in Solution”, *J. Phys. Chem. B* vol. 115, no. 9, pp.6040-6050, 2011.
6. U. Domańska, M. Królikowski, A. Pobudkowska, et al., “Solubility of Ionic Liquids in Water and Octan-1-ol and Octan-1-ol/water, or 2-phenylethanol/water Partition Coefficients”, *J. Chem. Thermo.*, vol. 55, pp. 225-233, 2012.
7. E. Ennis and S. Handy, “A Facile Route to C2-Substituted Imidazolium Ionic Liquids”, *Molecules*, vol. 14, no. 6, pp. 2235-2245, 2009.
8. T. Cremer, C. Kolbeck, K. Lovelock et al., “Towards a Molecular Understanding of Cation–Anion Interactions—Probing the Electronic Structure of Imidazolium Ionic Liquids by NMR Spectroscopy, X-ray Photoelectron Spectroscopy and Theoretical Calculations”, *Chem.: A Eur J.* vol. 16, no. 30, pp. 9018-9033, 2010.
9. W. Awad, J. Gilman, M. Nyden et al., “Thermal degradation studies of alkyl-imidazolium salts and their application in nanocomposites”, *Thermo. Acta*, vol. 409, pp. 3-11, 2004.
10. S. Handy and M. Okello, “The 2-Position on Imidazolium Ionic Liquids: Substitution and Exchange” *JOC*, vol. 70, no. 5, pp. 1915-1918, 2005.

11. T. Endo and K. Nishikawa, "Effects of Methylation at Position 2 of Cation Ring on Rotational Dynamics of Imidazolium-Based Ionic Liquids Investigated by NMR Spectroscopy: [C(4)mim]Br vs [C(4)C(1)mim]Br", *J. Phys. Chem. B*, vol. 114, no. 28, pp. 9201-9208, 2010.
12. Q. Lin, F. Huang, M. Liu et al. "Imidazolium-salt-based Fluorescent Chemosensor for Cu(II), Zn(II), and Co(II)", *Sensors and Materials*, vol. 20, no. 3, pp.143-150, 2008.
13. P. Sassi, M. Paolantoni, R. Cataliotti, et al., "Water/Alcohol Mixtures: A Spectroscopic Study of the Water-saturated 1-octanol Solution", *J. Phys. Chem. B* vol. 108, no. 50, pp.19557-19565, 2004.
14. J. DeBruijn, F. Busser, W. Seinen et al., "Determination of 1-octanol/water Partition Coefficients for Hydrophobic Organic Chemicals with the Slow-stirring Method", *Envir. Chem.*, vol. 8, pp. 499-512, 2009.
15. G. Veith, N. Austin, R. Morris, "A Rapid Method for Estimating log P for Organic Chemicals", *Water Res.*, vol. 13, pp. 43-47, 1979.
16. R. Rapaport, R. Eisenrelch, "Chromatographic Determination of 1-octanol-water Partition Coefficients (K_{ow} 's) for 58 PCB Polychlorinated biphenyl congeners", *Environ. Sci. Technol.* Vol. 18, pp. 163-170, 1984.
17. US EPA, "Product Properties Test Guidelines: Partition Coefficients (n -1-octanol/water), Estimation by Liquid Chromatography", OPPTS 830.7570, EPA 712-C-96-040, 1996.
18. H. DeVoe, M. Miller, S. Wasik, Generator Columns and High Pressure Liquid Chromatography for Determining Aqueous Solubilities and 1-octanol-water Partition Coefficients of Hydrophobic Substances", *J. Res. NBS*, vol. 86, no.4, pp. 361-366, 1981.
19. M. Harris, J. Logan, "Determination of log K_{ow} Values for Four Drugs", *J. Chem. Ed.* vol. 91, 915-918, 2014.
20. http://chemwiki.ucdavis.edu/Analytical_Chemistry/Analytical_Chemistry_2.0/07%3A_Collecting_and_Preparing_Samples/7G%3A_Liquid%E2%80%93Liquid_Extractions.(accessed October, 2015).
21. T. Cheng, Y. Zhao, X. Li et al., "Computation of 1-octanol-water Partition Coefficients by Guiding an Additive Method with Knowledge", *J. Chem. Info. and Mod.*, vol. 47, no. 6, pp.2140-2148.
22. D. Evans and S. Rubenstein, *ChemDraw Professional (Revision 15)* Perkin-Elmer, Waltham, MA, 2015.
23. K van Leeuwen, *Technical Guidance Document on Risk Assessment*, European Communities, Rome, 2003.
24. Sussex Drug Discovery Centre, sussexdrugdiscovery.wordpress.com/2015/02/03/not-all-logps-are-calculated-equal-clogp-and-other-short-stories/, (accessed October 2015).
25. M. Montalbán, M. Collado-González, R. Trigo et al., "Experimental Measurements of Octanol-Water Partition Coefficients of Ionic Liquids", *J. Ad. Chem Eng.*, vol. 5, no. 3, pp. 1-8, 2015.
26. D. Hâcu, E. Beckman, "Generation of hydrogen peroxide directly from H₂ and O₂ Using CO₂ as the solvent", *"Green Chemistry"*, vol. 3, no. 2, pp.80-86, 2001.
27. C. Margulis, H. Stern, B. Berne, "Computer Simulation of a Green Chemistry Room-temperature Ionic Solvent", *J. Phys. Chem. B*, vol. 106, no. 46, pp.12017-12021, 2002.
28. A. Chapeaux, L. Simoni, M. Stadther, et al. "Liquid Phase Behavior of Ionic Liquids with Water and 1-octanol/water Partition Coefficients", *J. Chem. Eng. Data*, vol. 52, no. 6, pp.2462-2467, 2007
29. S. Katkevica, A. Zicmanis, P. Mekss, Imidazolium and Pyridinium Salts---Solvents Influencing the Rate and Direction of the Fries, Beckmann, and Claisen Rearrangements", *Chem. Hetero. Comp.*, vol. 46, no. 2, pp.195-208, 2010
30. A. Cláudio, M. Freire, C. Freire, et al., "Extraction of Vanillin using Ionic-liquid-based Aqueous Two-phase Systems", *Sep. and Pur. Tech.* vol. 75, pp.39-47, 2010.
31. M. Freire, I. Marrucho, J. Coutinho, "High-performance extraction of Alkaloids Using Aqueous Two-phase systems with Ionic Liquids." *Green Chem.*, vol. 12, pp.1715-1718, 2010.
32. Y. Deng, P. Besse-Hoggan, M. Sanceime et al., "Influence of Oxygen Functionalities on the Environmental Impact of Imidazolium Based Ionic Liquids," *J. Haz. Mat.*, vol. 198, no. 30, pp165-174, 2011.
33. D Mackay, A. Bobra, W. Shiu, et al., "Relationships between Aqueous Solubilities and Octanol-water Partition Coefficients", *Chemosphere*, vol. 9, no. 11, pp.701-711, 1980.
34. S. Banerjee, S. Yalkowsky, S. Valvani, "Water Solubility and Octanol/water Partition Coefficients of Organics. Limitations of the Solubility-partition Coefficient Correlation", *Environ. Sci. and Tech.* vol. 14, no. 10, pp.1227-1229, 1980.
35. D. Hartmann, K. Shimizu, F. Siopa, et al., "Plasma Membrane Permeability by Ionic Liquids: A matter of Charge", *Green Chem.*, vol. 17, pp.4587-4598, 2015.