



GLOBAL JOURNAL OF SCIENCE FRONTIER RESEARCH: D
AGRICULTURE AND VETERINARY
Volume 15 Issue 5 Version 1.0 Year 2015
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Response of Broiler Chickens to Graded Levels of Urea Treated Rice Offal

By S.O. Onuh, E.E. Idogah & E. Ameh

Akperan Orshi College of Agriculture, Nigeria

Abstract- A total of ninety (90) day-old unsexed broiler chickens averaging 50.0 grammes were utilized for the purpose of accessing their response to graded levels of urea treated rice offal. These were randomly allocated into 3 equal groups of 30 birds in each treatment replicated 3 times with each replicate having 10 birds. Three (3) diets designated I, II and III were formulated such that urea treated rice offal was included at 0, 7.5% and 15.0% respectively. The results show that there were no significant differences ($P > 0.05$) in feed intake, weight gain and efficiency of feed utilization among birds fed the control diet and urea treated rice offal based diets. However, feed intake and weight gain were highest with birds fed 15.0% urea treated rice offal compared with those fed the control diet.

Keywords: broilers, chickens, urea, rice offa.

GJSFR-D Classification : FOR Code: 070799p



Strictly as per the compliance and regulations of :



Response of Broiler Chickens to Graded Levels of Urea Treated Rice Offal

S.O. Onuh ^α, E.E. Idogah ^σ & E. Ameh ^ρ

Abstract- A total of ninety (90) day-old unsexed broiler chickens averaging 50.0 grammes were utilized for the purpose of accessing their response to graded levels of urea treated rice offal. These were randomly allocated into 3 equal groups of 30 birds in each treatment replicated 3 times with each replicate having 10 birds. Three (3) diets designated I, II and III were formulated such that urea treated rice offal was included at 0, 7.5% and 15.0% respectively. The results show that there were no significant differences ($P>0.05$) in feed intake, weight gain and efficiency of feed utilization among birds fed the control diet and urea treated rice offal based diets. However, feed intake and weight gain were highest with birds fed 15.0% urea treated rice offal compared with those fed the control diet. Finally, feed cost per unit weight gain significantly improved ($P<0.05$) with birds fed the diet containing 15.0% urea treated rice offal compared with those fed other diets. However, there were no significant differences ($P>0.05$) in feed cost per unit weight gain among birds fed the control diet and those fed the diet containing 7.5% urea treated rice offal. On the basis of the results obtained, it may be recommended that urea treated rice offal could be included at up to 15.0% of the diet of broiler chickens without any adverse effect on their performance.

Keywords: broilers, chickens, urea, rice offa.

I. INTRODUCTION

Nigeria has the potential to produce about 200,000 metric tonnes of rice offal from the 500,000 metric tonnes of rice produced annually (Wudiri, 1991). The offal, therefore, makes up about 40% of the parboiled rice and contains husk, bran polishing and small quantities of broken grains. In spite of its abundance, it has been neglected as animal feeds because it contains high level of fibre and low protein and energy (Oyawoye and Nelson, 1999).

Maikano (2007) reported the proximate composition of rice offal thus; 94.42% dry matter, 5.09% crude protein, 30.39% crude fibre, 3.40% ether extract, 16.67% ash and 46.10% nitrogen free-extract. Several workers (Dafwang and Shwarmen, 1996; (Abasiekong, 1997; Awesu *et al.*, 2002) have reported that the high crude fibre (30 – 44%); mainly lignin and low protein contents have resulted in reduced voluntary feed intake

and low utilization in poultry feeding. This high fibre concentration results in poor nutrient utilization and consequent poor growth performance due to the presence of non-starch-polysaccharides (NSP) and phytate when fed to broiler chickens without any form of treatment.

The use of rice offal to replace cereal grains in poultry diets have been studied (Dafwang and Damang, 1995; Carew *et al.*, 2005) and has been successfully fed to broiler chickens at lower levels of inclusion (Amaefule *et al.*, 2006; Onuh, 2006; Maikano, 2007; Yakubu *et al.*, 2007) order to reduce feed costs. Higher levels of inclusion may therefore necessitate the development of strategies to increase the value of this by-product in order to reduce its fibre content. Alkali treatments of various fibrous materials (Faniyi and Ologhobo, 1999) and urea treatment (Isikwenu *et al.*, 2008; Onuh, 2011) have been reported to improve their nutritional qualities.

The study reported herein was conducted to determine the response of broiler chickens to graded levels of urea treated rice offal.

II. MATERIALS AND METHODS

The study was conducted in the Poultry Unit of the Department of Animal Husbandry, Akperan Orshi College of Agriculture, Yandev-Gboko, Benue State, Nigeria.

The rice offal, containing mainly the husk, used in the present study was collected from Rice Mill in Gboko, Benue State, Nigeria.

The urea treated rice offal was prepared according to the procedures outlined by Isikwenu *et al.* (2008). It was then sun dried for 2 days. The proximate chemical composition of untreated and urea treated rice offal are presented in Table 1.

Table 1 : Proximate chemical composition of urea treated rice offal

| Constituent | ^a Untreated | ^b Treated |
|-------------------|------------------------|----------------------|
| Crude protein (%) | 5.09 | 14.50 |
| Crude fibre (%) | 30.39 | 25.00 |
| Ether extract (%) | 3.40 | 7.37 |
| Ash (%) | 16.67 | 11.52 |

Sources: ^aMaikano (2007); ^bYakubu *et al.* (2007).

Author α: Department of Animal Husbandry.
e-mail: onuhsimokoh@gmail.com

Author σ: Department of Basic Sciences, Akperan Orshi College of Agriculture, Yandev, P.M.B. 181, Gboko-Benue State, Nigeria.

Author ρ: Department of Agricultural Education, Federal College of Education (Technical), Bichi-Kano State, Nigeria.

A total of ninety (90) day-old unsexed broiler chickens obtained from CHI hatchery, Ibadan averaging 50.00 grammes were utilized for the purpose of the study. These were randomly allocated into 3 equal groups of 30 birds in each treatment and brooded after a two day initial stabilization period on deep litter system. Each treatment was replicated 3 times with each replicate having 10 birds.

Three (3) diets designated I, II and III were formulated for broiler chickens such that urea treated rice offal was included at 0, 7.5% and 15.0% respectively. All diets were adequately fortified with vitamins and minerals. The compositions of the broiler starter diets are presented in Table 2 while those of finishing broiler diets are presented in Table 3.

Table 2 : Composition of broiler starter experimental diets

| Ingredient | Dietary Treatments | | |
|-------------------------------------|--------------------|-----------------|-----------------|
| | I Control | II 7.5% UTRO | III 15% UTRO |
| Maize | 45.00 | 40.50 | 33.00 |
| Full-fat Roasted Soyabean | 51.00 | 48.00 | 48.00 |
| Urea Treated Rice Offal | 0.00 | 7.50 | 15.00 |
| Bone Meal | 3.00 | 3.00 | 3.00 |
| Mineral-Vitamin Premix ⁺ | 0.25 | 0.25 | 0.25 |
| DL-Methionine | 0.25 | 0.25 | 0.25 |
| L-Lysine HCl | 0.25 | 0.25 | 0.25 |
| Common Salt | 0.25 | 0.25 | 0.25 |
| <i>Total</i> | <i>100.00</i> | <i>100.00</i> | <i>100.00</i> |
| <i>Calculated Analyses</i> | | | |
| Crude Protein (%) | 23.43 | 22.97 | 23.39 |
| Metabolizable Energy (Kcal/kg) | 3213 | 2961 | 2706 |
| Crude Fibre (%) | 3.71 | 5.33 | 7.05 |
| Methionine (%) | 0.67 | 0.64 | 0.62 |
| Lysine (%) | 1.79 | 1.69 | 1.67 |
| Methionine + Cystine (%) | 0.93 | 0.88 | 0.86 |
| Feed cost/kg (₦/kg) | 74.65 | 69.63 | 64.75 |

UTRO = Urea Treated Rice Offal

⁺Vitamin-mineral premix provided the following vitamins and minerals per kg of diet: A 15,000 I.U; D3 3000 I.U; E 30 I.U; K 2.5mg, B₁ 2.0mg; B₂ 6.0mg; B₆ 4.0mg; Niacin 40mg; B₁₂ 0.02mg; Pantothenic 10mg; Folic 1.0mg; Biotin 0.08mg; Choline Cl 500mg; Antioxidant 125mg; Mn 6mg; Zn 60mg; Fe 24mg; Cu 6mg; I 1.4mg; Se 0.24mg; Co 0.4mg. Product of Agricultural Technologies Nigeria Ltd. Marketed by S&D Farms Abeokuta.

Table 3 : Composition of finishing broiler experimental diets

| Ingredient | Dietary Treatments | | |
|-------------------------------------|--------------------|-----------------|-----------------|
| | I Control | II 7.5% UTRO | III 15% UTRO |
| Maize | 46.00 | 42.50 | 40.00 |
| Full-fat Roasted Soyabean | 38.00 | 34.00 | 29.00 |
| Urea Treated Rice Offal | 0.00 | 7.50 | 15.00 |
| Maize Offal | 6.00 | 6.00 | 6.00 |
| Palm Kernel Cake | 6.00 | 6.00 | 6.00 |
| Bone Meal | 3.00 | 3.00 | 3.00 |
| Mineral-Vitamin Premix ⁺ | 0.25 | 0.25 | 0.25 |
| DL-Methionine | 0.25 | 0.25 | 0.25 |
| L-Lysine HCl | 0.25 | 0.25 | 0.25 |
| Common Salt | 0.25 | 0.25 | 0.25 |
| <i>Total</i> | <i>100.00</i> | <i>100.00</i> | <i>100.00</i> |
| <i>Calculated Analyses</i> | | | |
| Crude Protein (%) | 19.88 | 19.13 | 18.10 |
| Metabolizable Energy (Kcal/kg) | 3105 | 2855 | 2605 |
| Crude Fibre (%) | 4.82 | 6.41 | 6.05 |
| Methionine (%) | 0.61 | 0.58 | 0.54 |
| Lysine (%) | 1.47 | 1.35 | 1.21 |
| Methionine + Cystine (%) | 0.85 | 0.80 | 0.74 |
| Feed cost/kg (₦/kg) | 68.84 | 63.76 | 58.64 |

UTRO = Urea Treated Rice Offal

⁺Vitamin-mineral premix provided the following vitamins and minerals per kg of diet: A 15,000 I.U; D₃ 3000 I.U; E 30 I.U; K 2.5mg, B₁ 2.0mg; B₂ 6.0mg; B₆ 4.0mg; Niacin 40mg; B₁₂ 0.02mg; Pantothenic 10mg; Folic 1.0mg; Biotin 0.08mg; Choline Cl 500mg; Antioxidant 125mg; Mn 6mg; Zn 60mg; Fe 24mg; Cu 6mg; I 1.4mg; Se 0.24mg; Co 0.4mg. Product of Agricultural Technologies Nigeria Ltd. Marketed by S&D Farms Abeokuta.

The birds were held on a basal diet for the first 2 days and monitored for problems that may be associated with hatchery defects and other sources of variations that could cause reduced performance and death which were independent of dietary treatments. The birds were reared according to standard procedures outlined by Dafwang and Ogundipe (1982). The birds were fed each of starter diets for 28 days and thereafter fed each of finishing diets for 21 days. The birds in each treatment were fed weighted amounts of their group diet to appetite daily and fresh water was offered *ad libitum* throughout the period of the study while necessary prophylaxis and vaccinations for broilers were administered. The left over feed was collected and weighed before another days' feeding to determine actual intake. The birds were weighed weekly to determine weight changes. Feed conversion ratio was computed by dividing daily feed intake by the corresponding weight gain. Feed cost per unit weight gain was computed as a product of feed conversion ratio and feed cost per kg of each diet at the time of conducting the study. The dietary feed cost was obtained from the market prices of the different ingredients at the time and locality of the study. Data on feed intake, weight gain, feed: gain ratio and feed cost per unit weight gain were recorded on replicate basis weekly for 49 days.

Data on each parameter were subjected to the analysis of variance (ANOVA) for Completely Randomized Design (CRD) and where significant differences were indicated, the means were separated using Duncan's Multiple Range Test (DMRT) according to the procedures of the Statistical Package (SPSS, 2006).

III. RESULTS

The summary of results of the response of broiler chickens to graded levels of urea treated rice offal is presented in Table 4.

Table 4 : Response of broiler chickens to graded levels of urea treated rice offal

| Parameters | Dietary Treatments | | | SEM |
|---|---------------------|---------------------|---------------------|---------------------|
| | I | II | III | |
| Average daily feed intake (g) | 96.77 | 101.16 | 109.43 | 10.30 ^{NS} |
| Average daily weight gain (g) | 45.27 | 43.03 | 49.53 | 4.95 ^{NS} |
| Feed conversion ratio | 2.14 | 2.35 | 2.21 | 0.30 ^{NS} |
| Feed cost per unit weight gain (₦/gain) | 184.43 ^b | 167.09 ^b | 139.87 ^a | 19.76 |

^{abc}Means followed by the same superscript in horizontal rows are not significantly different ($P>0.05$) from one another; SEM = Standard Error of Mean.

As the results have shown, there were no significant differences ($P>0.05$) in feed intake, weight gain and efficiency of feed utilization among birds fed the control diet and urea treated rice offal based diets. However, feed intake and weight gain were highest with birds fed 15.0% urea treated rice offal compared with those fed the control diet. Weight gain increased by 8.60% when 15.0% urea treated rice offal was fed compared with birds fed the control diet. Furthermore, feed cost per unit weight gain significantly improved ($P<0.05$) with birds fed the diet containing 15.0% urea treated rice offal compared with those fed other diets. However, there were no significant differences ($P>0.05$) in feed cost per unit weight gain among birds fed the control diet and those fed the diet containing 7.5% urea treated rice offal.

IV. DISCUSSION

In the present study, it was observed that 15.0% urea treated rice offal increased feed intake compared with birds fed other diets. The result of the present study is not consistent with those of Iheukwumere *et al.* (2001) and Yakubu *et al.* (2007) who reported lower feed intake with birds fed urea treated rice milling waste when compared to those fed untreated rice milling waste. The high feed intake in the present study could be attributed to the effect of the urea treatment on the fibre content of rice offal. Weight gain at 15.0% level of inclusion of urea treated rice offal was highest compared with birds fed diets fed other diets. The results of the present study is consistent with the reports of Amaefule *et al.* (2006) who reported that broilers fed urea treated rice milling wastes had significantly higher final body weight and daily weight gain than those fed other diets. Efficiency of feed utilization in the present study of birds fed 15.0% urea treated rice offal was better than those fed the diet containing 7.50% urea treated rice offal. The results of the present study which showed that urea treated rice offal resulted in better feed utilization agrees with earlier reports of Abu *et al.* (1999) in rabbits and Iheukwumere *et al.* (2001) in broilers who reported an increase in weight gain and efficiency of feed utilization when birds were placed on diets containing urea treated rice-milling waste. It is known that the chicken is known to be especially sensitive to dietary energy concentration

(Scott *et al.*, 1982). In the present study, the energy content of the diets decreased with increasing levels of urea treated rice offal. Since energy intake is a productive function of feed intake, the higher feed intake of birds fed the urea treated rice offal in the present study could have been responsible for their higher weight gain.

On the basis of the results obtained, it may be recommended that urea treated rice offal could be included at up to 15.0% of the diet of broiler chickens without any adverse effect on their performance.

REFERENCES RÉFÉRENCES REFERENCIAS

1. Abasiekong, S.F. (1997). Effects of termite culture on crude protein, fat and crude fibre contents of fibrous harvest residues. *Bio. Resource Tech.*, 62; 55 – 57.
2. Amaefule, K.J., Iheukwumere, F.C., Lawal, A.S and Ezekwonna, A.A. (2006). The effect of rice milling waste on the performance, nutrient retention, carcass and internal organ characteristics of finisher broilers. *Int. J. Poult. Sci.*, 5(1); 51 – 55.
3. Awesu, J.R., Bamgbose, A.M., Oduguwa, O.O., Fanimu A.O and Oguntola, E.B. (2002). Performance and nutrient utilization of cockerel finishers fed graded levels of rice milling waste. *Nig. J. Anim. Prod.*, 29; 181 – 188.
4. Carew, S.N., Tuleun, C.D and Yaakugh, I.D.I. (2005). The feed value of rice milling by- products in broiler finisher diets. *J. Sustainable Trop. Agric. Res.*, 13; 69 – 73.
5. Dafwang I.J and Shwarmen, E.B.N. (1996). Utilization of rice offal in practical rations for broilers. *Nig. J. Anim. Prod.*, 23(1); 21 – 23.
6. Dafwang, I.I and Damang, P. (1995). Rice offal in finishing diets for broilers. *J. Anim. Prod. Res.*, 15 (1); 131 – 139.
7. Dafwang, I.I and Ogundipe, S.O. (1982). Brooding and rearing of chicks on deep litter. Extension Bulletin No. 23, Poultry series No 3. Agricultural Extension, Research and Liaison Services, Ahmadu Bello University, Zaria, Nigeria. 29p.
8. Faniyi, G. F and Ologhobo, A. D. (1999). Effects of replacing brewers' dried grains with lye-and urea-

- treated cowpea and sorghum seed hulls in broiler diets. *Trop. Ani. Prod. Investigation*, 2; 69 – 82.
9. Ihenkwumere, F.C., Ezekwonna, A.A., Mwoche, G.N and Obeji, C.N. (2001) Effect of treated rice milling wastes on nutrient metabolizability, carcass yield and internal organs weights of finisher broilers. *Proceedings of the 6th Conference of Animal Science Association of Nigeria* held at the University of Maiduguri, September 17th –19th 2001. Pp 20 –23.
 10. Isikwenu, J. O., Omeje, S. I., Okagbare, G and Akpodiete, O. J. (2008). Effect of replacing groundnut cake with urea fermented brewer's dried grains in broiler chick's diets. *Ani. Res. Int.* 5(1): 795– 800.
 11. Maikano, A. (2007). Utilization of rice offal in practical ration of broilers. *The Zoologist*, 5; Pp. 1 – 7.
 12. Onuh, S. O. (2011). The effect of replacing full-fat soybeans with urea fermented sorghum brewers' grain in broiler starter diets. *Ani. Prod. Res. Advances*. 7(4); 282 – 285.
 13. Onuh, S.O. (2006). Evaluation of the performance of finishing broiler chickens fed different agro-industrial by-products. *J. Sustainable Trop. Agric. Res.*, 17; 113 – 115.
 14. Oyawoye, E.O and Nelson, F.S. (1999). Optimum level of inclusion of rice offal in the diet of young cockerels. *Conference of Nigerian Society of Animal Production*, 21st – 25th March 1999 held at the University of Ilorin, Ilorin Nigeria.
 15. Scott, M.L., M.C. Nesheim and R.J. Young, 1982. *Nutrition of the Chicken*. M.L. Scott and Associates, Ithaca. New York. 562 pages.
 16. SPSS (2006). *Statistical Package for Social Sciences*, Version 15.0.
 17. Wudiri, B.B. (1991). Production of cereal crops in Nigeria. From research to farmers' fields. A seminar paper delivered to staff and students at the Plateau State College of Agriculture, Lafia.
 18. Yakubu, B., Adegbola, T.A., Bogoro, S and Yussuf, H.B. (2007). Effect of urea treated and untreated rice offal on the performance of broilers: 1. Growth performances and economy of production. *J. Sustainable Dev. Agric. Env.*, (3); 7 – 13.