



GLOBAL JOURNAL OF SCIENCE FRONTIER RESEARCH: D  
AGRICULTURE AND VETERINARY  
Volume 15 Issue 6 Version 1.0 Year 2015  
Type : Double Blind Peer Reviewed International Research Journal  
Publisher: Global Journals Inc. (USA)  
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

## Nutritional Values of Smoked *Clarias Gariepinus* from Major Markets in Southwest, Nigeria

By Peter Taiwo Olagbemide

*Afe Babalola University, Nigeria*

**Abstract-** Smoked *Clarias gariepinus* from four major markets in the southwest, Nigeria were analyzed. The aim was to evaluate the nutritional values of the fish sold at the different markets. The results of the analyses showed that the nutritional values of the smoked fish are of good standard but nutritional and mineral compositions varied from market to market. The moisture content, protein, fat, fibre, ash and carbohydrate contents from the markets were in the range of 9.63 to 10.27%, 53.77 to 54.77%, 11.77 to 13.13%, 6.87 to 8.00%, 0.0 to 0.07% and 15.40 to 16.17% respectively. The range of the mineral compositions was 10.13 to 12.17mg/100g, 0.33 to 0.50mg/100g, 28.33 to 46.67mg/100g, 353.33 to 388.33mg/100g, 22.33 to 33.33mg/100g, and 271.67 to 305.00mg/100g for iron, zinc, magnesium, calcium, potassium and phosphorus respectively. Ascorbic acid, thiamine, niacin and riboflavin contents of the fish were in range of 0.17 to 0.27mg/100g, 0.05 to 0.07mg/100g, 0.24 to 0.28mg/100g, and 0.05 to 0.08mg/100g respectively.

**Keywords:** smoked fish, markets, southwest, *clarias gariepinus*, nutritional values.

**GJSFR-D Classification :** FOR Code: 079999



NUTRITIONAL VALUES OF SMOKED CLARIAS GARIEPINUS FROM MAJOR MARKETS IN SOUTHWEST NIGERIA

*Strictly as per the compliance and regulations of :*



RESEARCH | DIVERSITY | ETHICS

# Nutritional Values of Smoked *Clarias Gariepinus* from Major Markets in Southwest, Nigeria

Peter Taiwo Olagbemide

**Abstract** - Smoked *Clarias gariepinus* from four major markets in the southwest, Nigeria were analyzed. The aim was to evaluate the nutritional values of the fish sold at the different markets. The results of the analyses showed that the nutritional values of the smoked fish are of good standard but nutritional and mineral compositions varied from market to market. The moisture content, protein, fat, fibre, ash and carbohydrate contents from the markets were in the range of 9.63 to 10.27%, 53.77 to 54.77%, 11.77 to 13.13%, 6.87 to 8.00%, 0.0 to 0.07% and 15.40 to 16.17% respectively. The range of the mineral compositions was 10.13 to 12.17mg/100g, 0.33 to 0.50mg/100g, 28.33 to 46.67mg/100g, 353.33 to 388.33mg/100g, 22.33 to 33.33mg/100g, and 271.67 to 305.00mg/100g for iron, zinc, magnesium, calcium, potassium and phosphorus respectively. Ascorbic acid, thiamine, niacin and riboflavin contents of the fish were in range of 0.17 to 0.27mg/100g, 0.05 to 0.07mg/100g, 0.24 to 0.28mg/100g, and 0.05 to 0.08mg/100g respectively. There were significant differences ( $P < 0.05$ ) in the proximate, vitamin and mineral compositions between the paired samples from the markets. These variations could be explained as due to the different smoking devices employed, the duration and methods of storage and the effects of different fish handlers after smoking. It is therefore recommended that good smoking and storage methods should be adopted as well as maintenance of hygienic environment and practices by smoked fish handlers.

**Keywords:** smoked fish, markets, southwest, *clarias gariepinus*, nutritional values.

## I. INTRODUCTION

Fish makes up about 60% of world protein supply and developing countries derive more than 30% of their annual protein from fish (FAO 1994). Teutscher (1990) and Saisithi (1994) reported that fish provides between 30% and 80% of the total animal protein intake of the coastal people of West Africa. In Nigeria fish constitute 40% of animal protein intake (Olatunde, 1998). Fish demand is increasing as a result of the increasing world population, higher living standards and the good overall image of fish among consumers (Cahu *et al.*, 2004). In addition, the demand for fish is on the increase due to the health benefits of eating fish and due to increase in human population, the rinderpest disaster, and drought bane, which reduce the availability and affordability of red meat (Oshozekhai and Ngueku, 2014). Fish and fish products are highly nutritious with protein content of 15 to 20% and are

particularly efficient in supplementing the cereal and tuber diets widely consumed in Africa (Fagbenro *et al.*, 2005). Kreuzer and Heen (1962); Waterman (1976); Olomu (1995); Ojutiku *et al.* (2009) also highlighted that fish is rich in protein with amino acid composition very well suited to human dietary requirements comparing favorably with egg, milk and meat in the nutritional value of its protein. Fish also contains absorbable dietary minerals (Bruhiyan *et al.*, 1993). In Nigeria, fish is eaten fresh and smoked and form a much cherished delicacy that cut across socio-economic, age, religions and educational barriers (Adebayo *et al.*, 2008) and it is a rich source of protein commonly consumed due to the higher cost of meat and other sources of animal protein (Omolara and Omotayo, 2009). However, fish is highly perishable because it provides favourable medium for the growth of microorganisms after death (Aliya *et al.*, 2012; Oparaku and Mgbenka, 2012). An estimate of 40% postharvest losses of total fish landings have been reported in Nigeria (Akande, 1996). Fish spoilage in Nigeria is influenced to a large extent by high ambient temperatures, considerable distances of landing ports to points of utilization and poor as well as inadequate infrastructure for postharvest processing and landing (Saliu, 2008). Thus, it is imperative to process and preserve some of the fish caught in the period of abundance, so as to ensure an all year round supply. This will invariably reduce postharvest losses, increase the shelf-life of fish, and guarantee a sustainable supply of fish during off season with concomitant increase in the profit of the fishermen (Eyo, 1997). Proper preservation starts the moment fish is harvested until reaches the consumer's table (Oluborode *et al.*, 2010). A number of processing techniques is in operation in Nigeria. These include chilling, freezing, salting, canning, drying and smoking. However, smoking is the most popular method of fish processing (Eyo, 2000) and among the several methods of long term preservation of fish, smoking is perhaps the simplest method as it does not require sophisticated equipment or highly skilled workers (Olayemi *et al.*, 2011). Smoked-dried fish is the most acceptable form of fish product in Nigeria (Stolyhiro and Sikorski, 2005; Yanar, 2007). Smoking is the oldest and most common method of fish preservation in many developing countries (Krasemann 2004; Kumolu-Johnson *et al.*, 2010). It is a method of preservation effected by combination of drying and decomposition of naturally produced chemical resulting

**Author:** Department of Biological Sciences, Afe Babbalola University, Ado-Ekiti, Nigeria. e-mail: Petseko2004@yahoo.co.uk

from thermal breakdown of wood (Tobor, 2004). The smoke is produced by the process of incomplete combustion of wood in order to impart a characteristic flavour and colour to the fish. Smoke contributes to fish preservation and shelf life by drying, cooking, acting as an effective antioxidant, bacteriostatic and bactericidal agent as well as by depositing natural wood-smoke chemicals like tars, phenols and aldehydes; all of which provide a protective film on the surface of smoked fish and have powerful bactericidal action and prevent the growth of other microorganisms on the flesh of the fish (Gilbert and Knowles, 1995; Horner, 1997; Doe, 1998; Rorvik, 2000; Garrow and James, 2000; Daramola *et al.*, 2007; Ahmed *et al.*, 2010; Daramola *et al.*, 2013). Several methods are available for fish smoking and different smoked products have been developed in various parts of the world in relation to the properties of the locally available raw materials and the general level of technology (Olley *et al.*, 1988).

African catfish (*Clarias gariepinus*) is one of the most important fish species currently being cultured both inside and outside its natural range of tropical and subtropical environments (Adewolu *et al.*, 2008). Positive attributes such as resistance to diseases, high fecundity, and ease of larval production in captivity make it of commercial importance in aquaculture (Haylor, 1991). It is of great importance as it grows quickly, attains a large size, and is an edible fish with few spines in its flesh. It can withstand wide range of environmental conditions, including severe temperatures, as well as low oxygen. The importance of catfish itself cannot be overemphasized. According to Anoop *et al.*, (2009), it provides food for the populace, it allows for improved protein nutrition because it has a high biological value in terms of high protein retention in the body, higher protein assimilation as compared to other protein sources, low cholesterol content and one of the safest sources of animal protein.

The aim of this study is to evaluate the nutritional value of the smoked *Clarias gariepinus* sold at the major markets in Southwest, Nigeria.

## II. MATERIALS AND METHODS

### a) Sample collection

The fish samples used for this study were purchased from four major markets in Oyo and Ekiti States in Southwest, Nigeria. Two major markets (A and B) were selected in Ibadan, the capital of Oyo State and two major markets (C and D) in Ado-Ekiti, the capital of Ekiti State. These markets were selected because they are markets generally patronized by the populace. All samples were transported to the Department of Biological Sciences, Afe Babalola University for analyses.

### b) Analysis of samples

The proximate analyses of the all the samples from the various markets (A, B, C and D) for moisture, ash and carbohydrate contents were determined as described by AOAC (2005). Crude protein, fibre and fat contents were determined using the methods described by Pearson (1976). Mineral contents of all the samples were determined by atomic absorption spectrometry, flame photometry and spectrophotometry according to the methods of AOAC (2003). The phytochemical analysis for the presence of saponins, tannins, alkaloids, and cyanogenic glycosides in the samples were carried out according to the methods described by Harborne (1973) and Trease and Evans (1983). The vitamins in the samples from the four markets were determined by the official methods of the Association of official Analytical chemists (AOAC, 1990). Each analysis was carried out in triplicate.

### c) Statistical analysis

All assays were carried out in triplicate, and the means and standard error of means (SEM) were determined using SPSS version 20. Analysis of variance was performed to determine significant differences between the paired samples. Differences in paired samples performance for each nutrient and chemical composition were tested by the Student's t-test.  $<0.05$  implies significance.

## III. RESULTS

The proximate analyses of the samples from the different markets are shown in Table 1 while Table 2 shows the paired samples test of the proximate analyses of the samples from the markets. The values of the proximate analyses in the samples varied from market to market. The moisture content, protein, fat, fibre, ash and carbohydrate contents from the markets were in the range of 9.63 to 10.27%, 53.77 to 54.77%, 11.77 to 13.13%, 6.87 to 8.00%, 0.0 to 0.07% and 15.40 to 16.17% respectively. Samples from Market A and Market D recorded higher values of moisture contents than samples from Markets B and C. Table 2 shows that there were significant differences in the moisture content values between the samples from Market A and Markets B, C and between samples from Market D and Markets B, C. There was no significant difference in moisture values between samples from Market A and D and between samples from Market B and C. The protein content values of the samples from the markets were high but showed variation in the markets. There was a significant difference in the protein values of the samples between Market A and Market B and between Market B and Market C. The fat content showed significant difference only between samples from Market C and Market D. There was no significant different in the crude fibre and carbohydrates values of the samples from the markets.

Table 1 : Proximate Analyses of *Clarias gariepinus* from the four markets

Parameters	Market A	Market B	Market C	Market D
Moisture content (%)	10.13±0.09	9.63±0.09	9.67±0.09	10.27±0.09
Protein (%)	54.53±0.09	53.77±0.07	54.77±0.09	54.17±0.09
Ether extracts (fat) %	12.17±0.09	12.33±0.09	13.13±0.09	11.77±0.07
Ash (%)	7.63±0.09	8.00±0.07	6.87±0.09	7.53±0.09
Crude fibre (%)	0.07±0.03	0.00	0.00	0.03±0.03
Carbohydrate (by difference) %	15.43±0.09	16.17±0.12	15.40±0.17	16.03±0.15

Values are means ±SEM (Standard error of means) of triplicate samples.

Table 2 : Paired samples test of proximate analyses of *Clarias gariepinus* from the markets

Parameters	Paired samples	Diff.Mean	Sig. (2-Tailed)
Moisture content (%)	Market A-Market B	0.50±0.12	0.049
	Market A-Market C	0.47±0.09	0.034
	Market A-Market D	-0.13±0.03	0.057
	Market B-Market C	-0.03±0.17	0.860
	Market B-Market D	-0.63±0.09	0.019
	Market C-Market D	-0.60±0.12	0.035
Protein (%)	Market A-Market B	0.77±0.12	0.024
	Market A-Market C	-0.23±0.09	0.118
	Market A-Market D	0.37±0.09	0.053
	Market B-Market C	1.00±0.05	0.030
	Market B-Market D	-0.40±0.15	0.120
	Market C-Market D	0.60±0.15	0.590
Ether extract (fat) %	Market A-Market B	-0.17±0.09	0.200
	Market A-Market C	-0.97±0.09	0.080
	Market A-Market D	0.40±0.17	0.148
	Market B-Market C	-0.80±0.15	0.350
	Market B-Market D	0.57±0.13	0.051
	Market C-Market D	1.37±0.17	0.015
Ash content (%)	Market A-Market B	-0.37±0.13	0.111
	Market A-Market C	0.77±0.18	0.049
	Market A-Market D	0.10±0.15	0.580
	Market B-Market C	1.13±0.07	0.003
	Market B-Market D	0.47±0.03	0.005
	Market C-Market D	-0.67±0.09	0.017
Crude fibre (%)	Market A-Market B	0.07±0.03	0.184
	Market A-Market C	0.07±0.03	0.184
	Market A-Market D	0.03±0.03	0.423
	Market B-Market C	.....	.....
	Market B-Market D	-0.03±0.03	0.423
	Market C-Market D	-0.03±0.03	0.423
Carbohydrate (by difference) %	Market A-Market B	-0.73±0.07	0.008
	Market A-Market C	0.03±0.17	0.860
	Market A-Market D	-0.60±0.23	0.122
	Market B-Market C	0.77±0.23	0.081
	Market B-Market D	0.13±0.24	0.635
	Market C-Market D	-0.63±0.28	0.156

Significant difference ( $P < 0.05$ )

Table 3 shows the values of mineral compositions of the samples from the markets while Table 4 shows the paired samples test of the mineral compositions of the samples from the markets.

Table 3 : Mineral compositions of *Clarias gariepinus* from the four markets

Parameters (mg/100g)	Market A	Market B	Market C	Market D
Iron (Fe <sup>++</sup> )	11.33±0.09	10.13±0.09	12.17±0.09	11.57±0.09
Zinc (Zn <sup>++</sup> )	0.43±0.03	0.40±0.06	0.50±0.06	0.33±0.03
Magnesium (Mg <sup>++</sup> )	46.67±1.67	36.67±1.67	28.33±1.67	40.00±2.89
Calcium (Ca <sup>++</sup> )	353.33±6.01	373.33±7.26	358.33±4.41	388.33±1.67
Potassium (K <sup>+</sup> )	33.33±1.67	30.00±2.89	25.00±2.89	22.33±1.45
Phosphorus (PO <sub>4</sub> <sup>---</sup> )	280.01±2.89	288.33±4.41	271.67±4.41	305.00±2.89
Ca/P	1.26±0.02	1.30±0.01	1.32±0.04	1.27±0.02
Ca/Mg	7.60±0.36	10.22±0.47	12.75±0.84	9.81±0.67
Ca/K	10.65±0.53	12.71±1.36	14.67±1.76	17.54±1.19

Values are means ±SEM (Standard error of means) of triplicate samples.

Table 4 : Paired samples test of the mineral composition of *Clarias gariepinus* from the markets

Parameters	Paired samples	Diff. Mean	Sig. (2-Tailed)
Iron	Market A-Market B	1.20±0.15	0.016
	Market A-Market C	-0.83±0.18	0.042
	Market A-Market D	-0.23±0.17	0.300
	Market B-Market C	2.03±0.09	0.002
	Market B-Market D	1.43±0.03	0.001
	Market C-Market D	0.60±0.06	0.09
Zinc	Market A-Market B	0.03±0.07	0.667
	Market A-Market C	-0.07±0.03	0.184
	Market A-Market D	-----	-----
	Market B-Market C	-0.10±0.01	0.423
	Market B-Market D	0.07±0.07	0.423
	Market C-Market D	0.17±0.03	0.038
Magnesium	Market A-Market B	10.00±2.89	0.074
	Market A-Market C	18.33±1.67	0.008
	Market A-Market D	6.67±4.41	0.270
	Market B-Market C	8.33±3.33	0.130
	Market B-Market D	-3.33±1.67	0.184
	Market C-Market D	11.67±4.41	0.118
Calcium	Market A-Market B	20.00±2.89	0.020
	Market A-Market C	5.00±2.89	0.225
	Market A-Market D	35.00±5.00	0.020
	Market B-Market C	15.00±2.89	0.035
	Market B-Market D	15.00±5.77	0.122
	Market C-Market D	30.00±2.89	0.009
Potassium	Market A-Market B	3.33±1.69	0.184
	Market A-Market C	8.33±4.41	0.199
	Market A-Market D	11.00±2.08	0.034
	Market B-Market C	5.00±5.00	0.423
	Market B-Market D	7.67±2.33	0.081
	Market C-Market D	2.67±2.67	0.423
Phosphorus	Market A-Market B	8.33±4.41	0.199
	Market A-Market C	8.33±6.00	0.300
	Market A-Market D	25.00±5.00	0.038
	Market B-Market C	16.67±8.82	0.199
	Market B-Market D	16.67±7.26	0.149
	Market C-Market D	33.33±1.67	0.02
Ca/P	Market A-Market B	-0.04±0.01	0.093
	Market A-Market C	-0.06±0.02	0.122
	Market A-Market D	-0.01±0.00	0.057

	Market B-Market C	-0.02±0.03	0.551
	Market B-Market D	0.02±0.01	0.222
	Market C-Market D	0.05±0.02	0.148
Ca/Mg	Market A-Market B	-2.63±0.53	0.039
	Market A-Market C	-5.15±0.76	0.021
	Market A-Market D	-2.21±0.97	0.150
	Market B-Market C	-2.52±1.25	0.181
	Market B-Market D	0.42±0.58	0.550
	Market C-Market D	2.94±1.48	0.186
Ca/K	Market A-Market B	-2.05±0.85	0.138
	Market A-Market C	-4.01±2.09	0.195
	Market A-Market D	-6.89±1.10	0.025
	Market B-Market C	-1.96±2.52	0.517
	Market B-Market D	-4.84±1.15	0.052
	Market C-Market D	-2.88±1.43	0.181

Significant difference ( $P < 0.05$ )

The values of the mineral composition in the samples varied from market to market. The range was 10.13 to 12.17mg/100g, 0.33 to 0.50mg/100g, 28.33 to 46.67mg/100g, 353.33 to 388.33mg/100g, 22.33 to 33.33mg/100g, and 271.67 to 305.00mg/100g for iron, zinc, magnesium, calcium, potassium and phosphorus respectively. There was a significant difference in iron content of the samples between Market A and Markets B, C and between Market B and Markets C, D. There was a significant difference in the values of zinc in the samples between Market C and Market D. Magnesium in the samples showed significant difference between Market A and Market C. There was a significant difference in the values of calcium in the samples between Market A and Markets B, D; between Market B and Market C and between Market C and Market D. Potassium showed significant difference in the samples between Market A and Market D while Phosphorus showed significant difference in the samples between Market A and D and between Market C and Market D.

Table 5 shows phytochemical and vitamin compositions of the samples from the markets while Table 6 shows the paired samples test of the phytochemical and vitamin compositions of the samples from the markets. The phytate, saponin, ascorbic acid, thiamine, niacin and riboflavin contents were in the range of 21.67 to 28.33mg/100g, 0.00 to 0.17mg/100g, 0.17 to 0.27mg/100g, 0.05 to 0.07mg/100g, 0.24 to 0.28mg/100g, and 0.05 to 0.08mg/100g respectively. Thiamine and phytates showed no significant difference in their values in the samples between the markets while ascorbic acid and riboflavin showed significant difference in their values in the samples only between Market C and Market D. Niacin showed significant difference in its values in the sample between Market A and Market B and between Market B and Market C while saponin showed significant difference in its value in the samples between A and Market C and between Market C and Market D.

Table 5: Phytochemical and vitamin compositions of *Clarias gariepinus* from the markets

Parameters	Market A	Market B	Market C	Market D
Pyhtates	28.33±1.67	21.67±1.67	25.00±2.89	28.33±1.67
Saponins	0.17±0.03	0.17±0.09	0.00±0.00	0.17±0.03
Ascorbic acid	0.27±0.03	0.20±0.06	0.17±0.06	0.33±0.06
Thiamine	0.06±0.00	0.06±0.03	0.07±0.01	0.05±0.01
Niacin	0.24±0.01	0.27±0.01	0.23±0.01	0.28±0.01
Riboflavin	0.07±0.00	0.06±0.00	0.05±0.01	0.08±0.00

Values are means ±SEM (Standard error of means) of triplicate samples.

Table 6: Paired samples test of the phytochemical and vitamin of *Clarias gariepinus* from the markets

Parameters	Paired samples	Diff. Mean	Sig. (2-Tailed)
Phytates	Market A-Market B	6.67±1.67	0.057
	Market A-Market C	3.33±1.67	0.184
	Market A-Market D	0.00±2.89	1.000
	Market B-Market C	-3.33±3.33	0.423
	Market B-Market D	-6.67±3.33	0.184

	Market C-Market D	3.33±3.33	0.423
Saponins	Market A-Market B	0.00±0.12	1.000
	Market A-Market C	0.17±0.03	0.038
	Market A-Market D	0.00±0.06	1.000
	Market B-Market C	0.17±0.09	0.199
	Market B-Market D	0.00±0.06	1.000
	Market C-Market D	-0.17±0.03	0.038
	Ascorbic acid	Market A-Market B	0.07±0.07
Market A-Market C		-----	-----
Market A-Market D		-0.07±0.03	0.184
Market B-Market C		0.03±0.07	0.667
Market B-Market D		-0.13±0.09	0.270
Market C-Market D		-0.02±0.03	0.038
Thiamine	Market A-Market B	-0.00±0.00	0.423
	Market A-Market C	0.01±0.01	0.529
	Market A-Market D	0.01±0.00	0.184
	Market B-Market C	-0.00±0.01	0.667
	Market B-Market D	-----	-----
	Market C-Market D	0.01±0.01	0.184
Niacin	Market A-Market B	-0.03±0.01	0.035
	Market A-Market C	0.01±0.01	0.383
	Market A-Market D	-0.04±0.02	0.120
	Market B-Market C	0.04±0.01	0.023
	Market B-Market D	-0.01±0.02	0.580
	Market C-Market D	-0.05±0.01	0.067
Riboflavin	Market A-Market B	-----	-----
	Market A-Market C	0.02±0.01	0.199
	Market A-Market D	-0.02±0.00	0.038
	Market B-Market C	0.01±0.01	0.529
	Market B-Market D	-0.03±0.00	0.015
	Market C-Market D	-0.03±0.06	0.038

Significant difference ( $P < 0.05$ )

#### IV. DISCUSSION

The moisture content can be used as a pointer to the rate at which deterioration occurs in fish samples resulting in the early decomposition. The moisture content recorded in the samples in the Markets A-D is within the range (9-13%) recorded by Plahar *et al.* (1996) and is considered to be low enough to present little deterioration problems if storage conditions are properly controlled. The low moisture content is to reduce to minimum the conditions in the fish that allow for spoilage organisms and chemical activities. Kaneko (1976) reported that a lot of proteolytic, lipolytic deterioration and microbial proliferation are encouraged at moisture levels of 15% and above. The results of the proximate compositions in this study were slightly different from those of Adebowale *et al.* (2008) who reported the range of moisture, protein, fat and ash content of Nigerian smoked catfish to be 7.16-10.71, 33.66-66.04, 1.58-6.09 and 9.21-12.16%, respectively. The low crude fibre value recorded in the samples from the markets is due to the fact that the energy content in smoked *Clarias gariepinus* is high because crude fiber is considered as indigestible. The crude fibre content indicates the amount of cell walls in the feed. The fat levels in the samples from the four markets were below the range (15-33%) reported by Plahar *et al.* (1991) to

cause rancidity problems in storage. In this study, the crude protein formed the largest quantity of the dry matter in all the fish samples. This is in-line with the report that protein forms the largest quantity of dry matter in fish (Pannevis, 1993) and thus, smoked *Clarias gariepinus* is a good source of pure protein and would be more than enough to prevent malnutrition in children and adult who feed solely on this fish as a main source of protein. It also clear from the results of this study that smoked *Clarias gariepinus* is a good source of macro and micro mineral elements in spite of the processing effects of smoking and may contribute to health, growth and development of human beings. The ratios of the mineral compositions further point out the nutritional values of the fish as reported by Watts (2010) that determining nutritional interrelationships is much more important than knowing mineral level alone. Mineral ratios are often more important in determining nutritional deficiencies and excess; it is predictive of future metabolic dysfunctions or hidden metabolic dysfunction. The high Ca/P ratio observed in all the samples from the markets is of nutritional benefit, particularly for children and the aged who need higher intakes of calcium and phosphorus for bone formation and maintenance. Food is considered 'good' if the ratio is above one and 'poor' if the ratio is less than 0.5 while Ca/P ratio above two helps to increase the absorption of calcium in the small

intestine (Niemann *et al.*, 1992). Ca/K ratio is usually called thyroid ratio because calcium and potassium play a vital role in regulating thyroid activity and the ratio in this study is around the range (8-16) needed to maintain the regulation of thyroid activity in good balance (ARL, 2012). Ca/Mg ratio in the fish from the markets is within the range that enhances mental and emotional stability whereas ratio beyond 16 or less than 2 is associated with mental and emotional disturbances (ARL, 2012). The relatively small amount of zinc content recorded in smoked fish from all the markets is not surprising since zinc is a trace mineral and is needed only in small amounts by our bodies but has many important functions. It is needed for the body's defensive (immune) system to properly work; plays a role in cell division, cell growth, wound healing and the breakdown of carbohydrates and is also needed for the senses of smell and taste. The presence of riboflavin, niacin, thiamine and ascorbic acid in all the samples from the markets is a pointer to the nutritional value of smoked *Clarias gariepinus*. Riboflavin is important for body growth and the production of red blood cells; niacin helps maintain healthy skin and nerves and ensures that the digestive and nervous systems function properly; thiamine helps the body cells to change carbohydrates into energy and ascorbic acid helps the body to make collagen, an important protein used to make skin, cartilage, tendons, ligaments, and blood vessels and is also needed for healing wounds, and for repairing and maintaining bones and teeth. The high value of phytates recorded in all the samples from the markets indicates that fishes are likely sources of phytates to their environments. This is because ruminant animals (e.g., cattle, sheep, goats, buffalo) possess phytase producing flora for digesting phytates while non-ruminant animals (e.g., pigs, chickens, dogs, cats, fish) don't have phytase producing flora and as a consequence of low digestibility of phytates by fish, most of the phytates end up being excreted and make their way into their immediate environments. The relative small quantity of saponins in all the samples from the markets is likely to be due to their toxicity to cold-blooded animals at certain concentration. Francis *et al.* (2002) reported that saponins are toxic to fish.

However, there were variations in values of the proximate analyses and the chemical compositions of *Clarias gariepinus* from the markets sampled and these variations are likely to affect the wholeness, safety and shelf life of the products. Variations in proximate and chemical compositions of smoked fish are said to be caused by different factors, such as fish species, smoking methods, smoking time and salt concentration (Adegunwa *et al.*, 2013). Huda *et al.* (2010) reported that nutrient content of fish is influenced by several factors including smoking method and time and the nutrient composition of locally available foods. According to Swastawati (2004) differences in smoked fish flesh

composition are due to different fish species and smoking methods and Dvorak and Vognsrova (1997) reported that difference in smoke quality can make the end-products to differ with respect to nutritional esthetic quality. This is because, according to Kostyra and Pikielna (2006), different smoke sources produce different complex smoke compounds that could consist of mixture of various volatile and non volatile compounds, such as phenol, syringol and guaiacol and its derivatives that affect the quality of the smoked fish. In this study, the variations observed in proximate and chemical compositions of the samples from the different markets may be due to smoking methods and time and not because of difference in species. This is because traditional fish smoking devices are poorly constructed; the technology employed by local fishermen in smoking is not standardized and lack mechanisms for the control of smoke and heat production so that most parameters remain uncontrolled. Hence, essential drying parameters such as duration, air humidity and temperature which affect the efficiency of smoking and the quality of the final products (Olopade *et al.*, 2013) are not precisely determined and mastered. In addition, the variations in the nutritional value of *Clarias gariepinus* from different markets may also be due to the storage methods and the durations of storage after smoking. Smoking decreases the water activity in fish tissue (Sveinsdottir, 1998) and if the smoked fish is not properly stored afterward, the efforts involved in smoking may not yield the expected preservative effect. Jallow (1995) reported that fish with 10-15% moisture content has a shelf life of 3-9 months when stored properly. Thus the concentrations of chemicals in smoked fish are contingent on the storage time and temperature. It is therefore necessary to consider the recommendations of Daramola *et al.* (2007) that "intermittent sun-drying or mild smoking can be carried out on smoked fish to extend its shelf life; moisture content less than 10% should be maintained in stored smoked fish to reduce the growth of bacteria and moulds and preservatives such as pirimiphos-methyl (actelic) can be applied to preserve smoked fish." Thus, this study advocates the need for the adoption of good processing practices and storage methods of smoked fish. In addition, the people that are involved in the processing and selling of smoked fish should maintain hygienic environment and practices so as to ensure that safety standards are maintained in smoked fish and market worthiness of the products is preserved.

## V. CONCLUSION

The nutritional and chemical compositions of smoked *Clarias gariepinus* showed variations from market to market; the results however indicate that smoking method is an important preservation method which could preserve the nutritive values of fishes and



possibly reduce post-harvest losses. However, good smoking method should be adopted and hygienic and proper storage devices put in place.

## VI. ACKNOWLEDGMENTS

I would like to thank Mr. Adarabioyo, M. I. for helping with the statistical analyses of this work

## REFERENCES RÉFÉRENCES REFERENCIAS

- Adebayo-Tayo, B. C., Onilude, A. A. and Patrick, U. G. (2008). Mycofloral of smoke-dried fishes sold in Uyo, Eastern Nigeria. *World J. Agric. Science*, 4(3):346-350.
- Adebowale, B. A., Dongo, L. N., Jayeola, C. O. and Orisajo, S. B. (2008). Comparative quality assessment of fish (*Clarias gariepinus*) smoked with cocoa pod husk and three other different smoking materials. *J. Food. Technology*, 6:5-8.
- Adegunwa, M. O., Adebowale, A. A., Olisa, Z. G. and Bakare, H. A. (2013). Chemical and microbiological qualities of smoked herring (*sardinella eba*, valenciennes 1847) in Odeda, Ogun state, Nigeria. *International Journal of microbiology research and reviews*, vol. 1(5):085-087.
- Adewolu, M. A., Adeniji, C. A. and Adejobi, A. B. (2008). "Feed utilization, growth and survival of *Clarias gariepinus* (Burchell 1822) fingerlings cultured under different photoperiods," *Aquaculture*, vol. 283, (1-4):64-67.
- Ahmed, E. O., Ali, M. E., Kalid, R. A., Taha, H. M. and Mahammed, A. A. (2010). Investigating the quality changes of raw and hot smoked *Oreochromis niloticus* and *Clarias lazera*. *Pakistan Journal of Nutrition*, 9(5):481-484.
- Akande, G. R. (1996). "Post-harvest processing in Fisheries". A paper presented at training for officers of UNDP assisted programme on artisanal fisheries development, Ogun State at Federal College of Fisheries and Marine Technology, Lagos, pp. 1-20.
- Aliya, G., Humaid, K., Nasser, A., Sami, G., Aziz, K., Nashwa, M. and Ponnerassery, S. S. (2012). Effect of the freshness of starting material on the final product quality of dried salted shark. *Advance J. Food Sci. Technology*, 4(2): 60-63.
- Anoop K. R., Sundar K. S. G., Khan B. A., Lal S., (2009) Common Moorhen *Gallinula chloropus* in the diet of the African catfish *Clarias gariepinus* in Keoladeo Ghana National Park, India. *Indian Birds* 5(2):22-23.
- AOAC (1990). *Official methods of analysis of the Association of Official's Analytical Chemist* (15<sup>th</sup> edition) Williams S. (ed). Association of Official Analytical Chemists, Washington, D.C. PP. 152-164.
- AOAC (2003). *Official methods of analysis of the Association of Official's Analytical Chemist*, 17<sup>th</sup> Edn. Association of Official Analytical Chemists, Arlington, Virginia.
- AOAC (2005). *Official Methods of Analysis* (18th edition) Association of Official Analytical, Chemists International, Maryland, USA.
- ARL (2012). Basic ratios and their meaning. *Analytical research labs. Inc.* (602) 955-1580. [www.arlma.com/Articl/Ratio.Doc.htm](http://www.arlma.com/Articl/Ratio.Doc.htm).
- Bruhiyan, A. K. M., Ratnayake, W. M. N. and Aukman, R, G. (1993). Nutritional composition of raw fish and smoked Atlantic mackerel, oil and water soluble vitamins. *J. Food comp Anal.* 6:172-184.
- Cahu, C., Salen, P. and De Lorgeril, M. (2004). Farmed and wild fish in the prevention of cardiovascular diseases: assessing the possible differences in lipid nutritional values. *Nutr Metab Cardiovas*, 14:34-41.
- Daramola, J. A., Fasakin, E. A. and Adeparusi, E. O. (2007). Changes in Physicochemical and Sensory Characteristics of Smoked Dried Fish Species Stored at Ambient Temperature. *African Journal of Food, Agriculture Nutrition and Development*, 7 (6): 1684-5358.
- Daramola, J. A., Kester, C. T. and Allo, O. O. (2013). Biochemical changes of hot smoked African catfish (*Clarias gariepinus*) Samples from sango and Ota Markets in Ogun State. *The Pacific Journal of Science and Technology*, 14 (1): 380-386.
- Doe, P. E. (1998). *Fish drying and smoking, production and quality*. Technomic Publishing Co. Inc., Lancaster, PA., pp. 89-115.
- Dvorak, I. V. and Vognsrova, I. (1997). Available lysine in meat and meat products. *J. Sci. Food Agric.*, 16: 305.
- Eyo, A. A. (1997). Post harvest losses in the fisheries of Kainji Lake. *A consultancy report submitted to Nigerian/German (GTZ) kanji lake fisheries promotion project*, p. 75.
- Eyo, A. A. (2000). *Fish processing technology in the tropic*, University of Ibadan Press. pp. 165-168.
- Fagbenro, O. A., Akinbulumo, M. O., Adeparusi, O. E and Raji, A. A. (2005). Flesh yield, waste yield, proximate and mineral composition of four commercial West African freshwater food fishes. *J. Anim. Vet. Adv.*, 4(10):848-851.
- FAO (1994). Review of the state of world fishery resources: Marine Fisheries. *FAO Fisheries Circular*, No 920. Rome.
- Francis, G., Kerem, Z., Makkar, H. P. S. and Becker, K. (2002). The biological action of saponins in animal system: a review. *British Journal of nutrition*, 88: 587-605.
- Garrow, J. S. and James, W. P. T. (2000). *Human Nutrition and Dietetics*. London: Churchill Living Stone, pp. 84.

- Gilbert, J. and Knowles, M. E. (1995). The chemistry of smoked foods: A review. *Journal of Food Technology*, 10:245–261.
- Harborne, I. B. (1973). *Phytochemical methods: A guide to modern technology of plant analysis*, 2nd ed, New York, Chapman and Hall, pp: 88-185.
- Haylor, G. S. (1991). "Controlled hatchery production of *Clarias gariepinus* (Burchell 1822): growth and survival of fry at high stocking density," *Aquaculture & Fisheries Management*, vol. 22, (4) 405–422.
- Horner, W. F. A. (1997). Preservation of fish by curing, drying, salting and smoking. In: *Fish processing Technology*, Hall, G. M. (Ed). 2<sup>nd</sup> Edn., Blackie Academic and Professional, London; pp. 32-73.
- Huda, N., Deiri, R. S and Ahmed, R. (2010). Proximate, color and amino acid profile of Indonesians traditional smoked catfish. *J. Fish. Aquat. Science*, 5:106-112.
- Jallow, A. M. (1995). Contribution of Improved chorkor oven to artisanal fish smoking in the Gambia. Workshop on seeking improvement in Fish Technology in West Africa. *IDAF Technical Report*, No. 66.
- Kaneko, S. (1976). Smoked meat and microorganisms. *New Food Ind.* 18, 17-23. In *A review of Japanese studies. Fish smoking and drying. The effect of smoking and drying on the nutritional properties of fish* (ed. T. Moto)( 1988). Elsevier Applied Science (ed. J. R. Burt), pp. 91-120.
- Kostyra, E. and Pikielna, N. B. (2006.) Volatiles composition and flavour profile identity of smoke flavourings. *Food Quality and Preference*, 17: 85-95.
- Krasemann, S. (2004). Smoking seafood. "A History of smoke preservation" <http://www.3men.com/history.htm>.
- Kreuzer, R. and Heen, E. (1962). *Fish Nutrition*. London Fishing News Book Ltd. 445p.
- Kumolu-Johnson, N. F., Aladetohun, C. A. and Ndimele, A. (2010). The effect of smoking on the nutrient composition of the African cat fish (*Clarias gariepinus*). *African Journal of Biotechnology*, 9:73-76.
- Niemann, D. C., Butterworth, D. E. and Nieman, C. N. (1992). Pp. 237–312 in *Nutrition*. WmC. Brown, Dubuque, IA.
- Ojutiku, R. O., Kolo, R. J. and Mhammed, M. L. (2009). Comparative study of sun drying and solar tent drying of *Hyperopisus bebeoccidentalis*. *Pak. J. Nutr.* 8(7):955-957.
- Olatunde, A. A. (1998). Approach to the study of fisheries biology in Nigeria Inland water. *Proceedings of International Conference of two decades of Research in Lake Kainji*, pp. 335-341.
- Olayemi, F. F., Adedayo, M. R., Bamishaiye, E. I. and Awagu, E. F. (2011). Proximate composition of catfish (*Clarias gariepinus*) smoked in Nigerian stored products research institute (NSPRI) developed kiln. *Int. J. Fisheries Aquaculture*, 3 (5):96-98.
- Olley, J., Doe, P. E. Heruvati, E. S. (1988). The influence of drying and smoking on the nutritional properties of fish In: *Fish smoking and drying*, Editor. Burt, J. R. PP. 1-11 London and New York:Elsevier applied Science.
- Olomu, J. M. (1995). *Monogastric Animal Nutrition* Jachem Publications, Benin City. Pp. 165-200.
- Olopade, O. A., Taiwo, I. O. and Agbato, D.A. (2013). Effect of Traditional smoking Method on nutritive values and organoleptic properties of *Sarotherodon galilaeus* and *Oreochromis niloticus*. *International Journal of Applied Agricultural and Apicultural Research*, 9 (1&2): 91-97.
- Oluborode, G. B., Omorinkoba, W. S. and Bwala, R. L. (2010). Development and Construction of an Electric Furnace and Control System for fish drying. *Afr. J. Eng. Res.Dev. (Devon Science Publication)*, 3 (2):123-128. 2010.
- Omolaro O. O. and Omotayo O. D. (2009). Preliminary Studies on the effect of processing methods on the quality of three commonly consumed marine fishes in Nigeria. *BIOKEMISTRI*. 21(1): 1-7.
- Oparaku, N. F. and Mgbenka, B. O. (2012). Effects of electric oven and solar dryer on a proximate and water ctivity of *Clarias gariepinus* Fish. *European J. Sci. Res.* 81(1):139-144.
- Oshozekhai, F. E. and Ngueku, B. B. (2014). The use of wood shavings as an alternative fuel wood in fish smoking. *International Journal of Fauna and Biological Studies*, 1 (6): 126-130.
- Pannevis, M. C. (1993). Nutrition of ornamental fish. In: I Burger (Ed). *The Watham Book of Companion Animal Nutrition*. Pergamon Press Ltd, Oxford, England, pp. 85-96.
- Pearson, D. (1976). *The chemical analysis of foods*. 7<sup>th</sup> Ed. London, Churchill Livingstone, pp: 3-4.
- Plahar, W. A., Nti, C. A. and Steiner-Asiedu, M. (1996). Studies on fish consumption patterns and fish quality at household level in the middle belt districts of Ghana. A *Project Report submitted under the Ghana/Netherlands Regional Artisanal Fish Processing and Applied Research Project*. Food Research Institute, CSIR, Accra, Ghana.
- Plahar, W. A., Pace, R. D. and Lu, J. Y. (1991). Effects of storage methods on the quality of smoked-dry herring (*Sardinella eba*). *Jnl Sci. Fd Agric.* 57, 597- 610.
- Rorvik, L. M. (2000). *Listeria monocytogenes* in the smoked salmon industry, *International Journal of food Microbiology*, 62:183-190.

Saisithi, P. (1994). Traditional fermented fish source production. In *Fisheries Processing Biotechnological Applications*. Marthin A.M. (ed) Chapman and Hall, London pp. 201-217.

Saliu, J. K. (2008). Effect of smoking and frozen storage on the nutrient composition of some African fish. *Adv. Nat. Appl. Science*, 2(1): 16-20.

SPSS (Statistical package for Social Sciences) (1999). Computer Program, MS for Windows. *SPSS 10 for Windows*, Chicago, Illinois, USA.

Stolyhiro, A. and Sikorski, Z. E. (2005). Polycyclic aromatic hydrocarbons in smoked fish: A critical review. *Food Chem.*; 91:303-311.

Sveinsdottir, K. (1998). *The process of fish smoking and quality evaluation*. M. Sc. Dissertation, University of Denmark, Denmark.

Swastawati, F. (2004). The Effect of smoking duration on the quality and DHA composition of milkfish (*Chanos chanos* F). *Journal of Coastal Development*, Volume 7(3):137-142.

Teutscher, F. (1990). The Present and Potential role of small Inland Species on food supply and income generation in Eastern Southern Africa. In: *CIFA Tech. Pap. No. 19*, Rome, FAO 1992, pp. 18-21.

Tobor, J. G. (2004). A review of the fish industry in Nigeria and status of fish preservation method and future growth prerequisite to cope with anticipated increase in production. *NIOMR Technical Paper*, Nigerian Institute for Oceanography and Marine Research, Nigeria.

Trease, G. E. and Evans, W. C. (1983). *Textbook of pharmacognosy*. 12th Ed. Balliere Tindall and Company, pp: 343-383.

Waterman, A. (1976). The Production of Dried Fish. *FAO Fish. Tech. Paper*, 160:115-120.

Watts, D. L. (2010). HTHM mineral ratios-A brief discussion of their clinical importance. *Trace elements newsletters*, Volume 21 (1): 1-3.

Yanar, Y. (2007). Quality changes of hot smoked catfish (*Clarias gariepinus*) during refrigerated storage. *J. Muscle Foods*, 18:391-400.