



Assessment of Organophosphorus and Pyrethroid Pesticide Residues in *Lactuca Sativa* L. and *Solanum Macrocarpum* L. Cultivated in Benin

By Prudence Agnandji, Lucie Ayi-Fanou, Magloire Acakpo Nonvignon Gbaguidi, Boris Fresnel Cachon, Mathieu Hounha, Micheline Agassounon Djikpo Tchibozo, Winfred Seth Kofi Gbewonyo, Fabrice Cazier & Ambaliou Sanni

Université d'Abomey-Calavi

Abstract- Pesticides are harmful and toxic in nature. They are used for vegetable protection against pest attack. The present study aimed to determine the level of Organophosphorus and Pyrethroid residues in vegetables (*Solanum macrocarpum* L. and *Lactuca sativa* L.) Thirty one vegetable samples (16 samples of *Solanum macrocarpum* L. and 15 samples of *Lactuca sativa* L.) from four vegetable farms were collected. The samples were cleaned up before the extraction was performed. Analysis was then processed using the QuEChERS method. Residue analysis was performed using a GC-PFPD for Organophosphorus residues and GC-ECD for Pyrethroid residues. MedCalc Statistical Software version 15.0. was used for the statistical analysis. The results revealed that most of the samples are contaminated by Pyrethroid and Organophosphorus residues.

Keywords: levels contamination, organophosphorus, pyrethrinoid, vegetable crops, south of benin.

GJSFR-C Classification: FOR Code: 279999p



Strictly as per the compliance and regulations of:



Assessment of Organophosphorus and Pyrethroid Pesticide Residues in *Lactuca Sativa L.* and *Solanum Macrocarpum L.* Cultivated in Benin

Prudence Agnandji ^a, Lucie Ayi-Fanou ^a, Magloire Acakpo Nonvignon Gbaguidi ^b, Boris Fresnel Cachon ^c, Mathieu Hounha ^y, Micheline Agassounon Djikpo Tchibozo [§], Winfred Seth Kofi Gbewonyo ^x, Fabrice Cazier ^v & Ambaliou Sanni ^e

Abstract- Pesticides are harmful and toxic in nature. They are used for vegetable protection against pest attack. The present study aimed to determine the level of Organophosphorus and Pyrethroid residues in vegetables (*Solanum macrocarpum L.* and *Lactuca sativa L.*) Thirty one vegetable samples (16 samples of *Solanum macrocarpum L.* and 15 samples of *Lactuca sativa L.*) from four vegetable farms were collected. The samples were cleaned up before the extraction was performed. Analysis was then processed using the QuEChERS method. Residue analysis was performed using a GC-PFPD for Organophosphorus residues and GC-ECD for Pyrethroid residues. MedCalc Statistical Software version 15.0. was used for the statistical analysis. The results revealed that most of the samples are contaminated by Pyrethroid and Organophosphorus residues. Among Pyrethroid residues detected, 6.5% of cypermethrin and 43.75% of lambda cyhalothrin levels in *Solanum macrocarpum L.* have exceeded the maximum residue limits while in *Lactuca sativa L.* 40% of lambda cyhalothrin and 20% of fenvalerate levels exceeded also the maximum residue limits. For Organophosphorus analyzed, 6.25% of diazinon, 12.5% of chlorgenvinip, 18.75% of fenitrothion, 25% of dimethoate and pirimiphos methyl, 50% of profenofos, 56.25% of malathion and 75% of chlorpyrifos levels in *Solanum macrocarpum L.* exceeded maximum residue limits while 13.33 % of ethopropos and chlorgenvinip, 20% of profenofos and fenitrothion, 26.67 % of methamidophos, dimethoate and pirimiphos-methyl, 46.67%

Author a, b, c, d, e, f, g, h: Unité de Biochimie et Biologie Moléculaire, Laboratoire de Biologie Moléculaire et Environnement, Département de Biochimie et Biologie Cellulaire, Université d'Abomey-Calavi 04BP0320 Cotonou, Bénin. e-mail: afaluc@yahoo.fr

Author p: Unité de Recherche en Ecotoxicologie et Etude de Qualité (UREEQ), Laboratoire d'Etude et de Recherche en Chimie Appliquée (LERCA), Ecole Polytechnique d'Abomey-Calavi (EPAC), Université d'Abomey-Calavi (UAC) 01BP2009 Cotonou, Bénin.

Author o: Faculté des Sciences et Techniques de Dassa-Zoume, Université des Sciences, Technologie, Ingénierie et Mathématiques, Dassa-Zoume, Bénin.

Author s: Laboratoire des Normes et de Contrôle de Qualité Microbiologique, Nutritionnelle et Pharmacologique, Université d'Abomey-Calavi, 01 BP 1636 RP, Cotonou, Bénin.

Author x: Department of Biochemistry, Cell and Molecular Biology, School of Biological Sciences, College of Basic and Applied Sciences, University of Ghana, P.O. Box LG54, Legon.

Author v: Centre Commun de Mesures, Université du Littoral Côte d'Opale Dunkerque, France.

of malathion and 53.33% of chlorpyrifos levels in *Lactuca sativa L.* were exceeded the maximum residue limits. No significant difference was observed between the level of Pyrethroid and Organophosphorus residues which are above maximum residue limits in *Solanum macrocarpum L.* and *Lactuca sativa L.* The levels of Pyrethroid and Organophosphorus detected in vegetable samples show that the sanitary quality of these vegetables is affected.

Keywords: levels contamination, organophosphorus, pyrethrinoid, vegetable crops, south of benin.

Cite this Article: Prudence Agnandji, Lucie Ayi-Fanou, Magloire Acakpo Nonvignon Gbaguidi, Boris Fresnel Cachon, Mathieu Hounha, Micheline Agassounon Djikpo Tchibozo, Winfred Seth Kofi Gbewonyo, Fabrice Cazier, and Ambaliou Sanni, "Assessment of Organophosphorus and Pyrethroid Pesticide Residues in *Lactuca sativa L.* and *Solanum macrocarpum L.* cultivated in Benin." *World Journal of Analytical Chemistry*, vol. 6, no. 1 (2018): 4-12. doi: 10.12691/wjac-6-1-2.

I. INTRODUCTION

In Benin, vegetable farming is intensified in southern department's town such as Cotonou and Seme-Kpodji. Vegetables grown during this activity are most often vulnerable to attack by pests (weeds, bacteria, fungi and insect pests, etc.) [1]. A study carried out by Azandeme-Hounmalon *et al.* [2] in Southern Benin, on a population of mite fauna associated with local vegetables (*Amaranthus cruentus L.*, *Solanum macrocarpum L.*) showed that several species of mites attack these vegetables. One of them is *Tetranychus evansi*. The consequences of these attacks are the recording of losses in the harvest yield and the decline in monetary income. To remediate these problems, vegetables farmers use pesticides [3,4,5,6]. There are several groups of pesticides including insecticides such as Pyrethroids (Pyr) and Organophosphorus (Op). They are the insecticides which replaced Organochlorine insecticides [7].

In southern of Benin, active ingredients of synthetic Pyr and Op frequently used in vegetable farming are cypermethrin, lambda cyhalothrin, deltamethrin, profenofos, chlorpyfos, malathion [8]. These active substances have a protective role for vegetable crops against pests such as *aphids*, *beetles* and *lepidoptera* [9]. Pesticides are widely used (in fruit and vegetables) because of their susceptibility to insect and diseases attack. Consequently, food safety is a major public concern worldwide. During the last decades, the increasing demand for food safety has stimulated research regarding the risk associated with consumption of fruits and vegetables as they constitute to many part of human diet, contributing nutrients and vitamins. Therefore, residues of pesticides could affect the consumers, especially when these commodities are freshly consumed. The total dietary intake of pesticides residues that remain on agricultural commodities are known as carcinogens/or toxins and therefore it is desirable to reduce these residues [10]. Despite the fact that Pyr and Op pesticides are used by farmers in Benin to protect their vegetables, no data exist about Pyr and Op contamination levels in vegetables grown in Benin. The present study was designed to evaluate the level of contamination of Op and Pyr residues in exotic and local vegetable crops (*Solanum macrocarpum* L. and *Lactuca sativa* L.) cultivated in southern Benin.

II. MATERIAL AND METHODS

a) Study Areas

Vegetable samples were collected in 2017, during March to July in two farms sites located in Cotonou (intra-urban area) and Seme-kpodji (Suburban area). The map of this study was described in a previous publication [11].

b) Sampling

A total of 31 vegetable samples were collected from the study areas including 15 samples of *Lactuca sativa* L. and 16 *Solanum macrocarpum* L. These vegetables were chosen because they are very common in Beninese diet. The sampling plan consisted of delineating a 10 x 10 m plot in the middle of the field of each farmer. Five primary samples were chosen along the diagonals. These samples were roughly cut and mixed in situ to obtain a final sample of 750 g. Each sample was packed in aluminium foil, plastic bag and ice box. The samples were sent to the laboratory and it was stored at -20°C until the moment of the analysis.

c) Reagents and Glassware

All glassware used for extraction and cleaning was rigorously washed with tap water and detergent and rinsed twice with distilled water. They were again rinsed with acetone and finally dried in an oven. All solvents and reagents used were of the analytical grade supplied by VWR Chemical. Individual pesticide standards were

available at Ghana Standard Authority (GSA) laboratories in Agra.

d) Extraction and Clean-up

Vegetable sample (15 ± 0.1 g) was weighed and milled in a mill (MOULINEX Soup & Co Lm 901b1 Blender) before being transferred to a 50 mL centrifuge tube containing 15 mL of acetonitrile (ACN) HPLC grade. The tube was closed and shaken vigorously on a shaker (STUART SA7, VITESSE FIXE) for 1 min. After shaking, a mixture of 4 g magnesium sulphate, 1 g sodium chloride, 1 g trisodium citrate dihydrate and 0.5 g disodium hydrogenocitrate sesquihydrate was added. The tube was closed and immediately shaken vigorously using a shaker for 1 min and centrifuged for 10 min at 3000 rpm. An aliquot (6 mL) of the extract was transferred into centrifugation tube containing the following sorbents: 150 mg of anhydrous magnesium sulphate, 50 mg of PSA, 50 mg of black graphite Carbon and 50 mg of adsorbent C18 for purification. We added formic acid at 10 μ L per mL to the tube of extract (1%) then the tube was shaken vigorously for 1 min and centrifuged for 10 min at 3000 rpm. Thereafter, 4 mL of the extract supernatant was concentrated with a dessicator containing silica gel activated by heating in oven at 105 °C connected with a pump LABCONCO 117 (A65312906) N038-500. The residue was collected with 1 mL of the mixture of formic acid in ACN (1%) and introduced in vial.

e) Analysis of Insecticide Residues Using Gas Chromatography (GC) Method and Quality Control

A Varian CP3800 gas chromatograph Electron Capture Detector (ECD) and Pulsed Flame Photometric Detector (PFPD) were used to detect and quantify respectively Pyr and Op residues, in the vegetable samples from different sites. For Pyr, samples were injected into the column of the GC with an autosampler at a temperature of 270°C. The compounds were partitioned through the stationary phase, a capillary column (30 m + 10 m EZ Guard, internal diameter (i.d.) 0.25 mm, fused with silica coating VF- 5 ms, 0.25 μ m film) carried by solvents. The oven temperature was programmed at 70°C for 2 min, then 25°C/min till 180°C followed by 5°C/min up to 300°C. The carrier gas was nitrogen maintained at a constant flow rate of 1 ml/min. However, for the Op, the samples were injected into the column of the GC under the same conditions described in previously. The compounds were partitioned through the stationary phase, a capillary column (30 m x 0.25 mm internal diameter (i.d.) fused with silica capillary coating with VF- 1701 ms, 0.25 μ m film) carried by solvents. The oven temperature was programmed at 70°C for 2 min, then 25°C/min till 200°C followed by 20°C/min up to 250°C. The carrier gases were nitrogen (1 ml/min), Air 1 (17 ml/min), Hydrogen (14 ml/min) and Air 2 (10 ml/min).

The molecules of the insecticide residues were differentiated at different rates through the gas. The ECD and PFPD was used to detect differently Pyr and Op compound present. The detection temperatures were 300°C for the ECD and 280°C for the PFPD. These detectors were linked to a computerized integrated system which counts and records the signals as peaks, which were used to quantify insecticide residues present.

The residue of pesticide was identified based on comparison of the measured relative retention times to those of known standards, indicated in Table 1. The residue levels of Pyr or Op pesticides were quantitatively determined by the external standard method using peak area. Measurement was carried out within the linear range of the detector. The peak areas whose retention times coincide with the standards were extrapolated on their corresponding calibration curves to obtain the concentration.

The quality was assured through the analysis of solvent blanks, procedure blanks. The method was optimized and validated using spiked (together) with the internal standard to evaluate the recovery of compounds. The recoveries of internal standards ranged between 71.4% and 110.9% for all the Op and Pyr pesticide. The limit of detection was 10 ppb for Pyr and Op pesticides.

Table 1: Retention times of the pesticide residues

| Pesticides residues | Retention times (min) |
|----------------------------------|-----------------------|
| Organophosphorus Residues | |
| Methamidophos | 7.14 |
| Ethoprofos | 8.86 |
| Diazinon | 9.61 |
| Fonofos | 9.87 |
| Dimethoate | 10.46 |
| Pirimiphos-methyl | 10.65 |
| Chlorpyrifos | 10.87 |
| Malathion | 11.11 |
| Fenitrothion | 11.28 |
| Parathion-ethyl | 11.56 |
| Chlorfenvinphos | 11.76 |
| Profenofos | 12.56 |
| Pyrethroid residues | |
| Allethrin | 16.52 |
| Bifenthrin | 22.70 |
| Fenpropathrin | 23.07 |
| Lambda Cyhalothrin | 24.62 |
| Permethrin | 26.38 |
| Cyfluthrin | 27.58 |
| Cypermethrin | 28.17 |
| Fenvalerate | 29.91 |
| Deltamethrin | 31.18 |

f) Statistical Analysis

Results obtained are expressed in mean \pm standard deviation of the mean. Statistical analysis was performed by Student test unmatched to compare samples mean. The second test was Chi-square and it was used to determine the difference between the number of vegetable samples which the level in Pyr and Op residues are above the Maximum Residue Limits (MRLs). The statistical significance for each analysis was considered at $P < 0.05$. MedCalc Statistical Software version 15.0. was used for the statistical analysis.

III. RESULTS

a) Level of Pyrethroids Pesticides Concentration in *Solanum macrocarpum L.* and *Lactuca sativa L.*

Vegetables such as *Solanum macrocarpum L.* and *Lactuca sativa L.* are an important source of nutrition for human system and human diet. The contamination of those vegetables by pesticides could cause a threat to human health. Many studies about monitoring of Pyr pesticides shown their presence in vegetables. In our study we researched nine Pyr molecules in vegetables (*Solanum macrocarpum L.* and *Lactuca sativa L.*). The mean total concentration of Pyr residues (0.076 ± 0.248 mg/kg) found in *Solanum macrocarpum L.* is higher than the mean total concentration (0.070 ± 0.280 mg/kg) found in *Lactuca sativa L.*, but the difference is not statistically significant ($p > 0.05$).

b) Range, mean \pm SD of Pyrethroids Found in *Solanum macrocarpum L.* and *Lactuca sativa L.*

Table 2 shows minimum, maximum, mean \pm standard deviation of Pyr found in *Solanum macrocarpum L.* and *Lactuca sativa L.*

High concentration (0.045 mg/kg) of allethrin was detected in *Solanum macrocarpum L.* samples with mean of $88 \times 10^{-3} \pm 11.5 \times 10^{-3}$ mg/kg.

Bifenthrin and fenpropathrin weren't detected in *Lactuca sativa L.* samples but in *Solanum macrocarpum L.* their concentration reached 0.02 and 0.007 mg/kg. Their averages are $2.50 \times 10^{-3} \pm 5.45 \times 10^{-3}$ and $0.56 \times 10^{-3} \pm 1.79 \times 10^{-3}$ mg/kg.

The highest concentration of lambda cyhalothrin found in *Solanum macrocarpum L.* was 2.13 mg/kg with an average of $523.7 \times 10^{-3} \pm 564.1 \times 10^{-3}$ mg/kg. Residues of permethrin and fenvalerate were detected in *Lactuca sativa L.* at maximum concentration levels of 0.027 and 0.049 mg/kg respectively, with averages of $2.33 \times 10^{-3} \pm 6.96 \times 10^{-3}$ and $15.4 \times 10^{-3} \pm 14.6 \times 10^{-3}$ mg/kg. The highest concentration of cyfluthrin found in *Solanum macrocarpum L.* is 0.098 mg/kg with an average of $21.1 \times 10^{-3} \pm 22.3 \times 10^{-3}$ mg/kg. That of cypermethrin found in *Solanum macrocarpum L.* is 1.025 mg/kg with an average of $186.3 \times 10^{-3} \pm 260.3 \times 10^{-3}$ mg/kg. Similarly, the maximum level of deltamethrin residues found in *Solanum macrocarpum L.* is 0.028



mg/kg with an average of $11.9 \times 10^{-3} \pm 8.55 \times 10^{-3}$ mg/kg.

c) Frequency of Occurrence of Pyrethroid Residues in Samples of *Solanum macrocarpum L.* and *Lactuca sativa L.*

Figure 1 shows the frequencies of appearance of Pyr residues in the vegetable samples analyzed. From the analysis of the data in Figure 1 it is noted that the residues of bifenthrin and fenpropathrin were not detected in the samples of *Lactuca sativa L.* On the other hand, these residues are found at frequencies of 25% and 12.5% respectively in *Solanum macrocarpum L.* samples. The residues of permethrin and allelthrin are the least predominant in the two varieties of vegetables. They were detected at proportions of 6.25% and 25%, respectively, in the samples of *Solanum macrocarpum L.* While in *Lactuca sativa L.* samples, these proportions were 26.67% and 33.33%.

Deltamethrin residues were found in proportions of 60% and 93.75%, respectively, in the *Lactuca sativa L.* and *Solanum macrocarpum L.* samples. Fenvalerate was found to be 86.67% in the *Lactuca sativa L.*

samples. L. and 56.25% in samples of *Solanum macrocarpum L.* Cypermethrin, cyfluthrin and lambda cyhalothrin were found in all samples of *Solanum macrocarpum L.* and *Lactuca sativa L.* at frequencies greater than 73.33 %.

d) Maximum Residue Limit of Pyrethroids residues in *Solanum macrocarpum L.* and *Lactuca sativa L.*

Table 3 presents the Maximal Residue Level (MRL) and the percentage of the residues which the level are exceeded the limit.

Among Pyr residues analyzed, only three residues levels are above EU MRLs of European Union [12]. These residues are lambda cyhalothrin and cypermethrin in *Solanum macrocarpum L.*; lambda cyhalothrin and fenvalerate in *Lactuca sativa L.* Among Pyr residues which the concentration levels were above MRLs, only lambda cyhalothrin concentration was detected in 7/16 case (43.75%) in *Solanum macrocarpum L.* and 7/15 case (40%) in *Lactuca sativa L.* Fenvalerate concentration found in *Lactuca sativa L.* was above MRLs in 3/15 case (20%).

Table 2: Pyrethroids Pesticides residues concentrations in the tested vegetables (mg/kg)

| Pyrethroids Residues | <i>Solanum macrocarpum L.</i> | | <i>Lactuca Sativa L.</i> | |
|----------------------|-------------------------------|---|--------------------------|--|
| | Range | Mean \pm SD | Range | Mean \pm SD |
| Allelthrin | ND-0.046 | $2.88 \times 10^{-3} \pm 11.5 \times 10^{-3}$ | 0.003-0.011 | $1.80 \times 10^{-3} \pm 3.17 \times 10^{-3}$ |
| Bifenthrin | 0.001-0.020 | $2.50 \times 10^{-3} \pm 5.45 \times 10^{-3}$ | ND | ND |
| Fenpropathrin | 0.002-0.007 | $0.56 \times 10^{-3} \pm 1.79 \times 10^{-3}$ | ND | ND |
| Lambda Cyhalothrin | 0.067-2.127 | $523.7 \times 10^{-3} \pm 564.1 \times 10^{-3}$ | 0.017-1.713 | $675.9 \times 10^{-3} \pm 6.73 \times 10^{-3}$ |
| Permethrin | 0.003-0.014 | $1.94 \times 10^{-3} \pm 4.10 \times 10^{-3}$ | 0.001-0.027 | $2.33 \times 10^{-3} \pm 6.96 \times 10^{-3}$ |
| Cyfluthrin | 0.002-0.098 | $21.1 \times 10^{-3} \pm 22.3 \times 10^{-3}$ | 0.001-0.035 | $8.73 \times 10^{-3} \pm 10 \times 10^{-3}$ |
| Cypermethrin | 0.003-1.025 | $186.3 \times 10^{-3} \pm 260.3 \times 10^{-3}$ | 0.003-0.021 | $45.8 \times 10^{-3} \pm 103.5 \times 10^{-3}$ |
| Fenvalerate | 0.001-0.018 | $3.75 \times 10^{-3} \pm 5.08 \times 10^{-3}$ | 0.003-0.049 | $15.4 \times 10^{-3} \pm 14.6 \times 10^{-3}$ |
| Deltamethrin | 0.003-0.028 | $11.9 \times 10^{-3} \pm 8.55 \times 10^{-3}$ | 0.002-0.011 | $3.27 \times 10^{-3} \pm 3.69 \times 10^{-3}$ |




Figure 1: Frequency of Pyrethroid residues in *Solanum macrocarpum L.* and *Lactuca sativa L.*

Table 3: Maximum residue limits (MRLs) of Pyrethroids in vegetables samples and percentage of its above MRLs

| Pyrethroids residues | <i>Solanum macrocarpum L.</i> | | <i>Lactuca sativa L.</i> | | χ^2 | <i>p</i> -value |
|----------------------|-------------------------------|---|--------------------------|--|----------|-----------------|
| | EU MRL mg/kg | % of <i>Solanum macrocarpum L.</i> > EU MRL | EU MRL mg/kg | % of <i>Lactuca sativa L.</i> > EU MRL | | |
| Allethrin | -- | -- | -- | -- | -- | -- |
| Bifenthrin | -- | -- | 0.01 | 0 | -- | -- |
| Fenpropathrin | 0.01 | 0 | 0.5 | 0 | -- | -- |
| Lambda Cyhalothrin | 0.2 | 43.75 | 0.5 | 40 | 0.023 | 0.879 |
| Permethrin | 0.05 | 0 | 0.05 | 0 | -- | -- |
| Cyfluthrin | -- | -- | 1 | 0 | -- | -- |
| Cypermethrin | 0.7 | 6.25 | 2 | 0 | 0.00108 | 0.974 |
| Fenvalerate | 0.02 | 0 | 0.02 | 20 | 1.624 | 0.202 |
| Deltamethrin | 0.2 | 0 | 0.5 | 0 | -- | -- |

-- = No determined.

e) Level of Organophosphorus Pesticides in *Solanum macrocarpum L.* and *Lactuca sativa L.*

As previously reported, the mean total concentration (0.065 ± 0.360 mg/kg) of Op found in *Solanum macrocarpum L.* is higher than the mean total concentration (0.025 ± 0.106 mg/kg) found in *Lactuca sativa L.*, as indicated in Figure 2. But the difference is not statistically significant ($p > 0.05$).

f) Range, mean \pm SD of Organophosphorus Found in *Solanum macrocarpum L.* and *Lactuca sativa L.*

Table 4 show minimum, maximum, mean \pm standard deviation of Op found in *Solanum macrocarpum L.* and *Lactuca sativa L.*

For the Op, contamination levels ranged from non-determined (ND) to 4.809 mg/kg in the samples of *Solanum macrocarpum L.* and ND at 1.095 mg/kg in the

Lactuca sativa L samples. These high concentrations were recorded for the residues of dimethoate and chlorpyrifos, respectively. The average residues of Op vary from $0.62 \times 10^{-3} \pm 2.50 \times 10^{-3}$ mg/kg for fonofos at $1386 \times 10^{-3} \pm 178.9 \times 10^{-3}$ mg/kg for Malathion in *Solanum macrocarpum L.* On the other hand, in *Lactuca sativa L.* These mean concentrations per residue vary from $0.78 \times 10^{-3} \pm 2.94 \times 10^{-3}$ mg/kg for the diazinon at $203.6 \times 10^{-3} \pm 305.4 \times 10^{-3}$ mg/kg for chlorpyrifos.

In *Solanum macrocarpum L.* the two Op residues were not detected in all samples. These are diazinon and fonofos. In contrast, in *Lactuca sativa L.* residues of diazinon, ethopropofos, and parathion-ethyl were not detected in all samples. This finding shows that several Op molecules at varying doses are used on sites to control crop pests.




Figure 2: Mean of total concentration of Organophosphorus in *Solanum macrocarpum L.* and *Lactuca sativa L.*

Table 4: Pesticides residues concentrations in the tested vegetables (mg/kg)

| Organophosphorus Residues | <i>Solanum macrocarpum L.</i> | | <i>Lactuca sativa L.</i> | |
|---------------------------|-------------------------------|---|--------------------------|---|
| | Range (mg/kg) | Mean \pm SD (mg/kg) | Range (mg/kg) | Mean \pm SD (mg/kg) |
| Methamidophos | 0.011-0.014 | $2.25 \times 10^{-3} \pm 4.88 \times 10^{-3}$ | 0.008-0.016 | $5.50 \times 10^{-3} \pm 6.05 \times 10^{-3}$ |
| Ethopropos | 0.006-0.011 | $2.06 \times 10^{-3} \pm 3.82 \times 10^{-3}$ | ND-0.007 | $10^{-3} \pm 2.54 \times 10^{-3}$ |
| Diazinon | ND-0.232 | $14.5 \times 10^{-3} \pm 58 \times 10^{-3}$ | ND-0.011 | $0.78 \times 10^{-3} \pm 2.94 \times 10^{-3}$ |
| Fonofos | ND-0.01 | $0.62 \times 10^{-3} \pm 2.50 \times 10^{-3}$ | 0.009-0.012 | $2.86 \times 10^{-3} \pm 4.74 \times 10^{-3}$ |
| Dimethoate | 0.026-4.809 | $310 \times 10^{-3} \pm 1999.9 \times 10^{-3}$ | 0.025-0.034 | $8.27 \times 10^{-3} \pm 13.7 \times 10^{-3}$ |
| Pirimiphos-methyl | 0.009-0.028 | $6.94 \times 10^{-3} \pm 10 \times 10^{-3}$ | 0.008-0.014 | $5.15 \times 10^{-3} \pm 5.98 \times 10^{-3}$ |
| Chlorpyrifos | 0.015-0.704 | $219.8 \times 10^{-3} \pm 188.2 \times 10^{-3}$ | 0.025-1.095 | $203.6 \times 10^{-3} \pm 305.4 \times 10^{-3}$ |
| Malathion | 0.011-0.557 | $1386 \times 10^{-3} \pm 178.9 \times 10^{-3}$ | 0.017-0.266 | $34 \times 10^{-3} \pm 68.7 \times 10^{-3}$ |
| Fenitrothion | 0.021-0.03 | $4.81 \times 10^{-3} \pm 10.5 \times 10^{-3}$ | 0.023-0.025 | $5.14 \times 10^{-3} \pm 10.2 \times 10^{-3}$ |
| Parathion-ethyl | 0.01-0.015 | $1.56 \times 10^{-3} \pm 4.37 \times 10^{-3}$ | ND | -- |
| Chlorfenvinphos | 0.017-0.02 | $2.31 \times 10^{-3} \pm 6.34 \times 10^{-3}$ | 0.014-0.017 | $2.21 \times 10^{-3} \pm 5.66 \times 10^{-3}$ |
| Profenofos | 0.006-0.353 | $77.6 \times 10^{-3} \pm 109.9 \times 10^{-3}$ | 0.007-0.329 | $27.3 \times 10^{-3} \pm 84.3 \times 10^{-3}$ |

ND or -- = No determined.




Figure 3: Frequency of organophosphorus in *Solanum macrocarpum L.* and *Lactuca sativa L.*

g) Frequency of Organophosphorus in *Solanum macrocarpum L.* and *Lactuca sativa L.*

A total of twelve Op molecules were searched in two varieties of vegetables as *Solanum macrocarpum L.* and *Lactuca sativa L.* Among the Op residues detected in the vegetable samples, parathion ethyl was not found in *Lactuca sativa L.* but it was found in low proportion in *Solanum macrocarpum L.* (12.5%). Similarly residues such as methamidophos, ethopros, diazinon, dimethoate, fenitrothion, chlofenvinip are also found in proportions of less than 30% in all the studied vegetables. On the other hand, residues of chlorpyrifos, malathion and profenofos are the most ubiquitous with proportions varying between 53 and 100%, as indicated in Figure 3.

h) Maximum Residue Limits of Organophosphorus Residues in Vegetables and Percentage of Its above MRLs

Table 5 presents the MRLs and the percentage of the Op residues which the level are exceeded the limits.

Among Op residues analyzed in *Solanum macrocarpum L.* only three residues level were below EU MRLs [13]. These residues are methamidophos, ethopros and parathion ethyl. The concentration of the rest was found at the level above EU MRLs at different proportion; diazinon (6.25%), chlofenvinip (12.5%), fenitrothion (18.75%), dimethoate, (25%), pirimiphos methyl (25%), profenofos (50%) malathion (56.25%) and chlorpyrifos (75%).

Table 5: Maximum residue limits (MRLs) of Organophosphorus residues in vegetables and percentage of its above MRLs

| | <i>Solanum macrocarpum L.</i> | | <i>Lactuca sativa L.</i> | | χ^2 | <i>p</i> -value |
|-------------------|-------------------------------|---|--------------------------|--|----------|-----------------|
| | EU MRL | % of <i>Solanum macrocarpum L.</i> > EU MRL | EU MRL | % of <i>Lactuca sativa L.</i> > EU MRL | | |
| Methamidophos | 0.01 | 0 | 0.01 | 26.67 | 2.813 | 0.093 |
| Ethoprophos | 0.02 | 0 | 0.02 | 13.33 | 0.606 | 0.436 |
| Diazinon | 0.01 | 6.25 | 0.01 | 0 | 0.00108 | 0.974 |
| Fonofos | -- | -- | -- | -- | -- | -- |
| Dimethoate | 0.02 | 25 | 0.02 | 26.67 | 0.093 | 0.761 |
| Pirimiphos-methyl | 0.01 | 25 | 0.01 | 26.67 | 0.093 | 0.761 |
| Chlorpyrifos | 0.05 | 75 | 0.05 | 53.33 | 0.782 | 0.376 |
| Malathion | 0.02 | 56.25 | 0.02 | 46.67 | 0.030 | 0.862 |
| Fenitrothion | 0.01 | 18.75 | 0.01 | 20 | 0.135 | 0.714 |
| Parathion-ethyl | 0.05 | 0 | 0.05 | 0 | -- | -- |
| Chlorfenvinphos | 0.01 | 12.5 | 0.01 | 13.33 | 0.218 | 0.641 |
| Profenofos | 0.01 | 50 | 0.01 | 20 | 1.874 | 0.171 |

-- = no limit has fixed By EU.

For Op residues analyzed in *Lactuca sativa L.*, only two residues level were below MRLs of EU [13]. These residues are diazinon and parathion ethyl. The concentration of the rest were found at the level above EU MRLs in different proportion; ethoprophos and chlofenvinphos (13.33%); fenitrothion and profenofos (20%); dimethoate, pirimiphos methyl and methamidophos (26.67%); malathion (46.67%) and chlorpyrifos (53.33%).

IV. DISCUSSION

The investigation about Op and Pyr pesticides in two vegetables (*Solanum macrocarpum L.* and *Lactuca sativa L.*) grown in southern Benin, revealed that all the Op pesticides studied were found in *Solanum macrocarpum L.* leaves.

On the other hand, parathion-ethyl wasn't detected in *Lactuca sativa L.* The levels of Op residues ranged from no determined (ND) to 4.81 mg/kg in *Solanum macrocarpum L.* with mean of 0.065 ± 0.360 mg/kg. In *Lactuca sativa L.* the highest level detected is 1.1 ppm with mean of 0.025 ± 0.106 mg/kg. The Op residues are methamidophos, ethoprophos, diazinon, fonofos, dimethoate, pirimiphos-methyl, fenitrothion, parathion-ethyl, chlofenvinphos, malathion, chlorpyrifos and profenofos. Comparing our results with other studies, we note that diazinon was detected in vegetables in the range of 0.066 to 1.84 mg/kg [13].

These levels of contamination are superior to our finding which the ranges for this residue is ND to 0.232 mg/kg in *solanum macrocarpum L.* and ND to 0.011 mg/kg in *Lactuca sativa L.* [13]. For these two studies, the levels of diazinon in vegetables exceed the EU MRLs of 0.01 mg/kg. Quintero *et al.* [14] also found similar results in vegetables. Contrary to our results,

chlorpyrifos was found in Thailand at a concentration of 0.008 mg/kg in *Lactuca sativa L.* [15]. This level of contamination is lower than the EU MRLs which is still 0.05 mg/kg. These findings show that Op pesticides are widely used in market gardening in the world. In the present study, chlorpyrifos, malathion and profenofos were detected at a high frequency with levels above the EU MRLs (Table 5). Our results corroborate with of Wanwimolruk *et al.* [16] who found in 34 samples of Chinese cabbage tested, chlorpyrifos (7.2-37.700 ppb), dimethoate (0.5-239 ppb) and profenofos (0.5-183.000 ppb). In this study, the proportion of samples with contamination levels above the MRLs is 29% [16]. In addition, the analysis of cauliflower samples revealed the presence of Op residues such as chlorpyrifos and profenofos. However, none of the samples contained residues above their respective MRLs [17]. The analysis of tomato from Ghana revealed the presence of malathion in 50% of the samples at an average concentration of 0.027 ± 0.021 mg/kg while dimethoate was found in samples at a rate of 15%. Its average concentration in the samples was 0.155 ± 0.113 mg/kg [18]. Malathion and dimethoate levels in the tomato samples were higher than the EU MRLs [18]. These levels of contamination, which are sometimes found below the MRLs, are due to the washing of the leaves of the vegetables concerned either during watering or during rain season and do not contradict the massive use of Op.

For Pyr, we studied allethrin, bifenthrin, fenpropathrin, lamda cyhalothrin, permethrin, cyfluthrin, cypermethrin, fenvalerate, and deltamethrin residues. The average concentrations found are respectively 0.076 ± 0.248 mg/kg in *Solanum macrocarpum L.* and 0.070 ± 0.280 mg/kg in *Lactuca sativa L.* Allethrin has

no MRLs for the vegetable samples studied. Among bifenthrin, fenpropathrin and permethrin which have EU MRLs for the samples studied, they were detected in low frequency, without exceeding above EU MRLs. Akomea-Frempong *et al.* [13] also detected bifenthrin in vegetable (lettuce and cabbage) samples at an average concentration of 0.034 mg/kg. Bempah *et al.* [19] found permethrin in lettuce at a concentration range of 0.011 to 0.035 mg/kg with an average of 0.030 ± 0.009 mg/kg. This range of concentrations is higher than the level of permethrin we found in our samples of *Lactuca sativa L.* (0.001-0.027 mg/kg). For both studies, contamination levels of *Lactuca sativa L.* samples with permethrin are below the EU MRLs. Our levels of bifenthrin are lower than those found by González-Rodríguez *et al.* [20] who reported bifenthrin concentrations in lettuce of 0.02 to 0.05 mg/kg in Spain. Fenpropathrin was also detected in cucumber and tomato samples from Egypt at concentrations of 0.04 and 0.079 mg/kg, respectively (0.021-0.03 mg/kg) in *Solanum macrocarpum L* and (0.023- 0.025 mg/kg) in *Lactuca sativa L.* for this study [21]. These concentrations of fenpropathrin detected in the cucumber and tomato samples are higher than those we detected in our vegetables [21]. Permethrin, allethrin was found in eggplant in Ghana. Their respective mean concentrations are 0.003 mg/kg for Permethrin and 0.126 ± 0.018 mg/kg for allethrin [18]. In contrast, for the present study, Permethrin was detected in *Solanum macrocarpum L.* samples between 0.003 and 0.014 mg/kg. While in *Lactuca sativa L.* its concentration varies between 0.001 and 0.027 mg/kg. As for allethrin, it was detected between ND and 0.046 mg/kg in *Solanum macrocarpum L.* and 0.003 and 0.011 mg/kg in *Lactuca sativa L.* Permethrin was detected in 35% of the eggplant samples whereas in our case it was detected in 25% of the samples of *Solanum macrocarpum L.* and 27% in *Lactuca sativa L.* allethrin was detected in 100% of the samples [18]. In the present study, however, it is found only at 6.25% in the samples of *Solanum macrocarpum L.* and 33.33% in the samples of *Lactuca sativa L.* In contrast to our results, the residues of permethrin found in eggplant samples in Ghana were lower than the EU MRLs [19]. In contrast, consistent with our results, allethrin residues in the same samples exceeded the MRLs [18].

Our results are confirmed by Diop *et al.* [22], who found lambda cyhalothrin in lettuce from different farms sites of Senegal. The mean of it concentration is 0.222 mg/kg and it range is ND and 1.306 mg/kg. Less than 50% levels of lambda cyhalothrin found in lettuce samples are above EU MRLs.

All these observations show that Op and Pyr are massively used in market gardening in the world for pests control in farming. Their use is not without consequence on the sanitary quality of the vegetables whose contamination levels vary as well according to

the residues and the cultures with contents sometimes exceeding the maximum limits in residues. From the nutritional point of view, the level of Pyr and Op residues found in *Solanum macrocarpum L.* and *Lactuca sativa L.* samples at level above EU MRLs pose public health problem. Exposure to pesticides through a mouth is a major way for the people which aren't farm workers and don't live close to the vegetables fields. Many studies shown the health risk associated with the consumption of vegetables contaminated by pesticide residues [23,24,25,26].

This problem is especially encountered in poor countries by the proliferation of illegal trade in plant protection products, which are sometimes prohibited.

V. CONCLUSION

This study evaluates the contamination levels of Op and Pyr pesticides in vegetable crops in southern Benin revealed their presence at various concentrations showing their massive use in market gardening in southern Benin. The most detected residues are Pyr (lambda cyhalothrin) and Op (chlorpyrifos). By comparison of levels of residues with EU MRLs, lambda cyhalothrin, fenvalerate, diazinon, chlorgenvinp, fenitrothion, dimethoate, pirimiphos methyl, profenofos, malathion and chlorpyrifos were detected in samples of *Solanum macrocarpum L.* and/or *Lactuca sativa L.* at levels above the MRLs. These observations show that the use of pesticides in market gardening affects the sanitary quality of leafy vegetables such as *Solanum macrocarpum L.* and *Lactuca sativa L.* More studies should be done on quantifying these pesticide residues in other plants grown in Benin.

REFERENCES RÉFÉRENCES REFERENCIAS

1. Gregory, P. J., Johnson S. N., Newton A. C., Ingram J. S. I. (2009). Integrating pests and pathogens into the climate change/food security debate. *Journal of Experimental Botany*, 60: 2827-2838.
2. Azandeme-Hounmalon G. Y., Affognon, H. D., Assogba-komlan, F., Tamo, M., Fiaboe, K. K. M., Kreiter, S., Martin, T. (2014). Comportement des maraîchers face à l'invasion de *Tetranychus Evansi Baker & Pritchard* au Sud du Benin. AFPP-Dixième conférence internationale sur les ravageurs en Agriculture Montpellier-22 et 23 Octobre 2014.
3. Adjrah, Y., Dovlo, A., Karou1, S. D., Eklu-Gadegbeku, K., Agbonon, A., de Souza, C., Gbeassor, M. (2013). Survey of pesticide application on vegetables in the Littoral area of Togo. *Annals of Agricultural and Environmental Medicine*, 20(4), 715-720.
4. Gaupp-Berghausen, M., Hofer, M., Rewald, B., Zaller, J. G. (2015). Glyphosate-based herbicides reduce the activity and reproduction of earthworms and lead to increased soil nutrient concentrations. *Scientific Reports* 5: 12886.

5. Zakharov, S., Csomor, J., Urbanek, P., Pelclova, D. (2016). Toxic Epidermal Necrolysis After Exposure to Dithiocarbamate Fungicide Mancozeb. *Basic Clin Pharmacol Toxicol.* 118(1): 87-91.
6. Sun, X. (2015). History and Current Status of Development and Use of Viral Insecticides in China. *Viruses* 7: 306-319.
7. Elliott, M. N.F., Potter, J. C. (1978). The future of Pyrethroids in insect control. *Ann. Rev. Entomol.* 23:443-69.
8. Ahouangninou, C. A., Martin, T., Edorh, P., Siddick, I. A., Bio-Bangana, S., Dion, S., Onil, S. S., Boko M., Simon, S., Fagnomi, B.E. (2012). Characterization of health and environmental risks of pesticide used in market-gardening in the rural city of Tori-Bossito in Benin, west Africa. *J Environ Prot.*
9. Grzywacz, D., Rossbach, A., Rauf, A., Russell, D.A., Srinivasan, R., Shelton, A.M. (2010). Current control methods for diamondback moth and other brassica insect pests and the prospects for improved management with lepidopteran-resistant Bt vegetable brassica in Asia and Africa. *Crop protection* 29: 68-79.
10. Zawiyah, S., Che Man Y.B., Nazimah, S.A.H., Chin, C.K., Tsukamoto, I., Hamanya, A.H., Norhaizan, I. (2007). Determination of organochlorine and Pyrethroid pesticides in fruit and vegetables using SAX/PSA clean-up column. *Food Chem.* 102: 98-103.
11. Agnandji, P., Ayi-Fanou, L., Gbaguidi, A.N.M., Cachon, B.F., Hounha, M., Tchibozo, D.M., Cazier, F., Sanni, A. (2018). Distribution of Organochlorine Pesticides Residues in *Solanum macrocarpum* and *Lactuca sativa* Cultivated in South of Benin (Cotonou and Seme-kpodji). *American Journal of Food Science and Technology.* 6(1): 19-25.
12. EU. "EU-Pesticides Database", 2015.
13. Akomea-Fremppong, S., Ofosu, I. W., Owusu-Ansah E. J., Darko, G. (2017). Health risks due to consumption of pesticides in ready-to-eat vegetables (salads) in Kumasi, Ghana. *International Journal of Food Contamination.* 4:13.
14. Quintero, A., Caselles M J., Ettiene, G., Colmenares, N. G., Rami'rez, T., Deisy, Medina. (2008). Monitoring of Organophosphorus Pesticide Residues in Vegetables of Agricultural Area in Venezuela. *Bull Environ Contam Toxicol.* 81: 393-396.
15. Sapbamrer, R., Hongsibsong, S. (2014). Organophosphorus Pesticide Residues in Vegetables From Farms, Markets, and a Supermarket Around Kwan Phayao Lake of Northern Thailand. *Arch Environ Contam Toxicol* 67: 60-67.
16. Wanwimolruk, S., Kanchanamayoon, Onnicha., Phopin, K., Prachayasittikul, V. (2015). Food safety in Thailand 2: Pesticide residues found in Chinese kale (*Brassica oleracea*), a commonly consumed vegetable in Asian countries. *Science of the Total Environment* 532: 447-455.
17. Mandal, K., Singh, B. (2010). Magnitude and Frequency of Pesticide Residues in Farmgate Samples of Cauliflower in Punjab, India. *Bull Environ Contam Toxicol* 85:423-426.
18. Akoto, O., Gavor, Sandra., Appah, M. K., Apau, J. (2015). Estimation of human health risk associated with the consumption of pesticide-contaminated vegetables from Kumasi, Ghana. *Environ Monit Assess* 187: 244.
19. Bempah, C. K., Buah-Kwofie, A., Denutsui, D., Asomaning, J., Tutu, A. O. (2011). Monitoring of Pesticide Residues in Fruits and Vegetables and Related Health Risk Assessmentin Kumasi Metropolis, Ghana. *Research Journal of Environmental and Earth Sciences* 3(6): 761-771.
20. González-Rodríguez, R. M., Rial-Otero, R., Cancho-Grande, B., Simal-Gándara, J. (2008). Occurrence of fungicide and insecticide residues in trade samples of leafy vegetables. *Food Chem.* 107(3): 1342-7.
21. Ahmed, M. T., Greish, S., Ismail, S. M., Mosleh, Y., Loufty N. M. El Doussouki, A. (2013). Dietary Intake of Pesticides Based on Vegetable Consumption in Ismailia, Egypt. A Case Study» Human and Ecological Risk Assessment: An International Journal.
22. Diop, A., Diop, Y. M., Thiaré, D. D., Cazier, F., Sarr, S. O., Kasprowiak. A, Landy, D., Delattre, F. (2016). Monitoring survey of the use patterns and pesticides residues on vegetables in the Nyayes zone, Senegal. *Chemosphere* 144: 1715-1721.
23. Saha, N., Zaman, M. R. (2012). Evaluation of possible health risks of heavy metals by consumption of foodstuffs available in the central market of Rajshahi City, Bangladesh. *Environmental Monitoring and Assessment.*
24. Shen, L., Xia, B., Dai, X., (2013). Residues of persistent organic pollutant sinfrequently consumed vegetables and assessment of human health risk based on consum ption of vegetables in Huizhou, South China. *Chemosphere* 93: 2254-2263.
25. Arias, L. A., Bojaca, C., Ahumada, A.D., Schrevens, E. (2014). Monitoring of pesticide residues in tomato marketed in Bogota, Colombia. *Food Control* 35:213-217.
26. Yihua, L., Danyu, S., Shiliang, L., Zhanglin, N., Ming, D., Caifen, Y., Fubin, T. (2016). Residue levels and risk assessment of pesticides in nuts of China. *Chemosphere* 144: 645-651.