



GLOBAL JOURNAL OF SCIENCE FRONTIER RESEARCH: F
MATHEMATICS AND DECISION SCIENCES

Volume 20 Issue 6 Version 1.0 Year 2020

Type : Double Blind Peer Reviewed International Research Journal

Publisher: Global Journals

Online ISSN: 2249-4626 & Print ISSN: 0975-5896

New Integrals for Horn Hypergeometric Functions in Two Variables

By Jihad Ahmed Younis

Aden University

Abstract- In this paper, we establish several new integral representations of Euler-type involving Horn's functions $G_1, G_2, G_3, H_1, H_2, H_5, H_6$ and H_7 . Some corollaries have also been obtained as special cases of our main results.

Keywords: *gamma function, beta function, horn double functions, appell functions, eulerian integrals.*

GJSFR-F Classification: MSC 2010: 05B25



Strictly as per the compliance and regulations of:





New Integrals for Horn Hypergeometric Functions in Two Variables

Jihad Ahmed Younis

Abstract- In this paper, we establish several new integral representations of Euler-type involving Horn's functions $G_1, G_2, G_3, H_1, H_2, H_5, H_6$ and H_7 . Some corollaries have also been obtained as special cases of our main results.

Keywords: gamma function, beta function, horn double functions, appell functions, eulerian integrals.

I. INTRODUCTION

Integral representations of hypergeometric functions have found applications in divers fields such as mathematics, physics, statistics, and engineering. Hasanov et al. [9] studied some of the properties of the Horn type second-order double hypergeometric function H_2^* involving integral representations, differential equations, and generating functions. Choi et al. [6] introduced certain integral representations for Srivastava's triple hypergeometric functions H_A, H_B and H_C . Younis and Bin-Saad [19, 20] establish several integral representations and operational relations involving quadruple hypergeometric functions $X_i^{(4)}$ ($i = 38, 40, 45, 48, 50$). Younis and Nisar [21] introduce new integral representations of Euler-type for Exton's hypergeometric functions of four variables D_1, D_2, D_3, D_4 and D_5 . Also, in [2-5], authors introduced many integral representations for certain hypergeometric functions in four variables.

Let us recall the Gauss hypergeometric function ${}_2F_1$ is defined as (see, e.g., [14] and [16, Section 1.5])

$${}_2F_1(a, b; c; x) = \sum_{m=0}^{\infty} \frac{(a)_m (b)_m}{(c)_m} \frac{x^m}{m!}, (|x| < 1), \quad (1.1)$$

where $(a)_m$ is the well known Pochhammer symbol given by (see, e.g., [16, p. 2 and pp. 4-6])

$$(a)_m = \frac{\Gamma(a+m)}{\Gamma(a)} = \begin{cases} 1 & (m=0), \\ a(a+1)\dots(a+m-1) & (m \in \mathbb{N} := \{1, 2, \dots\}). \end{cases} \quad (1.2)$$

Euler's integral representation of ${}_2F_1$ is defined by (see, e.g., [14, p. 85] and [16, p. 65])

Author: Department of Mathematics, Aden University, Yemen.
e-mail: jihadalsaqaf@gmail.com

$${}_2F_1(a, b; c; x) = \frac{\Gamma(c)}{\Gamma(a)\Gamma(c-a)} \int_0^1 \alpha^{a-1} (1-\alpha)^{c-a-1} (1-x\alpha)^{-b} d\alpha,$$

$$(Re(a) > 0, Re(c-a) > 0).$$

Appell hypergeometric functions of two variables F_1, F_2 and F_3 are respectively defined by (see [17, p. 53, Eq. (4) - (6)])

$$F_1(a, b, c; d; x, y) = \sum_{m,n=0}^{\infty} \frac{(a)_{m+n}(b)_m(c)_n}{(d)_{m+n}} \frac{x^m y^n}{m! n!}, \quad (1.3)$$

$$F_2(a, b, c; d, e; x, y) = \sum_{m,n=0}^{\infty} \frac{(a)_{m+n}(b)_m(c)_n}{(d)_m(e)_n} \frac{x^m y^n}{m! n!} \quad (1.4)$$

and

$$F_3(a, b, c, d; e; x, y) = \sum_{m,n=0}^{\infty} \frac{(a)_m(b)_n(c)_m(d)_n}{(e)_{m+n}} \frac{x^m y^n}{m! n!}. \quad (1.5)$$

Integral representations of Euler type for the functions F_1, F_2, F_3 were already given by Appell [1, Chap. III]. For various integral representations of hypergeometric functions, the interested reader may refer to [8-10, 12, 13, 15, 18].

Other hypergeometric functions of two variables are the following Horn's functions $G_1, G_2, G_3, H_1, H_2, H_3, H_4, H_5, H_6$ and H_7 defined by (cf. [7], [8], [11])

$$G_1(a, b, c; x, y) = \sum_{m,n=0}^{\infty} (a)_{m+n}(b)_{n-m}(c)_{m-n} \frac{x^m y^n}{m! n!}, \quad (1.6)$$

$$G_2(a, b, c, d; x, y) = \sum_{m,n=0}^{\infty} (a)_m(b)_n(c)_{n-m}(d)_{m-n} \frac{x^m y^n}{m! n!}, \quad (1.7)$$

$$G_3(a, b; x, y) = \sum_{m,n=0}^{\infty} (a)_{2n-m}(b)_{2m-n} \frac{x^m y^n}{m! n!}, \quad (1.8)$$

$$H_1(a, b, c; d; x, y) = \sum_{m,n=0}^{\infty} \frac{(a)_{m-n}(b)_{m+n}(c)_n}{(d)_m} \frac{x^m y^n}{m! n!}, \quad (1.9)$$

$$H_2(a, b, c, d; e; x, y) = \sum_{m,n=0}^{\infty} \frac{(a)_{m-n}(b)_m(c)_n(d)_n}{(e)_m} \frac{x^m y^n}{m! n!}, \quad (1.10)$$

$$H_3(a, b, c; x, y) = \sum_{m,n=0}^{\infty} \frac{(a)_{2m+n}(b)_n}{(c)_{m+n}} \frac{x^m y^n}{m! n!}, \quad (1.11)$$

$$H_4(a, b, c, d; x, y) = \sum_{m,n=0}^{\infty} \frac{(a)_{2m+n}(b)_n}{(c)_m(c)_n} \frac{x^m y^n}{m! n!}, \quad (1.12)$$

$$H_5(a, b, c; x, y) = \sum_{m,n=0}^{\infty} \frac{(a)_{2m+n}(b)_{n-m}}{(c)_n} \frac{x^m y^n}{m! n!}, \quad (1.13)$$

$$H_6(a, b, c; x, y) = \sum_{m,n=0}^{\infty} (a)_{2m-n}(b)_{n-m}(c)_n \frac{x^m y^n}{m! n!}, \quad (1.14)$$

$$H_7(a, b, c; d; x, y) = \sum_{m,n=0}^{\infty} \frac{(a)_{2m-n}(b)_n(c)_n}{(d)_m} \frac{x^m y^n}{m! n!}. \quad (1.15)$$

Ref

7. A. Erdelyi, W. Magnus, F. Oberhettinger and F.G. Tricomi, *Higher Transcendental Functions*, Vol. I, McGraw-Hill Book Company, New York, Toronto and London, 1953.

In this paper, we aim to establish the further integral representation of Euler type for Horn double hypergeometric functions $G_1, G_2, G_3, H_1, H_2, H_5, H_6$ and H_7

II. MAIN RESULTS

Theorem 2.1. The following integral representations hold:

$$G_1(a, b, c; x, y) = \frac{\Gamma(b+b')}{\Gamma(b)\Gamma(b')} \int_0^\infty \alpha^{b-1} (1+\alpha)^{-(b+b')} H_1\left(c, a, b+b'; b'; \frac{x}{\alpha}, \frac{\alpha y}{(1+\alpha)}\right) d\alpha, \quad (Re(b) > 0, Re(b') > 0), \quad (1)$$

$$G_1(a, b, c; x, y) = \frac{\Gamma(b+b')\Gamma(c+c')}{2^{b+b'+c+c'-2}\Gamma(b)\Gamma(b')\Gamma(c)\Gamma(c')} \int_{-1}^1 \int_{-1}^1 (1+\alpha)^{b'-1} (1-\alpha)^{b-1} (1+\beta)^{c'-1} \\ \times (1-\beta)^{c-1} F_2\left(a, c+c', b+b'; b', c'; \frac{(1+\alpha)(1-\beta)x}{2(1-\alpha)}, \frac{(1-\alpha)(1+\beta)y}{2(1-\beta)}\right) d\alpha d\beta, \\ (Re(b) > 0, Re(b') > 0, Re(c) > 0, Re(c') > 0), \quad (2)$$

$$G_1(a, b, c; x, y) = \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)(S-R)^{a+b-1}} \int_R^S (\alpha-R)^{a-1} (S-\alpha)^{b+c-1} [(S-\alpha) - (\alpha-R)x]^{-c} \\ \times {}_2F_1\left(\frac{a+b}{2}, \frac{a+b+1}{2}; 1-c; \frac{-4[(\alpha-R)(S-\alpha) - (\alpha-R)^2 x]y}{(S-R)^2}\right) d\alpha, \\ (Re(a) > 0, Re(b) > 0, R < S), \quad (3)$$

$$G_1(a, b, c; x, y) = \frac{2M^{a+b}\Gamma(b+c)}{\Gamma(b)\Gamma(c)} \int_0^{\frac{\pi}{2}} (\sin^2 \alpha)^{b-\frac{1}{2}} (\cos^2 \alpha)^{c-\frac{1}{2}} (\cos^2 \alpha + M \sin^2 \alpha)^{-(b+c)} \\ \times [M - x \cot^2 \alpha - M^2 y \tan^2 \alpha]^{-a} d\alpha, \\ (Re(b) > 0, Re(c) > 0, M > 0). \quad (4)$$

Proof. To prove the result in equality (1) asserted in Theorem 2.1, let $\mathcal{U}$ denote the right-hand side of the equality (1). Then from the definition of Horn's function H_1 in (1.9), we obtain

$$\mathcal{U} = \frac{\Gamma(b+b')}{\Gamma(b)\Gamma(b')} \sum_{m,n=0}^{\infty} \frac{(c)_{m-n}(a)_{m+n}(b+b')_n}{(b')_m} \int_0^\infty \frac{\alpha^{b+n-m-1}}{(1+\alpha)^{b+b'+n}} d\alpha. \quad (5)$$

Employing the integral representation of the Beta function (see, e.g., [7, p. 9, Eq. (2)])

$$B(a, b) = \int_0^\infty \frac{\alpha^{a-1}}{(1+\alpha)^{a+b}} d\alpha, \quad (Re(a) > 0, Re(b) > 0),$$

in (5), we have

$$\mathcal{U} = \frac{\Gamma(b+b')}{\Gamma(b)\Gamma(b')} \sum_{m,n=0}^{\infty} \frac{(c)_{m-n}(a)_{m+n}(b+b')_n B(b+n-m, b'+m)}{(b')_m}. \quad (6)$$

Now applying well known beta function (see, e.g., [16, Section 1.1])

$$B(a, b) = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)},$$

in (6), we are led to the desired result. Then, the similar way we can easily get (2)-(4).

The following theorem can be proved, like Theorem 2.1. So the details are omitted.

Theorem 2.2. The following integral representations holds:

$$\begin{aligned} G_2(a, b, c, d; x, y) &= \frac{\Gamma(a+b)\Gamma(c+c')\Gamma(d+d')}{2^{a+b+c+c'+d+d'-6}\Gamma(a)\Gamma(b)\Gamma(c)\Gamma(c')\Gamma(d)\Gamma(d')} \int_{-1}^1 \int_{-1}^1 \int_{-1}^1 [(1+\alpha)^2]^{a-\frac{1}{2}} \\ &\times [(1-\alpha)^2]^{b-\frac{1}{2}} (1+\alpha^2)^{-(a+b)} [(1+\beta)^2]^{c'-\frac{1}{2}} [(1-\beta)^2]^{c-\frac{1}{2}} (1+\beta^2)^{-(c+c')} [(1+\gamma)^2]^{d'-\frac{1}{2}} \\ &\times [(1-\gamma)^2]^{d-\frac{1}{2}} (1+\gamma^2)^{-(d+d')} \\ &\times F_2\left(a+b, d+d', c+c'; c', d'; \frac{(1+\alpha)^2(1+\beta)^2(1-\gamma)^2x}{4(1+\alpha^2)(1-\beta)^2(1+\gamma^2)}, \frac{(1-\alpha)^2(1-\beta)^2(1+\gamma)^2y}{4(1+\alpha^2)(1+\beta^2)(1-\gamma)^2}\right) d\alpha d\beta d\gamma, \\ &(Re(a) > 0, Re(b) > 0, Re(c) > 0, Re(c') > 0, Re(d) > 0, Re(d') > 0), \end{aligned} \quad (7)$$

$$\begin{aligned} G_2(a, b, c, d; x, y) &= \frac{\Gamma(a+b)\Gamma(c+d)}{\Gamma(a)\Gamma(b)\Gamma(c)\Gamma(d)} \int_0^\infty \int_0^\infty (e^{-\alpha})^a (1-e^{-\alpha})^{b-1} (e^{-\beta})^c (1-e^{-\beta})^{a+b+d-1} \\ &\times \left[(1-e^{-\beta}) - e^{-\alpha+\beta} (1-e^{-\beta})^2 x - e^{-\beta} (1-e^{-\alpha}) y \right]^{-(a+b)} d\alpha d\beta, \\ &(Re(a) > 0, Re(b) > 0, Re(c) > 0, Re(d) > 0), \end{aligned} \quad (8)$$

$$\begin{aligned} G_2(a, b, c, d; x, y) &= \frac{\Gamma(a+c)\Gamma(b+d)(S_1-T_1)^a(R_1-T_1)^{b+c+d}(S_2-T_2)^b(R_2-T_2)^{a+c+d}}{\Gamma(a)\Gamma(b)\Gamma(c)\Gamma(d)} \\ &\times \int_{R_1}^{S_1} \int_{R_2}^{S_2} (\alpha-R_1)^{a-1} (S_1-\alpha)^{b+c+d-1} (\beta-R_2)^{b-1} (S_2-\beta)^{a+c+d-1} \\ &\times [(R_1-T_1)(S_1-\alpha)(S_2-R_2)(\beta-T_2) - (S_1-T_1)(\alpha-R_1)(R_2-T_2)(S_2-\beta)x]^{-(b+d)} \\ &\times [(S_1-R_1)(\alpha-T_1)(R_2-T_2)(S_2-\beta) - (R_1-T_1)(S_1-\alpha)(S_2-T_2)(\beta-R_2)y]^{-(a+c)} d\alpha d\beta, \\ &(Re(a) > 0, Re(b) > 0, Re(c) > 0, Re(d) > 0), \end{aligned} \quad (9)$$

$$\begin{aligned} G_2(a, b, c, d; x, y) &= \frac{2M^{a+c}\Gamma(c+d)}{\Gamma(c)\Gamma(d)} \int_0^\infty \cosh \alpha (\sinh^2 \alpha)^{a+d-\frac{1}{2}} (1+M \sinh^2 \alpha)^{-(c+d)} \\ &\times (M \sinh^2 \alpha - x)^{-a} (1-M \sinh^2 \alpha)^{-b} d\alpha, \\ &(Re(c) > 0, Re(d) > 0, M > 0). \end{aligned} \quad (10)$$

Theorem 2.3. The following integral representations hold:

$$G_3(a, b; x, y) = \frac{2M^a \Gamma(a+a')}{\Gamma(a)\Gamma(a')} \int_0^{\frac{\pi}{2}} (\sin^2 \alpha)^{a-\frac{1}{2}} (\cos^2 \alpha)^{a'-\frac{1}{2}} (\cos^2 \alpha + M \sin^2 \alpha)^{-(a+a')} \\ \times H_6 \left(b, a+a', 1-a'; \left(1 + \frac{1}{M} \cot^2 \alpha\right) x, -\frac{M^2 y \sin^2 \alpha \tan^2 \alpha}{\cos^2 \alpha + M \sin^2 \alpha} \right) d\alpha, \\ (Re(a) > 0, Re(a') > 0, M > 0), \quad (11)$$

$$G_3(a, b; x, y) = \frac{\Gamma(a+a')}{\Gamma(a)\Gamma(a')} \int_{-\frac{1}{2}}^{\frac{1}{2}} \left(\frac{1}{2} + \alpha\right)^{a'-1} \left(\frac{1}{2} - \alpha\right)^{a-1} \\ \times H_7 \left(b, \frac{a+a'}{2}, \frac{a+a'+1}{2}; a'; \frac{(1+2\alpha)x}{(1-2\alpha)}, (1-2\alpha)^2 y \right) d\alpha, \\ (Re(a) > 0, Re(a') > 0), \quad (12)$$

$$G_3(a, b; x, y) = \frac{(1+M_1)^a (1+M_2)^b \Gamma(a+a') \Gamma(b+b')}{\Gamma(a)\Gamma(a')\Gamma(b)\Gamma(b')} \int_0^1 \int_0^1 \alpha^{a-1} (1-\alpha)^{a'-1} (1+M_1\alpha)^{-(a+a')} \\ \times \beta^{b-1} (1-\beta)^{b'-1} (1+M_2\beta)^{-(b+b')} {}_2F_1 \left(\frac{b+b'}{2}, \frac{b+b'+1}{2}; a'; \frac{4(1+M_2)^2 (1-\alpha)\beta^2 x}{(1+M_1)\alpha(1+M_2\beta)^2} \right) \\ \times {}_2F_1 \left(\frac{a+a'}{2}, \frac{a+a'+1}{2}; b'; \frac{4(1+M_1)^2 \alpha^2 (1-\beta)y}{(1+M_2)(1+M_1\alpha)^2 \beta} \right) d\alpha d\beta, \\ (Re(a) > 0, Re(a') > 0, Re(b) > 0, Re(b') > 0, M_1 > -1, M_2 > -1), \quad (13)$$

$$G_3(a, b; x, y) = \frac{2(1+M)^{a+2b} \Gamma(a+b)}{\Gamma(a)\Gamma(b)} \int_0^{\frac{\pi}{2}} (\sin^2 \alpha)^{b-\frac{1}{2}} (\cos^2 \alpha)^{a-\frac{1}{2}} \\ \times [(1+M)(1+M \sin^2 \alpha) - (1+M)^3 x \sin^2 \alpha \tan^2 \alpha - y \cos^2 \alpha \cot^2 \alpha]^{-(a+b)} d\alpha, \\ (Re(a) > 0, Re(b) > 0, M > -1), \quad (14)$$

Theorem 2.4. The following integral representations hold:

$$H_1(a, b, c; d; x, y) = \frac{\Gamma(2d)}{\Gamma(a)\Gamma(2d-a)(S-R)^{2d-1}} \int_R^S (\alpha-R)^{2d-a-1} (S-\alpha)^{a-1} \\ \times F_1 \left(b, d + \frac{1}{2}, c; 2d-a; \frac{4(\alpha-R)(S-\alpha)x}{(S-R)^2}, \frac{(\alpha-R)y}{(S-\alpha)} \right) d\alpha, \\ (Re(a) > 0, Re(2d-a) > 0, R < S), \quad (15)$$

$$H_1(a, b, c; d; x, y) = \frac{2\Gamma(a+a')}{\Gamma(a)\Gamma(a')} \int_0^{\frac{\pi}{2}} (\sin^2 \alpha)^{a-\frac{1}{2}} (\cos^2 \alpha)^{a'-\frac{1}{2}} \\ \times F_2(b, a+a', c; d, a'; x \sin^2 \alpha, y \cot^2 \alpha) d\alpha, \\ (Re(a) > 0, Re(a') > 0), \quad (16)$$

$$H_1(a, b, c; d; x, y) = \frac{\Gamma(d)(S-T)^b(R-T)^{d-b}}{\Gamma(b)\Gamma(d-b)(S-R)^{d-a-1}} \int_R^S (\alpha-R)^{b-1}(S-\alpha)^{d-b-1}(\alpha-T)^{a-d} \\ \times [(S-R)(\alpha-T) - (S-T)(\alpha-R)x]^{-a} \\ \times {}_2F_1\left(c, 1+b-d; 1-a; \frac{(S-T)(\alpha-R)[(S-R)(\alpha-T) - (S-T)(\alpha-R)x]y}{(R-T)(S-R)(S-\alpha)(\alpha-T)}\right) d\alpha, \\ (Re(b) > 0, Re(d-b) > 0, T < R < S), \quad (17)$$

$$H_1(a, b, c; d; x, y) = \frac{(1+M)^{a+c}\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \int_0^1 \alpha^{a-1}(1-\alpha)^{b-1}(1+M\alpha)^{-(a+b)} \\ \times [(1+M)\alpha - (1-\alpha)y]^{-c} {}_2F_1\left(\frac{a+b}{2}, \frac{a+b+1}{2}; d; \frac{4(1+M)\alpha(1-\alpha)x}{(1+M\alpha)^2}\right) d\alpha, \\ (Re(a) > 0, Re(b) > 0, M > -1). \quad (18)$$

Corollary 2.5. Let $y = 0$ in (18). Then the following result holds true:

$${}_2F_1(a, b; d; x) = \frac{(1+M)^a\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \int_0^1 \alpha^{a-1}(1-\alpha)^{b-1}(1+M\alpha)^{-(a+b)} \\ \times {}_2F_1\left(\frac{a+b}{2}, \frac{a+b+1}{2}; d; \frac{4(1+M)\alpha(1-\alpha)x}{(1+M\alpha)^2}\right) d\alpha, \\ (Re(a) > 0, Re(b) > 0, M > -1). \quad (19)$$

Theorem 2.6. The following integral representations hold:

$$H_2(a, b, c, d; e; x, y) = \frac{\Gamma(b+c)\Gamma(2e)}{\Gamma(a)\Gamma(b)\Gamma(c)\Gamma(2e-a)(S_1-R_1)^{b+c-1}(S_2-R_2)^{2e-1}} \int_{R_1}^{S_1} \int_{R_2}^{S_2} \\ \times (\alpha-R_1)^{b-1}(S_1-\alpha)^{c-1}(\beta-R_2)^{a-1}(S_2-\beta)^{2e-a-1} \\ \times {}_F_1\left(b+c, e+\frac{1}{2}; d; 2e-a; \frac{4(\alpha-R_1)(\beta-R_2)(S_2-\beta)x}{(S_1-R_1)(S_2-R_2)^2}, \frac{(S_1-\alpha)(S_2-\beta)y}{(S_1-R_1)(\beta-R_2)}\right) d\alpha d\beta, \\ (Re(a) > 0, Re(b) > 0, Re(c) > 0, Re(2e-a) > 0, R_1 < S_1, R_2 < S_2), \quad (20)$$

$$H_2(a, b, c, d; e; x, y) = \frac{\Gamma(2e)}{2^{2(e-1)}\Gamma(a)\Gamma(2e-a)} \int_{-1}^1 [(1+\alpha)^2]^{2e-a-\frac{1}{2}} [(1-\alpha)^2]^{a-\frac{1}{2}} \\ \times (1+\alpha^2)^{-e} F_3\left(b, c, e+\frac{1}{2}; d; 2e-a; \left(\frac{1-\alpha^2}{1+\alpha^2}\right)^2 x, \left(\frac{1+\alpha}{1-\alpha}\right)^2 y\right) d\alpha, \\ (Re(a) > 0, Re(2e-a) > 0), \quad (21)$$

$$H_2(a, b, c, d; e; x, y) = \frac{(1+M)^a\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \int_0^1 \alpha^{a-1}(1-\alpha)^{b-1}(1+M\alpha)^{-(a+b)} \\ \times H_7\left(a+b, c, d; e; \frac{(1+M)\alpha(1-\alpha)x}{(1+M\alpha)^2}, \frac{(1+M\alpha)y}{(1+M\alpha)\alpha}\right) d\alpha, \\ (Re(a) > 0, Re(b) > 0, M > -1), \quad (22)$$

$$\begin{aligned}
 H_2(a, b, c, d; e; x, y) &= \frac{2M^{e-b}\Gamma(e)}{\Gamma(b)\Gamma(e-b)} \int_0^{\frac{\pi}{2}} (\sin^2 \alpha)^{e-b-\frac{1}{2}} (\cos^2 \alpha)^{b-\frac{1}{2}} (\cos^2 \alpha + M \sin^2 \alpha)^{a-e} \\
 &\times [(\cos^2 \alpha + M \sin^2 \alpha) - x \cos^2 \alpha]^{-a} {}_2F_1\left(c, d; 1-a; \left[\frac{x \cos^2 \alpha}{(\cos^2 \alpha + M \sin^2 \alpha)} - 1\right] y\right) d\alpha, \\
 &(Re(b) > 0, Re(e-b) > 0, M > 0).
 \end{aligned} \tag{23}$$

Theorem 2.7. The following integral representations hold:

$$\begin{aligned}
 H_5(a, b; c; x, y) &= \frac{\Gamma(2c)}{\Gamma(b)\Gamma(2c-b)} \int_0^\infty (e^{-\alpha})^b (1-e^{-\alpha})^{2c-b-1} \\
 &\times H_3\left(a, c + \frac{1}{2}; 2c-b; (e^\alpha - 1)x, 4e^{-\alpha}(1-e^{-\alpha})y\right) d\alpha, \\
 &(Re(b) > 0, Re(2c-b) > 0),
 \end{aligned} \tag{24}$$

$$\begin{aligned}
 H_5(a, b; c; x, y) &= \frac{\Gamma(b+b')}{2^{b+b'-1}\Gamma(b)\Gamma(b')} \int_{-1}^1 (1+\alpha)^{b'-1} (1-\alpha)^{b-1} \\
 &\times H_4\left(a, b+b'; b', c; \left(\frac{1+\alpha}{1-\alpha}\right)x, \frac{(1-\alpha)}{2}y\right) d\alpha, \\
 &(Re(b) > 0, Re(b') > 0),
 \end{aligned} \tag{25}$$

$$\begin{aligned}
 H_5(a, b; c; x, y) &= \frac{2M^a\Gamma(c)}{\Gamma(a)\Gamma(c-a)} \int_0^\infty \cosh \alpha (\sinh^2 \alpha)^{a-\frac{1}{2}} (1 + M \sinh^2 \alpha)^{b-c} \\
 &\times [(1 + M \sinh^2 \alpha) - M y \sinh^2 \alpha]^{-b} \\
 &\times {}_2F_1\left(\frac{1+a-c}{2}, \frac{a-c}{2} + 1; 1-b; -\frac{4M^2 x \sinh^4 \alpha [(1 + M \sinh^2 \alpha) - M y \sinh^2 \alpha]}{(1 + M \sinh^2 \alpha)}\right) d\alpha, \\
 &(Re(a) > 0, Re(c) > 0, M > 0),
 \end{aligned} \tag{26}$$

$$\begin{aligned}
 H_5(a, b; c; x, y) &= \frac{2M^{c-a}\Gamma(c)}{\Gamma(a)\Gamma(c-a)} \int_0^\infty \cosh \alpha (\sinh^2 \alpha)^{c-a-\frac{1}{2}} (1 + M \sinh^2 \alpha)^{b-c} \\
 &\times [(1 + M \sinh^2 \alpha) - y]^{-b} \\
 &\times {}_2F_1\left(\frac{1+a-c}{2}, \frac{a-c}{2} + 1; 1-b; -\frac{4x [(1 + M \sinh^2 \alpha) - M y \sinh^2 \alpha]}{M^2 \sinh^4 \alpha (1 + M \sinh^2 \alpha)}\right) d\alpha, \\
 &(Re(a) > 0, Re(c) > 0, M > 0).
 \end{aligned} \tag{27}$$

Corollary 2.8. Let $x = 0$ in (26). Then the following result holds true:

$$\begin{aligned}
 {}_2F_1(a, b; c; y) &= \frac{2M^a\Gamma(c)}{\Gamma(a)\Gamma(c-a)} \int_0^\infty \cosh \alpha (\sinh^2 \alpha)^{a-\frac{1}{2}} (1 + M \sinh^2 \alpha)^{b-c} \\
 &\times [(1 + M \sinh^2 \alpha) - M y \sinh^2 \alpha]^{-b} d\alpha, \\
 &(Re(a) > 0, Re(c) > 0, M > 0).
 \end{aligned} \tag{28}$$

Theorem 2.9. *The following integral representations hold:*

$$\begin{aligned}
 H_6(a, b, c; x, y) &= \frac{\Gamma(a+c)\Gamma(b+b')}{\Gamma(a)\Gamma(b)\Gamma(b')\Gamma(c)} \int_{-\frac{1}{2}}^{\frac{1}{2}} \int_{-\frac{1}{2}}^{\frac{1}{2}} \left(\frac{1}{2} + \alpha\right)^{c-1} \left(\frac{1}{2} - \alpha\right)^{a-1} \\
 &\quad \times \left(\frac{1}{2} + \beta\right)^{b'-1} \left(\frac{1}{2} - \beta\right)^{b-1} \\
 &\quad F_3\left(\frac{a+c}{2}, \frac{b+b'}{2}, \frac{a+c+1}{2}, \frac{b+b'+1}{2}; b'; \frac{(1-2\alpha)^2(1+2\beta)x}{(1-2\beta)}, \frac{(1+2\alpha)(1-4\beta^2)y}{(1-2\alpha)}\right) d\alpha d\beta, \\
 &\quad (Re(a) > 0, Re(b) > 0, Re(b') > 0, Re(c) > 0), \tag{29}
 \end{aligned}$$

$$\begin{aligned}
 H_6(a, b, c; x, y) &= \frac{\Gamma(a+a')}{\Gamma(a)\Gamma(a')} \int_0^\infty \alpha^{a-1} (1+\alpha)^{-(a+a')} G_2\left(a+a', c, b, 1-a'; \frac{-\alpha^2 x}{(1+\alpha)}, \frac{-y}{\alpha}\right) d\alpha, \\
 &\quad (Re(a) > 0, Re(a') > 0), \tag{30}
 \end{aligned}$$

$$\begin{aligned}
 H_6(a, b, c; x, y) &= \frac{\Gamma(b+b')}{2^{b+b'-2}\Gamma(b)\Gamma(b')} \int_{-1}^1 [(1+\alpha)^2]^{b'-\frac{1}{2}} [(1-\alpha)^2]^{b-\frac{1}{2}} (1+\alpha^2)^{-(b+b')} \\
 &\quad \times H_6\left(a, 1-b', c; -\left(\frac{1+\alpha}{1-\alpha}\right)^2 x, -\left(\frac{1-\alpha}{1+\alpha}\right)^2 y\right) d\alpha, \\
 &\quad (Re(b) > 0, Re(b') > 0), \tag{31}
 \end{aligned}$$

$$\begin{aligned}
 H_6(a, b, c; x, y) &= \frac{4(1+M_1)^{a+b+b'}(1+M_2)^{b'}\Gamma(a+c)\Gamma(b+b')}{\Gamma(a)\Gamma(b)\Gamma(b')\Gamma(c)} \int_0^{\frac{\pi}{2}} \int_0^{\frac{\pi}{2}} (\sin^2 \alpha)^{a-\frac{1}{2}} \\
 &\quad \times (\cos^2 \alpha)^{c-\frac{1}{2}} (\sin^2 \beta)^{b'-\frac{1}{2}} (\cos^2 \beta)^{b-\frac{1}{2}} (1+M_1 \sin^2 \alpha)^{-(a+c)} \\
 &\quad \times [(1+M_1)(1+M_2 \sin^2 \beta) - y \cot^2 \alpha \cos^2 \beta]^{-(b+b')} \\
 &\quad \times {}_2F_1\left(\frac{a+c}{2}, \frac{a+c+1}{2}; b'; \frac{4(1+M_1)^2(1+M_2)x \sin^4 \alpha \tan^2 \beta}{(1+M_1 \sin^2 \alpha)^2}\right) d\alpha d\beta, \\
 &\quad (Re(a) > 0, Re(b) > 0, Re(b') > 0, Re(c) > 0, M_1 > -1, M_2 > -1). \tag{32}
 \end{aligned}$$

Theorem 2.10. *The following integral representations hold:*

$$\begin{aligned}
 H_7(a, b, c; d; x, y) &= \frac{\Gamma(a+a')(S-T)^a(R-T)^{a'}}{\Gamma(a)\Gamma(a')(S-R)^{a+a'-1}} \int_R^S (\alpha-R)^{a-1} (S-\alpha)^{a'-1} (\alpha-T)^{-(a+a')} \\
 &\quad \times {}_2F_1\left(\frac{a+a'}{2}, \frac{a+a'+1}{2}; d; \left(\frac{2(S-T)(\alpha-R)}{(S-R)(\alpha-T)}\right)^2 x\right) {}_2F_1\left(b, c; a'; \frac{(R-T)(S-\alpha)y}{(S-T)(\alpha-R)}\right) d\alpha, \\
 &\quad (Re(a) > 0, Re(a') > 0, T < R < S), \tag{33}
 \end{aligned}$$

$$H_7(a, b, c; d; x, y) = \frac{\Gamma(d)}{\Gamma(b)\Gamma(d-b)} \int_0^\infty (e^{-\alpha})^b (1 - e^{-\alpha})^{d-b-1} \\ \times H_6\left(a, 1+b-d, c; (e^{-\alpha}-1)x, \frac{y}{(1-e^{-\alpha})}\right) d\alpha, \\ (Re(b) > 0, Re(d-b) > 0), \quad (34)$$

$$H_7(a, b, c; d; x, y) = \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)(S-R)^{a+b-1}} \int_R^S (\alpha-R)^{a+c-1} (S-\alpha)^{b-1} \\ \times [(\alpha-R) - (S-\alpha)y]^{-c} {}_2F_1\left(\frac{a+b}{2}, \frac{a+b+1}{2}; d; 4\left(\frac{\alpha-R}{S-R}\right)^2 x\right) d\alpha, \\ (Re(a) > 0, Re(b) > 0, R < S), \quad (35)$$

$$H_7(a, b, c; d; x, y) = \frac{2\Gamma(a+a')}{\Gamma(a)\Gamma(a')} \int_0^{\frac{\pi}{2}} (\sin^2 \alpha)^{a'-\frac{1}{2}} (\cos^2 \alpha)^{a-\frac{1}{2}} \\ \times H_2(1-a', a+a', b, c; d; -x \cot^2 \alpha \cos^2 \alpha, -y \tan^2 \alpha) d\alpha, \\ (Re(a) > 0, Re(a') > 0). \quad (36)$$

Corollary 2.11. Let $y = 0$ in (32). Then the following result holds true:

$${}_2F_1\left(\frac{a}{2}, \frac{a+1}{2}; d; 4x\right) = \frac{\Gamma(a+a')(S-T)^a (R-T)^{a'}}{\Gamma(a)\Gamma(a')(S-R)^{a+a'-1}} \int_R^S (\alpha-R)^{a-1} (S-\alpha)^{a'-1} \\ \times (\alpha-T)^{-(a+a')} {}_2F_1\left(\frac{a+a'}{2}, \frac{a+a'+1}{2}; d; 4\left(\frac{(S-T)(\alpha-R)}{(S-R)(\alpha-T)}\right)^2 x\right) d\alpha, \\ (Re(a) > 0, Re(a') > 0, T < R < S). \quad (37)$$

REFERENCES RÉFÉRENCES REFERENCIAS

1. P. Appell, *Sur les fonctions hypergjeomjetriques de deux variables*, J. Math. Pures Appl., (3)8 (1882), 173216.
2. M. G. Bin-Saad and J. A. Younis, *Certain quadruple hypergeometric series and their integral representations*, Appl. Math., 14 (2019), 1085-1098.
3. Bin-Saad, M.G. and Younis, J. A., *Certain integrals associated with hypergeometric functions of four variables*, Earthline J. Math. Sci., 2 (2019), 325-341.
4. M. G. Bin-Saad and J. A. Younis, *Certain integral representations of some quadruple hypergeometric series*, Palestine J. Math., 9 (2020), 132-141.
5. M. G. Bin-Saad and J. A. Younis, *Some integrals connected with a new quadruple hypergeometric series*, Universal J. Math. Appl., 3 (2020), 19-27.

6. J. Choi, A. Hasanov, H.M. Srivastava and M. Turaev, *Integral representations for Srivastava's triple hypergeometric functions*, Taiwanese J. Math. 15 (2011), 2751-2762.
7. A. Erdjelyi, W. Magnus, F. Oberhettinger and F.G. Tricomi, *Higher Transcendental Functions*, Vol. I, McGraw-Hill Book Company, New York, Toronto and London, 1953.
8. H. Exton, *Multiple Hypergeometric Functions and Applications*, Halsted Press, New York, London, Sydney and Toronto, 1976.
9. A. Hasanov, M.G. Bin Saad and A. Ryskan, *Some properties of Horn type second order double hypergeometric series*, Bulletin KRASEC. Phys. Math. Sci., 21 (2018), 32-47.
10. A. Hasanov, H.M. Srivastava and M. Turaev, *Decomposition formulas for some triple hypergeometric functions*, J. Math. Anal. Appl., 324 (2006), 955-969.
11. J. Horn, *Ueber die Convergenz der Hypergeometrische Reihen Zweierund Dreier Veränderlichen*, Math. Ann., 34 (1889), 544-600.
12. S. Jun, I. Kim and A.K. Rathie, *On a new class of Eulerians type integrals involving generalized hypergeometric functions*, Aust. J. Math. Anal. Appl. 2019, 16, 10.
13. P.O.M. Olsson, *hypergeometric function of two variables of importance in perturbation theory*, I, Ark. Fys., B30 (1965), 187191.
14. E.D. Rainville, *Special Functions*, Macmillan Company, New York, 1960, Reprinted by Chelsea Publishing Company, Bronx, New York, 1971.
15. G. Sharma and A.K. Rathie, *Integrals of hypergeometric series*, Vij. Pari. Anu. Patri., 34(1991), 26-29.
16. H.M. Srivastava and J. Choi, *Zeta and q-Zeta Functions and Associated Series and Integrals*, Elsevier Science Publishers, Amsterdam, London and New York, 2012.
17. H.M. Srivastava and H.L. Manocha, *A Treatise on Generating Functions*, Halsted Press, Ellis Horwood Limited, Chichester, John Wiley and Sons, New York, Chichester, Brisbane and Toronto, 1984.
18. H.M. Srivastava and R. Panda, *Some hypergeometric transformations involving Horn's function H₃*, Rend. Mat., 6 (1973), 859-864.
19. J.A. Younis and M.G. Bin-Saad, *Integral representations involving new hypergeometric functions of four variables*, J. Frac. Calc. Appl., 10 (2019), 77-91.
20. J.A. Younis, Maged G. Bin-Saad, *Integral representations and operational relations involving some quadruple hypergeometric functions*, J. Frac. Calc. Appl., 11 (2020), 62-74.
21. J. A. Younis, K. S. Nisar, *Several Euler-type integrals involving Extons quadruple hypergeometric series*, J. Math. Computer Sci., 21 (2020), 286-295.