



In Vitro Testing of the Trichoderma Species as Bio-control Agents of (*Neofusicoccum. Mangiferae. Crous*) Pathogen of Branch Disease in Shade and Fruit Trees

By Fatin Abdalla Osman, Mohamed Abbaker Hassan & Taha Mohamed Shareef

University of Zalingei

Abstract- This study carried out in 2013 to test the fungicidal potentiality of three Trichoderma species under laboratory condition. The laboratory experiments were conducted at the laboratory of plant quarantine- Administration of Plant Protection Ministry of Agriculture. Any Trichoderma specie was considered as treatment in addition to one synthetic fungicide (Amistar-Top) for comparison, each treatment replicated 6 times and arranged in CRD. The result revealed that: Amistar-Top at concentration of 200 ppm, had given significant inhibition compared with the three species of Trichoderma with different isolates of the pathogen. However, there was no significant difference between inhibition percentages from Amistar-Top in some treatments. More over *T. harzianum* with *N. mangiferae* isolated from *Mangifera indica* and *T. viride* with the fungal isolate from *F. benjamina* were more effective than Amistar-Top.

Keywords: *trichoedrma spp., pathogen, neofusicoccum. mangiferae .crous, inhibition.*

GJSFR-D Classification: DDC Code: 628.161 LCC Code: QR105.5



Strictly as per the compliance and regulations of:



In Vitro Testing of the Trichoderma Species as Bio-control Agents of (Neofusicoccum. Mangiferae. Crous) Pathogen of Branch Disease in Shade and Fruit Trees

Fatin Abdalla Osman ^a, Mohamed Abbaker Hassan ^a & Taha Mohamed Shareef ^b

Abstract This study carried out in 2013 to test the fungicidal potentiality of three Trichoderma species under laboratory condition. The laboratory experiments were conducted at the laboratory of plant quarantine- Administration of Plant Protection Ministry of Agriculture. Any Trichoderma specie was considered as treatment in addition to one synthetic fungicide (Amistar-Top) for comparison, each treatment replicated 6 times and arranged in CRD. The result revealed that: Amistar-Top at concentration of 200 ppm, had given significant inhibition compared with the three species of Trichoderma with different isolates of the pathogen. However, there was no significant difference between inhibition percentages from Amistar-Top in some treatments. More over *T. harzianum* with *N. mangiferae* isolated from *Mangifera indica* and *T. viride* with the fungal isolate from *F. benjamina* were more effective than Amistar-Top.

Keywords: *trichoedrma spp.*, pathogen, *neofusicoccum. mangiferae .crous*, inhibition.

I. INTRODUCTION

One of the strategies used to control pathogens is mycoparasitism whereby a species or strain of fungus directly attacks and feeds on other fungi (Harman, 2000). The antagonistic ability of *Trichoderma* species was discovered 70 years ago. *Trichoedrma spp.* are now the most common fungal biological control agents throughout the world. The primary mechanism of antagonism in *Trichoderma* is mycoparaasitism (Chet, 1987).

Over the past several decades, various attempts to control plant diseases have been made for eradication or prevention through the development of systemic fungicides. Continued and repeated application of fungicides has disturbed the biological control by natural enemies and let to out-break in disease and development of resistance to various types of fungicides toxicity to non-target organisms and environmental problems (Hayes and Law, 1991). In addition, use of fungicides is uneconomic due to longevity of trees that necessitates repeated application

over a long period of time. Triazoles antifungal such as tebuconazole are now widely used for treatment of fungal infections due to their broad spectra activity and improved safety profile compared to other fungicides (Kamai et al., 2002).

Due to the importance of bio-agents to control plant diseases, this work is considered a contribution in this aspect especially, to test inhibition activity of three *Trichoderma* species to *N. mangiferae*.

II. MATERIALS AND METHODS

a) Effect of *Trichoderma* Species on the Growth of *N. Mangiferae*

Three species of the genus *Trichoderma* were obtained from the laboratory of plant quarantine-Administration of Plant Protection, Ministry of Agriculture. Active pure culture from each species was prepared by transferring samples from each species into PDA plates and incubated at 28°C for seven days (until the Petri dishes completely covered). Five mm discs of PDA from seven days old culture of each isolate of *N. mangiferae* and the same size disc from seven days old cultures of *Trichoderma* species were placed 3 cm apart from each other and 3 cm from the edge of the plate. The antagonistic potentiality of the three *Trichoderma* species against *N. mangiferae* isolates was measured as inhibition % considering the radial growth in the opposite directions from the centre of the inoculums discs as control (R) and the growth towards the disc of pathogen as treatment (R₁). The inhibition percent was calculated according to the following formula;

$$I \% = R - R_1 / R$$

I % = Inhibition percentage

Whereas R= the growth length from the centre of the inoculums disc towards the edge of the plate

R₁= the growth towards *Trichoderma* disc

b) Effect of Amistar-Top on the Radial Growth of *N. Mangiferae*

Five mm discs of PDA from seven days old culture of each isolate of *N. mangiferae* and placed in

Author a: Assistant Prof. Department of Crop Protection and Environmental Studies U. of Alzaeim Alazhari- Sudan.

e-mail: Fatinwadi44@gmail.com



Author a p: Department of Crop Protection, University of Zalingei.



Table 1: Inhibition Percentages of Radial Growth of Four *N. Mangiferae* Isolates Treated with Three Trichoderma Species

Treatments (Trichoderma species)	Isolates of <i>N. Mangiferae</i>			
	Host plants			
	Inhibition%			
Citrus lemon	49.80a	53.02a	66.90 a	
T. harzianum	78.44a	64.30a	68.54a	78.82 a
T. viride	78.76a	54.96a	50.86a	72.78a
LSD	17.63	18.47	35.77	18.47

Mean with the same letter(s) in the column are not significantly different at $P < 0.05$.



III. RESULTS

a) Effect of Trichoderma Species on the Radial Growth of Four Isolates of *N. Mangiferae*

The biological interaction between *N. mangiferae* isolates and *Trichoderma* spp. resulted into reduction of the radial growth of the *N. mangiferae* *In vitro*. The inhibition percent ranged between 49.8% and 82.14% (Fig-1). *Trichoderma viride* was the most effective against *N. mangiferae* isolated from *C. lemon*, it has resulted in 82.14% inhibition percent. The least inhibition percent was of *N. mangiferae* was that of *T. koningei* with *N. mangiferae* isolated from *F. nitida*. However the statistical analysis showed no significant differences between the means of inhibition percent regarding different *Trichoderma* spp. (Table 1).

Fig.-1: Effect of Trichoderma Spp. on the Radial Growth of *N. Mangiferae*

b) *Effect of Different Concentration of Amistar-Top on the Radial Growth of N. Mangiferae Isolates*

The systemic fungicide Amistar-top was very effective in reduction of radial growth of *N. mangiferae* isolates. Three concentrations of the fungicide have been tested *In vitro*: 100, 200 and 500 ppm. The results showed that there was no significant different between

the two concentrations of 200ppm and 500ppm, but they differed significantly from the concentration of 100ppm. This result means that the concentrations 200ppm and 500ppm have the same efficacy (Table-2) and (Fig-2).

Table-2: Effect of Amistar- Top on the Radial Growth of Four Isolates of *N. Mangiferae*

Concentrations	Isolates of <i>N. mangiferae</i>			
	Host plants			
	<i>C. lemon</i>	<i>F. benjamina</i>	<i>F. nitida</i>	<i>M. indica</i>
	Inhibition %			
100ppm	55.98b	52.96 b	52.04b	51.06b
200ppm	88.08a	85.16a	82.72a	85.00a
500ppm	90.44a	89.64a	90.32a	90.75a
LSD	8.077	19.92	17.30	12.43

Mean with the same letter(s) in the column are not significantly different at $P < 0.05$.




Fig.-2: Effect of Amistar Top on the Radial Growth of *N. Mangiferae*

c) *Effect of Three Trichoderma Species on the Growth of N. Mangiferae Isolates Compared with Amistar-Top*

The bio-agents tested were compared with the fungicide Amistar-Top tested in the laboratory showed that *Trichodema* spp. have approximately similar effect as Amistar-Top on the pathogen isolates radial growth and (Fig-3). Amistar-top with *N. mangiferae* isolated from *M. indica* was significantly better than *T. harzianum*. However, there was no significant differences between Amistar-Top effect and *T. viride* with *N. mangiferae* isolated from *Citrus lemon*, *Ficus nitida* and *mangiferae*

indica. On the other hand Amistar-top showed its significant superior to *T. viride* with *N. mangiferae* isolated from *F. benjamina*.






Fig.- 3: Comparison of Inhibition Effect (%) of Three Trichoderma Spp. and Amistar on Four Isolates of *N. Mangiferae*

IV. DISCUSSION

Trichoderma spp. plays a major role as biocontrol agents, owing to their capability of improving crop yield by multiple roles, such as bio-pesticide, bio-herbicide and plant growth promotion (Mausam, 2007). Currently the roles of BCAs are a well-established fact and become increasingly crucial, and in several cases complementary or even replacing the synthetic chemical components where antagonistic fungi play an important part (Templeton, and Heiny, 1989). Bio-interaction between *Trichoderma* spp. and plant pathogenic fungi tested in the laboratory showed that *Trichoderma* species were differently effective in inhibiting the different isolates of the pathogen (*N. mangiferae*). *T. viride* gave the highest inhibitory action on *N. mangiferae* isolated from symptomatic lime tree with 80% inhibition. In counterpart *T. harzianum* have the least inhibition with *N. mangiferae* isolated from *F. benjamina* with 49.8% inhibition. This inhibition of the pathogen can be attributed to the different biological interactions between *Trichoderma* spp. and the pathogen. The systemic fungicide Amstar-Top was used as reference revealed that it is highly effective in reducing the radial growth of different isolates of the pathogen, exactly 88.08% inhibition. This is due to the fact that in case of synthetic chemical fungicides the formulations applied are very accurately calculated according to the recommendations of the company. On the other hand in case of bio-control agents it is difficult to setup formulations for application as the activity of the control agent is subjected to biological factors and

environmental conditions. This fact may give the farmer evident to rely on chemicals more than bio-agents application.

REFERENCES RÉFÉRENCES REFERENCIAS

1. Chet, I. (1987). Trichoderma application, mode of action, and potential as a biological control agent of soilborne plant fungi. In: chet, I. (ed.) Innovative Approaches to Plant Disease Control. John Wiley and Sons, New York, pp. 137-160.
2. Harman, G. E. (2000). The myth and dogmas of biocontrol: changes in perceptions derived from research on *Trichoderma harzianum* stain T-22. *Plant Dis.* 84: 377-393.
3. Hayes, W. J. Jr, Laws E. R. Jr. (1991). Handbook of pesticide toxicology, vols 1-3. New York: Academic Press. Pp. 1576.
4. Kamai, Y; Harasaki, T; Fukuoka, T; Ohya, S; Uchida, K; Yamaguchi, H. and Kuwahara, S; (2002). *In vitro* and *in vivo* activities of CS-758 (R-120758), a new triazole antifungal agent. *Antimicrob Agents Chemother* 46, pp 367-370.
5. Mausam, V; Satinder, K.; Brar, R. D.; Tyagi, R. Y.; Surampalli, J. R. (2007). Antagonistic fungi *Trichoderma* spp. Pansophy of Biological Control. *Biochemical Engineering Journal*. 37: 1-20.
6. Templeton, G. E. and Heiny, D. K. (1989). Improvement of fungi to enhance mycoherbicide potential, In: J. M. Whipps, R.D. Lumsdon (Eds.). biotechnology of fungi for improving plant growth, Cambridge University Press. UK. pp. 127-151.