



GLOBAL JOURNAL OF SCIENCE FRONTIER RESEARCH: F
MATHEMATICS AND DECISION SCIENCES
Volume 22 Issue 1 Version 1.0 Year 2022
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Fractional order Riemann Curvature Tensor in Differential Geometry

By Wedad Saleh
Taibah University


Abstract- This study discussed some interesting aspects and features of fractional curvature in the differential manifold. In particular, Riemannian fractional curvature tensor, Livi-Civita fractional connection and Bianchi fractional identity are presented.

GJSFR-F Classification: DDC Code: 516.36 LCC Code: QA641



Strictly as per the compliance and regulations of:





Fractional Order Riemann Curvature Tensor in Differential Geometry

Wedad Saleh

Abstract- This study discussed some interesting aspects and features of fractional curvature in the differential manifold. In particular, Riemannian fractional curvature tensor, Livi-Civita fractional connection and Bianchi fractional identity are presented.

I. INTRODUCTION

In mathematics, several special functions appear in many applications such as the Gamma function that plays some significant roles in the theory of integral differential equations in particular fractional calculus. Thus, we begin with some definitions, for the details we refer to [1, 15, 8].

The Gamma function of a positive integer η is again a positive integer, while the gamma function $\Gamma(-\eta)$ of a negative integer changes to infinities. The Gamma function any positive η value is defined as follows:

$$\Gamma(\eta) = \int_0^{\infty} t^{\eta-1} e^{-t} dt.$$

The Gamma function $\Gamma(\eta)$ is considered as a generalization of the factorial and $\Gamma(\eta)$ is defined for $\eta > 0$ by the integral

$$\Gamma(\eta) = \int_0^{\infty} t^{\eta-1} e^{-t} dt.$$

In the classical sense since $\Gamma(0) = \frac{\Gamma(1)}{0}$, then it follows that $\Gamma(\eta)$ is not defined for integers $\eta \leq 0$. However, the extension formula gives finite values for $\Gamma(\eta)$, for $\Re(\eta) \leq 0$ since $\Gamma(\eta)$ is analytic everywhere except at $\eta = 0, -1, -2, \dots$, and the residue at $\eta = k$ is given by

$$\text{Res}_{\eta=k} \Gamma(\eta) = \frac{(-1)^k}{k!}.$$

Author: Department of Mathematics, Taibah University, Al- Medina, Saudi Arabia.
e-mail: wed_10_777@hotmail.com

Now, if $\eta > 0$, then

$$\Gamma(\eta + 1) = \eta\Gamma(\eta). \quad (1.1)$$

Equation (1.1) can be used to define $\Gamma(\eta)$ for $\eta < 0$ and $\eta \neq -1, -2, \dots$ and further, this is one of the most important formulas that were satisfied by the Gamma function.

Even though the Gamma function is defined as a locally summable function on the real line by [17]

$$\Gamma(\eta) = \int_0^\infty t^{\eta-1} e^{-t} dt, \quad \eta > 0. \quad (1.2)$$

In the classical sense, $\Gamma(\eta)$ function was not defined for the negative integer thus, there was an open problem to give a satisfactory definition. However, by using the neutral limit, it has been shown in [21] that the Gamma function (1.2) is defined as follows:

$$\Gamma(\eta) = N - \lim_{\varepsilon \rightarrow 0} \int_\varepsilon^\infty t^{\eta-1} e^{-t} dt$$

for $\eta \neq 0, -1, -2, \dots$, and this function can be defined by neutral limit such as

$$\begin{aligned} \Gamma(-n) &= N - \lim_{\varepsilon \rightarrow 0} \int_\varepsilon^\infty t^{-n-1} e^{-t} dt \\ &= \int_1^\infty t^{-n-1} e^{-t} dt \\ &\quad + \int_0^1 t^{-n-1} \left[e^{-t} - \sum_{i=0}^n \frac{(-1)^i}{i!} t^i \right] dt - \sum_{i=0}^{n-1} \frac{(-1)^i}{i!(n-i)}, n \in \mathbb{N}. \end{aligned}$$

It was also proven in [20] the existence of r the derivative of the Gamma function and defined it by equation

$$\begin{aligned} \Gamma^{(r)}(0) &= N - \lim_{\varepsilon \rightarrow 0} \int_\varepsilon^\infty t^{-1} \ln^r t e^{-t} dt \\ &= \int_1^\infty t^{-1} \ln^r t e^{-t} dt + \int_0^1 t^{-1} \ln^r t [e^{-t} - 1] dt \\ \Gamma^{(r)}(-n) &= N - \lim_{\varepsilon \rightarrow 0} \int_\varepsilon^\infty t^{-n-1} \ln^r t e^{-t} dt \end{aligned}$$

R_{ef}

21. B. Fisher, A. Kill, cman and D. Nicholas. On the beta function and the neutrix product of distributions. *Integral Transforms Spec. Funct.*, 7 1-2(1998), 35–42.

$$\begin{aligned}
 &= \int_1^\infty t^{-n-1} \ln^r t e^{-t} dt \\
 &\quad + \int_0^1 t^{-n-1} \ln^r t \left[e^{-t} - \sum_{i=0}^n \frac{(-1)^i}{i!} t^i \right] dt \\
 &\quad - \sum_{i=0}^{n-1} \frac{(-1)^i}{i!} r! (n-i)^{-r-1}
 \end{aligned}$$

for $r \in \mathbb{N}_0$ and $n \in \mathbb{N}$. Also,

$$\Gamma(-r) = \frac{(-1)^r}{r!} (r) - \frac{(-1)^r}{r!} \gamma$$

for $r = 1, 2, \dots$, where

$$(r) = \sum_{i=1}^r \frac{1}{i}.$$

Thus, the definition can be extended to the whole real line where,

$$\Gamma(0) = \Gamma'(1) = -\gamma,$$

where γ denotes Euler's constant, see [22].

For a function $f: V \subset \mathbb{R} \rightarrow \mathbb{R}$ with $0 \in V$, the fractional derivative of order α is defined by:

$$\frac{d^\alpha}{dt^\alpha} f(t) = \frac{1}{\Gamma(-\alpha)} \int_0^t \frac{f(s) - f(0)}{(t-s)^{1+\alpha}} ds, \quad \alpha < 0 \quad (1.3)$$

$$\frac{d^\alpha}{dt^\alpha} f(t) = \frac{1}{\Gamma(n-\alpha)} \frac{d^n}{dt^n} \int_0^t \frac{f(s) - f(0)}{(t-s)^{\alpha-n+1}} ds, \quad \alpha > 0 \quad (1.4)$$

where n is the first integer greater than or equal to α .

The relation (1.3) gives a fractional integral and (1.4) gives a fractional derivative.

We express some of the operators of fractional derivatives, see for example, [4, 7, 9, 10, 12, 16].

1. $\frac{d^\alpha}{dt^\alpha} t^\gamma = \frac{\Gamma(1+\gamma)}{\Gamma(1+\gamma+\alpha)} t^{\gamma-\alpha}, \quad \alpha \in \mathbb{R} \text{ or } (\alpha \in \mathbb{C}) \text{ and } 1+\gamma \neq 0, -1, \dots, -n,$

2. $\frac{d^n}{dt^n} \frac{d^\alpha}{dt^\alpha} f(t) = \frac{d^{n+\alpha}}{dt^{n+\alpha}} f(t), \quad n \in \mathbb{N},$
3. $\frac{d^\alpha}{dt^\alpha} (f_1(t) + f_2(t)) = \frac{d^\alpha}{dt^\alpha} f_1(t) + \frac{d^\alpha}{dt^\alpha} f_2(t),$
4. $\frac{d^\alpha}{dt^\alpha} (Cf(t)) = C \frac{d^\alpha}{dt^\alpha} f(t),$ where C is a constant,
5. $\frac{d^\alpha}{dt^\alpha} f(\beta t) = \beta^\alpha \frac{d^\alpha}{[d(\beta t)]^\alpha} f(\beta t).$

It is well known that fractional calculus is an essential and advantageous branch of mathematics, having a broad range of applications at almost every department of sciences. Techniques of fractional calculus have been employed in the modeling of many different phenomena in engineering, physics, and mathematics. The problem in fractional calculus is not only essential but also quite challenging, which usually involves complicated mathematical solution techniques. However, a general solution theory for almost every issue in this area has yet to be established. Each application has developed its approaches and implementations. Consequently, a single standard method for the problems in fractional calculus has not emerged yet. Therefore, finding reliable and efficient solution techniques along with fast implementation methods are significantly essential and still active research areas.

Further, it is also realized that the operators of fractional integration and derivation have physical and geometric interpretations, which streamline along with their utilization for related issues in various fields of science (see [2, 8, 10, 11, 12, 14, 18, 19]). Moreover, the fractional differential calculus on a differential manifold is studied in [2, 3, 4, 6, 13]. Even though fractional calculus is a handy and important topic, however, the research on geometric interpretation and applications are limited, and not many in current literature. Thus, in this study, we focus on the Riemannian curvature tensor, Livi-Civita connection and Bianchi's identity on fractional differentiable manifolds and discuss some related properties. We also give some examples.

II. FRACTIONAL DIFFERENTIAL CALCULUS ON MANIFOLDS

Assume that N be an m -dimensional differential manifold (V, x_i) a local coordinate system on N and $V_0 = \{x \in V : 0 \leq x_i \leq b_i, i = 1, 2, \dots, m\}$ [5].

For a function $f: V_0 \rightarrow \mathbb{R}$, the fractional derivative with respect to x_i :

$$\partial_i^\alpha f(x) = \frac{1}{\Gamma(n-\alpha)} \partial_{x_i}^n \int_0^{x_i} \frac{f(x_1, \dots, x_{i-1}, s, x_{i+1}, \dots, x_m) - f(x_1, \dots, x_{i-1}, 0, x_{i+1}, \dots, x_m)}{(x_i - s)^{\alpha-n+1}} ds,$$

R_{ef}

2. M. Axtell, M.E. Bise, Fractional Calculus Application in Control Systems. IEEE, Nat, Aerospace and Electronics Conf., New York, (1990) 563–566.

where $\partial_{x_i}^n = \frac{\partial}{\partial x_i} \circ \frac{\partial}{\partial x_i} \circ \dots \circ \frac{\partial}{\partial x_i}$ (n times, i is fixed, $\alpha \geq 0$).

For $\alpha \in (0, 1), \gamma > -1$,

$$\partial_i^\alpha (x_i)^\gamma = \frac{\Gamma(1+\gamma)}{\Gamma(1+\gamma-\alpha)} \quad ; \partial_i^\alpha = \delta_i^j.$$

A fractional vector field $V \subset N$ is an object of the form $X^\alpha = X_i^\alpha \partial_i^\alpha$, where $X_i^\alpha \in \mathfrak{S}_V(N)$ $i = 1, \dots, m$.

The fractional vector fields on V and χ_V^α is generated by the operators $\partial_i^\alpha, i = 1, 2, \dots, m$ are denoted by χ_V^α . If $c: x = x(t), t \in I$ is a parameterized curve in U then the fractional tangent vector field of c is given by

$$x^\alpha(t) = \frac{1}{\Gamma(1+\alpha)} \partial_t^\alpha x_i(t) \partial_i^\alpha.$$

A fractional covariant derivative is given by

$$\nabla_{X^\alpha} Y^\alpha = X_i^\alpha (\partial_i^\alpha Y_j^\alpha + \tilde{\Gamma}_{ik}^j Y_k^\alpha) \partial_j^\alpha$$

where $X^\alpha, Y^\alpha \in \chi_U^\alpha$ and $\tilde{\Gamma}_{ik}^j$ the functions defining the coefficients of a fractional linear connection on N . They are determined by the relations

$$\nabla_{\partial_i^\alpha} \partial_k^\alpha = \tilde{\Gamma}_{ik}^j \partial_j^\alpha.$$

Since it is essential to study fractional vector fields on a differentiable manifold N . For $\mathbb{R}^n$, there is an obvious way to do this. Recall that $\chi^\alpha(\mathbb{R}^n)$ denotes the space of fractional differentiable vector fields defined on $\mathbb{R}$. Examples are the fractional vector fields $\frac{\partial^\alpha}{\partial u_1^\alpha}, \dots, \frac{\partial^\alpha}{\partial u_n^\alpha}$ determined by the natural coordinate functions $u_1, \dots, u_n$.

Definition 2.1. Fractional Riemannian metric F on m -dimensional manifold N defines for every point $p \in N$, the scalar product of fractional tangent vectors in the fractional tangent space $T_p^\alpha N$ depending on the point p .

Let $A^\alpha = A_i^\alpha \partial_i^\alpha$ and $B^\alpha = B_j^\alpha \partial_j^\alpha$ any two fractional vectors tangent to the manifold N at the point p with coordinates $x = (x_1, \dots, x_m)$ ($A^\alpha, B^\alpha \in T_p^\alpha N$) the scalar product is equal to

$$\begin{aligned} \langle A^\alpha, B^\alpha \rangle_F |_p &= A_i^\alpha(x) \tilde{g}_{ij}(x) B_j^\alpha(x) \\ &= (A_1^\alpha, \dots, A_n^\alpha) \begin{pmatrix} \tilde{g}_{11} & \dots & \tilde{g}_{1n} \\ \vdots & \dots & \vdots \\ \tilde{g}_{n1} & \dots & \tilde{g}_{nn} \end{pmatrix} \begin{pmatrix} B_1^\alpha \\ \vdots \\ B_n^\alpha \end{pmatrix} \end{aligned}$$

where

1. $F(A^\alpha, B^\alpha) = F(B^\alpha, A^\alpha)$, i.e., $\tilde{g}_{ij} = \tilde{g}_{ji}$ (symmetricity condition).
2. $F(A^\alpha, A^\alpha) > 0$ if $A^\alpha \neq 0$, i.e. $\tilde{g}_{ij} u_i^\alpha u_j^\alpha \geq 0$, $\tilde{g}_{ij} u_i^\alpha u_j^\alpha = 0$ iff $u_1^\alpha = \dots = u_n^\alpha = 0$ (positive definiteness).
3. $F(A^\alpha, B^\alpha) |_{p=x}$, i.e. $\tilde{g}_{ij}(x)$ are smooth function where $0 < \alpha < 1$.

Components of tensor field F in coordinate system are matrix valued functions $\tilde{g}_{ij}(x)$

$$F = \tilde{g}_{ij}(x) d^\alpha x_i \otimes d^\alpha x_j.$$

III. RULE OF TRANSFORMATION FOR ENTRIES OF THE MATRIX $\tilde{g}_{ij}(x)$

$\tilde{g}_{ij}(x)$ - entries of the matrix $\| \tilde{g}_{ij} \|$ are components of tensor field F in a given coordinate system.

How do these components transform under transformation of coordinates $\{x_i\} \rightarrow \{x_{i'}\}$?

$$\begin{aligned} F &= \tilde{g}_{ij} d^\alpha x_i \otimes d^\alpha x_j \\ &= \tilde{g}_{ij} \left(\frac{\partial x_i^\alpha}{\partial x_{i'}^\alpha} d^\alpha x_{i'} \right) \otimes \left(\frac{\partial x_j^\alpha}{\partial x_{j'}^\alpha} d^\alpha x_{j'} \right) \\ &= \frac{\partial x_i^\alpha}{\partial x_{i'}^\alpha} \tilde{g}_{ij} \frac{\partial x_j^\alpha}{\partial x_{j'}^\alpha} d^\alpha x_{i'} \otimes d^\alpha x_{j'} \\ &= \tilde{g}_{i'j'} d^\alpha x_{i'} \otimes d^\alpha x_{j'}. \end{aligned}$$

Hence,

$$\tilde{g}_{i'j'} = \frac{\partial x_i^\alpha}{\partial x_{i'}^\alpha} \tilde{g}_{ij} \frac{\partial x_j^\alpha}{\partial x_{j'}^\alpha}.$$

Example 2.2. Consider $\mathbb{R}^2$ with fractional polar coordinates in the domain $y > 0$, $X = (r^\alpha \cos^\alpha \varphi, r^\alpha \sin^\alpha \varphi)$, then

$$\frac{\partial^\alpha X}{\partial r^\alpha} = (\alpha! \cos^\alpha \varphi, \alpha! \sin^\alpha \varphi).$$

$$\frac{\partial^\alpha X}{\partial \varphi^\alpha} = (\alpha! e^{i\alpha\pi} r^\alpha \sin^\alpha \varphi, \alpha! r^\alpha \cos^\alpha \varphi).$$

$$\tilde{g}_{ij} = \begin{pmatrix} (\alpha!)^2 [\cos^{2\alpha} \varphi + \sin^{2\alpha} \varphi] & (\alpha!)^2 r^\alpha [e^{2\alpha\pi} + 1] \sin^\alpha \varphi \cos^\alpha \varphi \\ (\alpha!)^2 r^\alpha [e^{2\alpha\pi} + 1] \sin^\alpha \varphi \cos^\alpha \varphi & (\alpha!)^2 r^{2\alpha} [e^{2\alpha\pi} \sin^{2\alpha} \varphi + \cos^{2\alpha} \varphi] \end{pmatrix}$$

We have that

$$\begin{aligned} F &= (\alpha!)^2 [\cos^{2\alpha} \varphi + \sin^{2\alpha} \varphi] (dr^\alpha)^2 \\ &\quad + 2(\alpha!)^2 r^\alpha [e^{i\alpha\pi} + 1] \sin^\alpha \varphi \cos^\alpha \varphi dr^\alpha d\varphi^\alpha \\ &\quad + (\alpha!)^2 r^{2\alpha} [e^{2i\alpha\pi} \sin^{2\alpha} \varphi + \cos^{2\alpha} \varphi] (d\varphi^\alpha)^2. \end{aligned}$$

Notice that , as expected, when $\alpha = 1$, one recovers the classical formula.

$$F = (dr)^2 + r^2 (d\varphi)^2.$$

Table 1: $C = \cos \varphi, S = \sin \varphi$

$\alpha =$	0.1	0.2
$\tilde{g}_{rr}$	$0.905[C^{0.2} + S^{0.2}]$	$0.843[C^{0.4} + S^{0.4}]$
$\tilde{g}_{r\varphi} = \tilde{g}_{\varphi r}$	$0.905r^{0.1}[i+1]S^{0.2}C^{0.2}$	0
$\tilde{g}_{\varphi\varphi}$	$0.905r^{0.2}[-S^{0.2} - C^{0.2}]$	$0.843r^{0.4}[-S^{0.4} + C^{0.4}]$
F	$0.905[C^{0.2} + S^{0.2}]dr^{0.2}$ $+1.8101r^{0.1}[i+1]S^{0.2}C^{0.2}(dr)^{.1}(d\varphi)^{.1}$ $+0.905r^{0.2}[-S^{0.2} - C^{0.2}](d\varphi)^{.2}$	$0.843[C^{0.4} + S^{0.4}]dr^{0.4}$ $+0.843r^{0.4}[-S^{0.4} + C^{0.4}](d\varphi)^{0.4}$

 Table 2: $C = \cos \varphi, S = \sin \varphi$

$\alpha =$	0.3	0.4
$\tilde{g}_{rr}$	$0.805[C^{0.6} + S^{0.6}]$	$0.787[C^{0.8} + S^{0.8}]$
$\tilde{g}_{r\varphi} = \tilde{g}_{\varphi r}$	$0.805r^{0.3}[[i+1]S^{0.3}C^{0.3}]$	$1.574r^{0.4}S^{0.3}C^{0.3}$
$\tilde{g}_{\varphi\varphi}$	$0.805r^{0.6}[-S^{0.6} + C^{0.6}]$	$0.787r^{0.8}[S^{0.8} + C^{0.8}]$
F	$0.805[C^{0.6} + S^{0.6}](dr)^{0.6}$ $+1.61r^{0.3}[[i+1]S^{0.3}C^{0.3}](dr)^{.3}(d\varphi)^{.3}$ $+0.805r^{0.6}[-S^{0.6} + C^{0.6}](d\varphi)^{.6}$	$0.787[C^{0.8} + S^{0.8}](dr)^{0.8}$ $+3.148r^{0.4}S^{0.4}C^{0.4}dr^{.4}d\varphi^{.4}$ $+0.787r^{0.8}[S^{0.8} + C^{0.8}](d\varphi)^{.8}$

Remark 2.1. Let N is an m -dimensional Riemannian manifold with fractional metric tensor $\tilde{g}$, then we shall denote the fractional derivatives of the elements of tensor $\tilde{g}$ as follows:

$$\tilde{g}_{ij,k} = \frac{\partial^\alpha}{\partial x_k^\alpha} \tilde{g}_{ij},$$

and

$$\begin{aligned} \tilde{g}_{ij,kl} &= \frac{\partial^\alpha}{\partial x_l^\alpha} \frac{\partial^\alpha}{\partial x_k^\alpha} \tilde{g}_{ij} \\ &= \frac{\partial^{2\alpha}}{\partial x_l^\alpha \partial x_k^\alpha} \tilde{g}_{ij}, i, j, k, l = 1, \dots, n. \end{aligned}$$

Definition 2.3. Asymmetric fractional connection is called Levi-Civita fractional connection if it is compatible with metric, i.e., if it preserves the scalar product:

$$\partial_{X^\alpha} \langle Y^\alpha, Z^\alpha \rangle = \langle \nabla_{X^\alpha}^\alpha Y^\alpha, Z^\alpha \rangle + \langle Y^\alpha, \nabla_{X^\alpha}^\alpha Z^\alpha \rangle$$

for arbitrary fractional vector fields X^α, Y^α , and Z^α .

In local coordinates Christoffel symbols of Levi-Civita fractional connection are given by:

$$\tilde{\Gamma}_{ij}^k = \frac{1}{2} \tilde{g}^{kl} (\partial_j^\alpha \tilde{g}_{il} + \partial_i^\alpha \tilde{g}_{lj} - \partial_l^\alpha \tilde{g}_{ij}).$$

Table 4: $C = \cos \varphi, S = \sin \varphi$

$\alpha =$	0.5	0.6
$\tilde{g}_{rr}$	$0.785[C + S]$	$0.798[C^{1.2} + S^{1.2}]$
$\tilde{g}_{r\varphi} = \tilde{g}_{\varphi r}$	$0.785r^{0.5}[i + 1]S^{0.5}C^{0.5}$	0
$\tilde{g}_{\varphi\varphi}$	$0.785r[-S + C]$	$0.798r^{1.2}[S^{1.2} + C^{1.2}]$
F	$0.785[C + S]r(dr)$ $+1.57r^{0.5}[i + 1]S^{0.5}C^{0.5}(dr)^{0.5}(d\varphi)^{0.5}$ $0.785r[-S + C]d\varphi$	$0.798[C^{1.2} + S^{1.2}](dr)^{1.2}$ $+$ $+0.798r^{1.2}[S^{1.2} + C^{1.2}](d\varphi)^{1.2}$

 Table 5: $C = \cos \varphi, S = \sin \varphi$

$\alpha =$	0.7	0.8
$\tilde{g}_{rr}$	$0.826[C^{1.4} + S^{1.4}]$	$0.867[C^{1.6} + S^{1.6}]$
$\tilde{g}_{r\varphi} = \tilde{g}_{\varphi r}$	$0.826r^{0.7}[i + 1]S^{0.7}C^{0.7}$	$1.734r^{0.8}S^{0.8}C^{0.8}$
$\tilde{g}_{\varphi\varphi}$	$0.826r^{1.4}[-S^{1.4} + C^{1.4}]$	$0.867r^{1.6}[S^{1.6} + C^{1.6}]$
F	$0.826[C^{1.4} + S^{1.4}](dr)^{1.4}$ $+1.652r^{0.7}[i + 1]S^{0.7}C^{0.7}(dr)^{0.7}(d\varphi)^{0.7}$ $+0.826r^{1.4}[-S^{1.4} + C^{1.4}](d\varphi)^{1.4}$	$0.867[C^{1.6} + S^{1.6}](dr)^{1.6}$ $+3.468r^{0.8}S^{0.8}C^{0.8}dr^{0.8}d\varphi^{0.8}$ $+0.867r^{1.6}[S^{1.6} + C^{1.6}](d\varphi)^{1.6}$

$\alpha =$	0.9	1
$\tilde{g}_{rr}$	$0.925[C^{1.8} + S^{1.8}]$	1
$\tilde{g}_{r\varphi} = \tilde{g}_{\varphi r}$	$0.925r^{0.9}[i + 1]S^{0.9}C^{0.9}$	0
$\tilde{g}_{\varphi\varphi}$	$0.925r^{1.8}[-S^{1.8} + C^{1.8}]$	r^2
F	$0.925[C^{1.8} + S^{1.8}](dr)^{1.8}$ $+1.85r^{0.9}[i + 1]S^{0.9}C^{0.9}(dr)^{0.9}(d\varphi)^{0.9}$ $+0.925r^{1.8}[-S^{1.8} + C^{1.8}](d\varphi)^{1.8}$	$(dr)^2$ $+$ $r^2(d\varphi)^2$

Proof. Since

$$\partial_j^\alpha e_i = \tilde{\Gamma}_{ij}^m e_m \quad (2.1)$$

$$\tilde{\Gamma}_{ij}^m e_m e_l = (\partial_j^\alpha e_i) e_l \quad (2.2)$$

$$\begin{aligned} \tilde{\Gamma}_{ij}^m g_{ml} &= \partial_j^\alpha (e_i \cdot e_l) - e_i (\partial_j^\alpha e_l) \\ &= \partial_j^\alpha g_{il} - \tilde{\Gamma}_{lj}^m e_m e_i \\ &= \partial_j^\alpha g_{il} - \tilde{\Gamma}_{lj}^m g_{mi}, \end{aligned} \quad (2.3)$$

then

$$\tilde{\Gamma}_{ij}^m g_{ml} + \tilde{\Gamma}_{lj}^m g_{mi} = \partial_j^\alpha g_{il}, \quad (2.4)$$

which implies that

$$\tilde{\Gamma}_{ij}^m \tilde{g}_{ml} + \tilde{\Gamma}_{lj}^m \tilde{g}_{mi} = \partial_j^\alpha \tilde{g}_{il}. \quad (2.5)$$

In this equation, the index m is a dummy, so only the indices i, j , and l are specified. We can cyclically permute these indices to generate two more equations:

$$\tilde{I}_{jl}^m \tilde{g}_{mi} + \tilde{I}_{il}^m \tilde{g}_{mj} = \partial_l^\alpha \tilde{g}_{ji} \quad (2.6)$$

$$\tilde{I}_{li}^m \tilde{g}_{mj} + \tilde{I}_{ji}^m \tilde{g}_{ml} = \partial_i^\alpha \tilde{g}_{lj} \quad (2.7)$$

since $\tilde{I}_{ij}^m = \tilde{I}_{ji}^m$, then

$$\tilde{I}_{lj}^m \tilde{g}_{mi} + \tilde{I}_{il}^m \tilde{g}_{mj} = \partial_l^\alpha \tilde{g}_{ij} \quad (2.8)$$

$$\tilde{I}_{il}^m \tilde{g}_{mj} + \tilde{I}_{ij}^m \tilde{g}_{ml} = \partial_i^\alpha \tilde{g}_{lj}. \quad (2.9)$$

We can now add (2.5) to (2.9) and subtract (2.8) to get

$$2\tilde{I}_{ij}^m \tilde{g}_{ml} = \partial_j^\alpha \tilde{g}_{il} + \partial_i^\alpha \tilde{g}_{lj} - \partial_l^\alpha \tilde{g}_{ij}$$

$$2\tilde{I}_{ij}^m \tilde{g}_{ml} \tilde{g}^{kl} = \tilde{g}^{kl} (\partial_j^\alpha \tilde{g}_{il} + \partial_i^\alpha \tilde{g}_{lj} - \partial_l^\alpha \tilde{g}_{ij})$$

since $\tilde{g}_{ml} \tilde{g}^{kl} = \delta_m^k$, then

$$\tilde{I}_{ij}^k = \frac{1}{2} \tilde{g}^{kl} (\partial_j^\alpha \tilde{g}_{il} + \partial_i^\alpha \tilde{g}_{lj} - \partial_l^\alpha \tilde{g}_{ij}),$$

we can write

$$\tilde{I}_{ij}^k = \frac{1}{2} \tilde{g}^{kl} (\tilde{g}_{il,j} + \tilde{g}_{lj,i} - \tilde{g}_{ij,l}).$$

Example 2.2. For 2-dimensional polar coordinates $X = (r^\alpha \cos^\alpha \varphi, r^\alpha \sin^\alpha \varphi)$.

The metric tensor and its inverse here are:

$$\tilde{g}_{ij} = \begin{pmatrix} (\alpha!)^2 [\cos^{2\alpha} \varphi + \sin^{2\alpha} \varphi] & (\alpha!)^2 r^\alpha [e^{2\alpha\pi} + 1] \sin^\alpha \varphi \cos^\alpha \varphi \\ (\alpha!)^2 r^\alpha [e^{\alpha\pi i} + 1] \sin^\alpha \varphi \cos^\alpha \varphi & (\alpha!)^2 r^{2\alpha} [e^{2\alpha\pi i} \sin^{2\alpha} \varphi + \cos^{2\alpha} \varphi] \end{pmatrix}$$

$$\tilde{g}^{ij} = \begin{pmatrix} A^{-1}(\alpha!)^2 r^{2\alpha} [e^{2\alpha\pi i} \sin^{2\alpha} \varphi + \cos^{2\alpha} \varphi] & -A^{-1}(\alpha!)^2 r^\alpha [e^{2\alpha\pi} + 1] \sin^\alpha \varphi \cos^\alpha \varphi \\ -A^{-1}(\alpha!)^2 r^\alpha [e^{2\alpha\pi} + 1] \sin^\alpha \varphi \cos^\alpha \varphi & A^{-1}(\alpha!)^2 [\cos^{2\alpha} \varphi + \sin^{2\alpha} \varphi] \end{pmatrix}.$$

where

$$A = (\alpha!)^4 r^{2\alpha} \{ [\cos^{2\alpha} \varphi + \sin^{2\alpha} \varphi] [e^{2\alpha\pi i} \sin^{2\alpha} \varphi + \cos^{2\alpha} \varphi] - [e^{\alpha\pi i} + 1] \sin^{2\alpha} \varphi \cos^{2\alpha} \varphi \}$$

Therefore,

$$\partial_r^\alpha \tilde{g}_{ij} = \begin{pmatrix} 0 & (\alpha!)^3 [e^{\alpha\pi i} + 1] \sin^\alpha \varphi \cos^\alpha \varphi \\ (\alpha!)^3 [e^{\alpha\pi i} + 1] \sin^\alpha \varphi \cos^\alpha \varphi & (\alpha!)(2\alpha)! r^\alpha [e^{2\alpha\pi i} \sin^{2\alpha} \varphi + \cos^{2\alpha} \varphi] \end{pmatrix}$$

$$\partial_\varphi^\alpha \tilde{g}_{ij} = \begin{pmatrix} (\alpha!)(2\alpha)! [e^{\alpha\pi i} + 1] \cos^\alpha \varphi \sin^\alpha \varphi & (\alpha!)^3 r^\alpha [e^{\alpha\pi i} + 1] [e^{\alpha\pi i} \sin^{2\alpha} \varphi + \cos^{2\alpha} \varphi] \\ (\alpha!)^3 r^\alpha [e^{\alpha\pi i} + 1] [e^{\alpha\pi i} \sin^{2\alpha} \varphi + \cos^{2\alpha} \varphi] & (\alpha!)(2\alpha)! r^{2\alpha} [e^{\alpha\pi i} + 1] \sin^\alpha \varphi \cos^\alpha \varphi \end{pmatrix}.$$

Then ,

$$\tilde{I}_{rr}^r = \frac{1}{2} \tilde{g}^{rl} (\partial_r^\alpha \tilde{g}_{rl} + \partial_r^\alpha \tilde{g}_{lr} - \partial_l^\alpha \tilde{g}_{rr})$$

$$= -A^{-1}(\alpha!)^5 r^\alpha [e^{\alpha\pi i} + 1]^2 \sin^{2\alpha} \varphi \cos^{2\alpha} \varphi,$$

$$\tilde{I}_{r\varphi}^r = \frac{1}{2} \tilde{g}^{rl} (\partial_\varphi^\alpha \tilde{g}_{rl} + \partial_r^\alpha \tilde{g}_{l\varphi} - \partial_l^\alpha \tilde{g}_{r\varphi})$$

$$= 0$$

$$= \tilde{I}_{\varphi r}^r,$$

$$\tilde{I}_{\varphi\varphi}^r = \frac{1}{2} \tilde{g}^{rl} (\partial_\varphi^\alpha \tilde{g}_{\varphi l} + \partial_\varphi^\alpha \tilde{g}_{l\varphi} - \partial_l^\alpha \tilde{g}_{\varphi\varphi})$$

$$= (\alpha!)^2 r^\alpha (2A)^{-1} \{ [e^{2\alpha\pi i} \sin^{2\alpha} \varphi + \cos^{2\alpha} \varphi] [2(\alpha!)^3 r^\alpha [e^{\alpha\pi i} + 1] [e^{\alpha\pi i} \sin^{2\alpha} \varphi + \cos^{2\alpha} \varphi] \\ - (\alpha!)(2\alpha)! r^\alpha [e^{2\alpha\pi i} \sin^{2\alpha} \varphi + \sin^{2\alpha} \varphi] - (\alpha!)(2\alpha)! r^\alpha [e^{\alpha\pi i} + 1] \sin^{2\alpha} \varphi \cos^{2\alpha} \varphi \},$$

$$\tilde{I}_{rr}^\varphi = \frac{1}{2} \tilde{g}^{\varphi l} (\partial_\varphi^\alpha \tilde{g}_{rl} + \partial_r^\alpha \tilde{g}_{lr} - \partial_l^\alpha \tilde{g}_{rr})$$

$$= -(\alpha!)^2 (2A)^{-1} \{ (\alpha!)(2\alpha)! r^\alpha [e^{\alpha\pi i} + 1]^2 \sin^{2\alpha} \varphi \cos^{2\alpha} \varphi + [\cos^{2\alpha} \varphi + \sin^{2\alpha} \varphi] [e^{\alpha\pi i} + 1] \\ \times [(\alpha!)^3 r^\alpha [e^{\alpha\pi i} \sin^{2\alpha} \varphi + \cos^{2\alpha} \varphi] + (\alpha!)^3 \sin^\alpha \varphi \cos^\alpha \varphi - (\alpha!)(2\alpha)! \cos^\alpha \varphi \sin^\alpha \varphi \},$$

$$\tilde{I}_{r\varphi}^\varphi = \frac{1}{2} \tilde{g}^{\varphi l} (\partial_\varphi^\alpha \tilde{g}_{rl} + \partial_r^\alpha \tilde{g}_{l\varphi} - \partial_l^\alpha \tilde{g}_{r\varphi})$$

$$= (\alpha!)^2 (2A)^{-1} \{ -(\alpha!)(2\alpha)!^2 r^\alpha [e^{\alpha\pi i} + 1] \cos^{2\alpha} \varphi \sin^{2\alpha} \varphi \\ + (\alpha!)(2\alpha)! r^\alpha [\cos^{2\alpha} \varphi + \sin^{2\alpha} \varphi] [e^{2\alpha\pi i} \sin^{2\alpha} \varphi + \cos^{2\alpha} \varphi] \}$$

$$= \tilde{I}_{\varphi r}^\varphi,$$

$$\tilde{I}_{\varphi\varphi}^\varphi = \frac{1}{2} \tilde{g}^{\varphi l} (\partial_\varphi^\alpha \tilde{g}_{\varphi l} + \partial_\varphi^\alpha \tilde{g}_{l\varphi} - \partial_l^\alpha \tilde{g}_{\varphi\varphi})$$

$$= (\alpha!)^2 r^{2\alpha} (2A)^{-1} \{ -[e^{\alpha\pi i} + 1] \sin^\alpha \varphi \cos^\alpha \varphi [2(\alpha!)^3 [e^{\alpha\pi i} + 1] [e^{\alpha\pi i} \sin^{2\alpha} \varphi + \cos^{2\alpha} \varphi] \\ - (\alpha!)(2\alpha)! [e^{2\alpha\pi i} \sin^{2\alpha} \varphi + \cos^{2\alpha} \varphi] + (\alpha!)(2\alpha)! [e^{\alpha\pi i} + 1] [\sin^{2\alpha} \varphi + \cos^{2\alpha} \varphi] \sin^\alpha \varphi \cos^\alpha \varphi \}.$$

IV. FRACTIONAL CURVATURE

Definition 3.1. The fractional curvature $\tilde{R}$ of order α of a Riemannian manifold N is a correspondence that associates to every pair $X^\alpha, Y^\alpha \in \chi^\alpha$ a mapping $\tilde{R}(X^\alpha, Y^\alpha): \chi^\alpha(N) \times \chi^\alpha(N) \rightarrow \chi^\alpha(N)$ given by

$$\tilde{R}(X^\alpha, Y^\alpha)Z^\alpha = \nabla_{X^\alpha}^\alpha \nabla_{Y^\alpha}^\alpha Z^\alpha - \nabla_{Y^\alpha}^\alpha \nabla_{X^\alpha}^\alpha Z^\alpha - \nabla_{[X^\alpha, Y^\alpha]}^\alpha Z^\alpha,$$

where $Z^\alpha \in \chi^\alpha$ and ∇^α is the fractional Riemannian connection.

Remark 3.1.

$$\begin{aligned}
 \tilde{R}(X^\alpha, Y^\alpha)Z^\alpha &= \nabla_{X^\alpha}^\alpha \nabla_{Y^\alpha}^\alpha Z^\alpha - \nabla_{Y^\alpha}^\alpha \nabla_{X^\alpha}^\alpha Z^\alpha - \nabla_{[X^\alpha, Y^\alpha]}^\alpha Z^\alpha \\
 &= -(\nabla_{Y^\alpha}^\alpha \nabla_{X^\alpha}^\alpha Z^\alpha - \nabla_{X^\alpha}^\alpha \nabla_{Y^\alpha}^\alpha Z^\alpha - \nabla_{[Y^\alpha, X^\alpha]}^\alpha Z^\alpha) \\
 &= -\tilde{R}(Y^\alpha, X^\alpha)Z^\alpha.
 \end{aligned}$$

Proposition 3.2. *The fractional curvature $\tilde{R}$ of a Riemannian manifold has the following properties:*

1. $\tilde{R}$ is bilinear in $\chi^\alpha(N) \times \chi^\alpha(N)$, that is,

$$\tilde{R}(fX^\alpha + gY^\alpha, Z^\alpha)W^\alpha = f\tilde{R}(X^\alpha, Z^\alpha)W^\alpha + g\tilde{R}(Y^\alpha, Z^\alpha)W^\alpha,$$

$$\tilde{R}(X^\alpha, fY^\alpha + gZ^\alpha)W^\alpha = f\tilde{R}(X^\alpha, Y^\alpha)W^\alpha + g\tilde{R}(X^\alpha, Z^\alpha)W^\alpha,$$

where $f, g \in \mathfrak{S}(M)$, $X^\alpha, Y^\alpha, Z^\alpha, W^\alpha \in \chi^\alpha(N)$

2. For any $X^\alpha, Y^\alpha \in \chi^\alpha(N)$, $\tilde{R}(X^\alpha, Y^\alpha)$ is linear

$$\tilde{R}(X^\alpha, Y^\alpha)(Z^\alpha + W^\alpha) = \tilde{R}(X^\alpha, Y^\alpha)Z^\alpha + \tilde{R}(X^\alpha, Y^\alpha)W^\alpha,$$

$$\tilde{R}(X^\alpha, Y^\alpha)(fZ^\alpha) = f\tilde{R}(X^\alpha, Y^\alpha)Z^\alpha,$$

where $Z^\alpha, W^\alpha \in \chi^\alpha(N)$

Proof. 1.

$$\begin{aligned}
 &\tilde{R}(fX^\alpha + gY^\alpha, Z^\alpha)W^\alpha \\
 &= \nabla_{fX^\alpha + gY^\alpha}^\alpha \nabla_{Z^\alpha}^\alpha W^\alpha - \nabla_{Z^\alpha}^\alpha \nabla_{fX^\alpha + gY^\alpha}^\alpha W^\alpha - \nabla_{[fX^\alpha + gY^\alpha, Z^\alpha]}^\alpha W^\alpha \\
 &= (f \nabla_{X^\alpha}^\alpha + g \nabla_{Y^\alpha}^\alpha) \nabla_{Z^\alpha}^\alpha W^\alpha - \nabla_{Z^\alpha}^\alpha (f \nabla_{X^\alpha}^\alpha W^\alpha + g \nabla_{Y^\alpha}^\alpha W^\alpha) \\
 &\quad - \nabla_{f[X^\alpha, Z^\alpha] + g[Y^\alpha, Z^\alpha] - (Z^\alpha f)X^\alpha - (Z^\alpha g)Y^\alpha}^\alpha W^\alpha \\
 &= f \nabla_{X^\alpha}^\alpha \nabla_{Z^\alpha}^\alpha W^\alpha + g \nabla_{Y^\alpha}^\alpha \nabla_{Z^\alpha}^\alpha W^\alpha - (Z^\alpha f) \nabla_{X^\alpha}^\alpha W^\alpha \\
 &\quad - f \nabla_{Z^\alpha}^\alpha \nabla_{X^\alpha}^\alpha W^\alpha - (Z^\alpha g) \nabla_{Y^\alpha}^\alpha W^\alpha - g \nabla_{Z^\alpha}^\alpha \nabla_{Y^\alpha}^\alpha W^\alpha - f \nabla_{[X^\alpha, Z^\alpha]}^\alpha W^\alpha \\
 &\quad - g \nabla_{[Y^\alpha, Z^\alpha]}^\alpha W^\alpha + (Z^\alpha f) \nabla_{X^\alpha}^\alpha W^\alpha + (Z^\alpha g) \nabla_{Y^\alpha}^\alpha W^\alpha \\
 &= f(\nabla_{X^\alpha}^\alpha \nabla_{Z^\alpha}^\alpha W^\alpha - \nabla_{Z^\alpha}^\alpha \nabla_{X^\alpha}^\alpha W^\alpha - \nabla_{[X^\alpha, Z^\alpha]}^\alpha W^\alpha) \\
 &\quad + g(\nabla_{Y^\alpha}^\alpha \nabla_{Z^\alpha}^\alpha W^\alpha - \nabla_{Z^\alpha}^\alpha \nabla_{Y^\alpha}^\alpha W^\alpha - \nabla_{[Y^\alpha, Z^\alpha]}^\alpha W^\alpha) \\
 &= f\tilde{R}(X^\alpha, Z^\alpha)W^\alpha + g\tilde{R}(Y^\alpha, Z^\alpha)W^\alpha.
 \end{aligned}$$

Also,

$$\begin{aligned}
 \tilde{R}(X^\alpha, fY^\alpha + gZ^\alpha)W^\alpha &= -\tilde{R}(fY^\alpha + gZ^\alpha, X^\alpha)W^\alpha \\
 &= -f\tilde{R}(Y^\alpha, X^\alpha)W^\alpha - g\tilde{R}(Z^\alpha, X^\alpha)W^\alpha \\
 &= f\tilde{R}(X^\alpha, Y^\alpha)W^\alpha + g\tilde{R}(X^\alpha, Z^\alpha)W^\alpha.
 \end{aligned}$$

2.

$$\begin{aligned}
 & \tilde{R}(X^\alpha, Y^\alpha)(Z^\alpha + W^\alpha) \\
 &= \nabla_{X^\alpha}^\alpha \nabla_{Y^\alpha}^\alpha (Z^\alpha + W^\alpha) - \nabla_{Y^\alpha}^\alpha \nabla_{X^\alpha}^\alpha (Z^\alpha + W^\alpha) - \nabla_{[X^\alpha, Y^\alpha]}^\alpha (Z^\alpha + W^\alpha) \\
 &= \nabla_{X^\alpha}^\alpha \nabla_{Y^\alpha}^\alpha Z^\alpha + \nabla_{X^\alpha}^\alpha \nabla_{Y^\alpha}^\alpha W^\alpha - \nabla_{Y^\alpha}^\alpha \nabla_{X^\alpha}^\alpha Z^\alpha \\
 &\quad - \nabla_{Y^\alpha}^\alpha \nabla_{X^\alpha}^\alpha W^\alpha - \nabla_{[X^\alpha, Y^\alpha]}^\alpha Z^\alpha - \nabla_{[X^\alpha, Y^\alpha]}^\alpha W^\alpha \\
 &= (\nabla_{X^\alpha}^\alpha \nabla_{Y^\alpha}^\alpha Z^\alpha - \nabla_{Y^\alpha}^\alpha \nabla_{X^\alpha}^\alpha Z^\alpha - \nabla_{[X^\alpha, Y^\alpha]}^\alpha Z^\alpha) \\
 &\quad + (\nabla_{X^\alpha}^\alpha \nabla_{Y^\alpha}^\alpha W^\alpha - \nabla_{Y^\alpha}^\alpha \nabla_{X^\alpha}^\alpha W^\alpha - \nabla_{[X^\alpha, Y^\alpha]}^\alpha W^\alpha) \\
 &= \tilde{R}(X^\alpha, Y^\alpha)Z^\alpha + \tilde{R}(X^\alpha, Y^\alpha)W^\alpha.
 \end{aligned}$$

Also,

$$\begin{aligned}
 & \tilde{R}(X^\alpha, Y^\alpha)(fZ^\alpha) \\
 &= \nabla_{X^\alpha}^\alpha \nabla_{Y^\alpha}^\alpha (fZ^\alpha) - \nabla_{Y^\alpha}^\alpha \nabla_{X^\alpha}^\alpha (fZ^\alpha) - \nabla_{[X^\alpha, Y^\alpha]}^\alpha (fZ^\alpha) \\
 &= \nabla_{X^\alpha}^\alpha ((Y^\alpha f)Z^\alpha + f \nabla_{Y^\alpha}^\alpha Z^\alpha) - \nabla_{Y^\alpha}^\alpha ((X^\alpha f)Z^\alpha + f \nabla_{X^\alpha}^\alpha Z^\alpha) \\
 &\quad - ([X^\alpha, Y^\alpha]f)Z^\alpha + f \nabla_{[X^\alpha, Y^\alpha]}^\alpha Z^\alpha \\
 &= X^\alpha(Y^\alpha f)Z^\alpha + (Y^\alpha f) \nabla_{X^\alpha}^\alpha Z^\alpha + (X^\alpha f) \nabla_{Y^\alpha}^\alpha Z^\alpha + f \nabla_{X^\alpha}^\alpha \nabla_{Y^\alpha}^\alpha Z^\alpha \\
 &\quad - Y^\alpha(X^\alpha f)Z^\alpha + (X^\alpha f) \nabla_{Y^\alpha}^\alpha Z^\alpha + (Y^\alpha f) \nabla_{X^\alpha}^\alpha Z^\alpha + f \nabla_{Y^\alpha}^\alpha \nabla_{X^\alpha}^\alpha Z^\alpha \\
 &\quad - ([X^\alpha, Y^\alpha]f)Z^\alpha - f \nabla_{[X^\alpha, Y^\alpha]}^\alpha Z^\alpha \\
 &= ([X^\alpha, Y^\alpha]f)Z^\alpha + f(\nabla_{X^\alpha}^\alpha \nabla_{Y^\alpha}^\alpha Z^\alpha - \nabla_{Y^\alpha}^\alpha \nabla_{X^\alpha}^\alpha Z^\alpha - \nabla_{[X^\alpha, Y^\alpha]}^\alpha Z^\alpha) - ([X^\alpha, Y^\alpha]f)Z^\alpha \\
 &= f\tilde{R}(X^\alpha, Y^\alpha)Z^\alpha.
 \end{aligned}$$

Proposition 3.3 (Bianchi Fractional Identity).

$$\tilde{R}(X^\alpha, Y^\alpha)Z^\alpha + \tilde{R}(Y^\alpha, Z^\alpha)X^\alpha + \tilde{R}(Z^\alpha, X^\alpha)Y^\alpha = 0.$$

Proof.

$$\begin{aligned}
 & \tilde{R}(X^\alpha, Y^\alpha)Z^\alpha + \tilde{R}(Y^\alpha, Z^\alpha)X^\alpha + \tilde{R}(Z^\alpha, X^\alpha)Y^\alpha \\
 &= \nabla_{X^\alpha}^\alpha \nabla_{Y^\alpha}^\alpha Z^\alpha - \nabla_{Y^\alpha}^\alpha \nabla_{X^\alpha}^\alpha Z^\alpha - \nabla_{[X^\alpha, Y^\alpha]}^\alpha Z^\alpha \\
 &\quad + \nabla_{Y^\alpha}^\alpha \nabla_{Z^\alpha}^\alpha X^\alpha - \nabla_{Z^\alpha}^\alpha \nabla_{Y^\alpha}^\alpha X^\alpha - \nabla_{[Y^\alpha, Z^\alpha]}^\alpha X^\alpha \\
 &\quad + \nabla_{Z^\alpha}^\alpha \nabla_{X^\alpha}^\alpha Y^\alpha - \nabla_{X^\alpha}^\alpha \nabla_{Z^\alpha}^\alpha Y^\alpha - \nabla_{[Z^\alpha, X^\alpha]}^\alpha Y^\alpha \\
 &= \nabla_{X^\alpha}^\alpha [Y^\alpha, Z^\alpha] + \nabla_{Y^\alpha}^\alpha [Z^\alpha, X^\alpha] + \nabla_{Z^\alpha}^\alpha [X^\alpha, Y^\alpha] \\
 &\quad - \nabla_{[X^\alpha, Y^\alpha]}^\alpha Z^\alpha - \nabla_{[Y^\alpha, Z^\alpha]}^\alpha X^\alpha - \nabla_{[Z^\alpha, X^\alpha]}^\alpha Y^\alpha \\
 &= [X^\alpha, [Y^\alpha, Z^\alpha]] + [Y^\alpha, [Z^\alpha, X^\alpha]] + [Z^\alpha, [X^\alpha, Y^\alpha]] = 0.
 \end{aligned}$$

Notes

In local coordinates

$$\tilde{R}(\partial_i^\alpha, \partial_j^\alpha) \partial_k^\alpha = \tilde{R}_{ijk}^l \partial_l^\alpha,$$

and

$$\begin{aligned} \tilde{R}_{ijkm} &= \left\langle \tilde{R}(\partial_i^\alpha, \partial_j^\alpha) \partial_k^\alpha, \partial_m^\alpha \right\rangle \\ &= \left\langle \tilde{R}_{ijk}^l \partial_l^\alpha, \partial_m^\alpha \right\rangle \\ &= \tilde{R}_{ijk}^l \langle \partial_l^\alpha, \partial_m^\alpha \rangle \\ &= \tilde{R}_{ijk}^l \tilde{g}_{ml}. \end{aligned}$$

The fractional Riemannian curvature tensor acts on fractional vector fields as follows:

$$\tilde{R}(X^\alpha, Y^\alpha, Z^\alpha, W^\alpha) = \left\langle \tilde{R}(X^\alpha, Y^\alpha) Z^\alpha, W^\alpha \right\rangle.$$

Proposition 3.4. 1. $\tilde{R}_{ijkl} + \tilde{R}_{jkil} + \tilde{R}_{kijl} = 0$.

$$2. \quad \tilde{R}_{ijkl} = -\tilde{R}_{ikjl}.$$

$$3. \quad \tilde{R}_{ijkl} = -\tilde{R}_{ijlk}.$$

$$4. \quad \tilde{R}_{ijkl} = \tilde{R}_{klij}.$$

Proof. 1. is just the Bianchi fractional identity again.

2.

$$\begin{aligned} \tilde{R}_{ijkl} &= \left\langle \tilde{R}(\partial_i^\alpha, \partial_j^\alpha) \partial_k^\alpha, \partial_l^\alpha \right\rangle \\ &= \left\langle -\tilde{R}(\partial_j^\alpha, \partial_i^\alpha) \partial_k^\alpha, \partial_l^\alpha \right\rangle \\ &= -\left\langle \tilde{R}(\partial_j^\alpha, \partial_i^\alpha) \partial_k^\alpha, \partial_l^\alpha \right\rangle \\ &= -\tilde{R}_{jikl}. \end{aligned}$$

3. is equivalent to $\tilde{R}_{ijkk} = 0$, whose proof follows:

$$\begin{aligned} \tilde{R}_{ijkk} &= \left\langle \tilde{R}(\partial_i^\alpha, \partial_j^\alpha) \partial_k^\alpha, \partial_k^\alpha \right\rangle \\ &= \left\langle \nabla_{\partial_i^\alpha}^\alpha \nabla_{\partial_j^\alpha}^\alpha \partial_k^\alpha - \nabla_{\partial_j^\alpha}^\alpha \nabla_{\partial_i^\alpha}^\alpha \partial_k^\alpha - \nabla_{[\partial_i^\alpha, \partial_j^\alpha]}^\alpha \partial_k^\alpha, \partial_k^\alpha \right\rangle, \end{aligned}$$

but

$$\left\langle \nabla_{\partial_j^\alpha}^\alpha \nabla_{\partial_i^\alpha}^\alpha \partial_k^\alpha, \partial_k^\alpha \right\rangle = \partial_j^\alpha \left\langle \nabla_{\partial_i^\alpha}^\alpha \partial_k^\alpha, \partial_k^\alpha \right\rangle - \left\langle \nabla_{\partial_i^\alpha}^\alpha \partial_k^\alpha, \nabla_{\partial_j^\alpha}^\alpha \partial_k^\alpha \right\rangle,$$

and

$$\left\langle \nabla_{[\partial_i^\alpha, \partial_j^\alpha]}^\alpha \partial_k^\alpha, \partial_k^\alpha \right\rangle = \frac{1}{2} [\partial_i^\alpha, \partial_j^\alpha] \langle \partial_k^\alpha, \partial_k^\alpha \rangle,$$

then

$$\tilde{R}_{ijkk} = \partial_j^\alpha \left\langle \nabla_{\partial_i^\alpha}^\alpha \partial_k^\alpha, \partial_k^\alpha \right\rangle - \partial_i^\alpha \left\langle \nabla_{\partial_j^\alpha}^\alpha \partial_k^\alpha, \partial_k^\alpha \right\rangle + \frac{1}{2} [\partial_i^\alpha, \partial_j^\alpha] \langle \partial_k^\alpha, \partial_k^\alpha \rangle$$

$$\begin{aligned}
 &= \frac{1}{2} \partial_j^\alpha (\partial_i^\alpha \langle \partial_k^\alpha, \partial_k^\alpha \rangle) - \frac{1}{2} \partial_i^\alpha (\partial_j^\alpha \langle \partial_k^\alpha, \partial_k^\alpha \rangle) + \frac{1}{2} [\partial_i^\alpha, \partial_j^\alpha] \langle \partial_k^\alpha, \partial_k^\alpha \rangle \\
 &= -\frac{1}{2} [\partial_i^\alpha, \partial_j^\alpha] \langle \partial_k^\alpha, \partial_k^\alpha \rangle + \frac{1}{2} [\partial_i^\alpha, \partial_j^\alpha] \langle \partial_k^\alpha, \partial_k^\alpha \rangle = 0.
 \end{aligned}$$

4. By Bianchi fractional identity we have

$$\tilde{R}_{ijkl} + \tilde{R}_{jkil} + \tilde{R}_{kijl} = 0$$

$$\tilde{R}_{jkli} + \tilde{R}_{klji} + \tilde{R}_{ljki} = 0$$

$$\tilde{R}_{klij} + \tilde{R}_{likj} + \tilde{R}_{iklj} = 0$$

$$\tilde{R}_{lij k} + \tilde{R}_{ijlk} + \tilde{R}_{jlik} = 0$$

summing the equations above, we obtain

$$2\tilde{R}_{kijl} + 2\tilde{R}_{ljki} = 0,$$

then

$$\tilde{R}_{kijl} = -\tilde{R}_{ljki} = \tilde{R}_{jlik}.$$

Proposition 3.5. *The following expression holds*

$$2\tilde{R}_{ijk m} = \tilde{g}_{jm,ki} + \tilde{g}_{km,ji} - \tilde{g}_{jk,mi} - \tilde{g}_{im,kj} - \tilde{g}_{km,ij} + \tilde{g}_{ik,mj} - 2\tilde{\Gamma}_{jk}^r \tilde{\Gamma}_{im}^s \tilde{g}_{rs} + 2\tilde{\Gamma}_{ik}^r \tilde{\Gamma}_{jm}^s \tilde{g}_{rs}.$$

Proof. From the definition of the Christoffel symbols, $\nabla_{\partial_i}^\alpha \partial_j^\alpha = \tilde{\Gamma}_{ij}^k \partial_k^\alpha$,

$$\begin{aligned}
 2 \langle \nabla_{\partial_i}^\alpha \partial_j^\alpha, \partial_m^\alpha \rangle &= 2 \langle \tilde{\Gamma}_{ij}^k \partial_k^\alpha, \partial_m^\alpha \rangle \\
 &= 2\tilde{\Gamma}_{ij}^k \tilde{g}_{mk} \\
 &= \tilde{g}_{im,j} + \tilde{g}_{jm,i} - \tilde{g}_{ij,m},
 \end{aligned}$$

an appropriate rearrangement of the indices yields the following expression:

$$\begin{aligned}
 2 \langle \nabla_{\partial_j}^\alpha \partial_k^\alpha, \partial_m^\alpha \rangle &= 2 \langle \tilde{\Gamma}_{jk}^i \partial_i^\alpha, \partial_m^\alpha \rangle \\
 &= 2\tilde{\Gamma}_{jk}^i \tilde{g}_{im} \\
 &= \tilde{g}_{jm,k} + \tilde{g}_{km,j} - \tilde{g}_{jk,m}.
 \end{aligned} \tag{3.1}$$

$$\partial_i^\alpha \langle \nabla_{\partial_j}^\alpha \partial_k^\alpha, \partial_m^\alpha \rangle = \langle \nabla_{\partial_i}^\alpha \nabla_{\partial_j}^\alpha \partial_k^\alpha, \partial_m^\alpha \rangle + \langle \nabla_{\partial_j}^\alpha \partial_k^\alpha, \nabla_{\partial_i}^\alpha \partial_m^\alpha \rangle$$

whence, by (3.1)

$$\begin{aligned}
 2 \langle \nabla_{\partial_i}^\alpha \nabla_{\partial_j}^\alpha \partial_k^\alpha, \partial_m^\alpha \rangle + 2 \langle \nabla_{\partial_j}^\alpha \partial_k^\alpha, \nabla_{\partial_i}^\alpha \partial_m^\alpha \rangle &= 2\partial_i^\alpha \langle \nabla_{\partial_j}^\alpha \partial_k^\alpha, \partial_m^\alpha \rangle \\
 &= \partial_i^\alpha (\tilde{g}_{im} g^{il} (\partial_k^\alpha g_{jl} + \partial_j^\alpha g_{kl} - \partial_l^\alpha g_{jk})) \\
 &= \tilde{g}_{jm,ki} + \tilde{g}_{km,ji} - \tilde{g}_{jk,mi}.
 \end{aligned} \tag{3.2}$$

By switching i and j we also have that

$$2 \langle \nabla_{\partial_j}^\alpha \nabla_{\partial_i}^\alpha \partial_k^\alpha, \partial_m^\alpha \rangle + 2 \langle \nabla_{\partial_i}^\alpha \partial_k^\alpha, \nabla_{\partial_j}^\alpha \partial_m^\alpha \rangle = \tilde{g}_{im,kj} + \tilde{g}_{km,ij} - \tilde{g}_{ik,mj}. \tag{3.3}$$

Combining (3.2) and (3.3) yields

$$2 \left\langle \nabla_{\partial_i}^\alpha \nabla_{\partial_j}^\alpha \partial_k^\alpha, \partial_m^\alpha \right\rangle - 2 \left\langle \nabla_{\partial_j}^\alpha \nabla_{\partial_i}^\alpha \partial_k^\alpha, \partial_m^\alpha \right\rangle = \\ \tilde{g}_{jm,ki} + \tilde{g}_{km,ji} - \tilde{g}_{jk,mi} - \tilde{g}_{im,kj} - \tilde{g}_{km,ij} + \tilde{g}_{ik,mj} \\ - 2 \left\langle \nabla_{\partial_j}^\alpha \partial_k^\alpha, \nabla_{\partial_i}^\alpha \partial_m^\alpha \right\rangle + 2 \left\langle \nabla_{\partial_i}^\alpha \partial_k^\alpha, \nabla_{\partial_j}^\alpha \partial_m^\alpha \right\rangle.$$

By definition

$$\tilde{R}(\partial_i^\alpha, \partial_j^\alpha) \partial_k^\alpha = \nabla_{\partial_i}^\alpha \nabla_{\partial_j}^\alpha \partial_k^\alpha - \nabla_{\partial_j}^\alpha \nabla_{\partial_i}^\alpha \partial_k^\alpha$$

whence

$$2\tilde{R}_{ijkm} = 2 \left\langle \tilde{R}(\partial_i^\alpha, \partial_j^\alpha) \partial_k^\alpha, \partial_m^\alpha \right\rangle \\ = 2 \left\langle \nabla_{\partial_i}^\alpha \nabla_{\partial_j}^\alpha \partial_k^\alpha, \partial_m^\alpha \right\rangle - 2 \left\langle \nabla_{\partial_j}^\alpha \nabla_{\partial_i}^\alpha \partial_k^\alpha, \partial_m^\alpha \right\rangle,$$

so we have proven that

$$2\tilde{R}_{ijkm} = \\ \tilde{g}_{jm,ki} + \tilde{g}_{km,ji} - \tilde{g}_{jk,mi} - \tilde{g}_{im,kj} - \tilde{g}_{km,ij} + \tilde{g}_{ik,mj} \\ - 2 \left\langle \nabla_{\partial_j}^\alpha \partial_k^\alpha, \nabla_{\partial_i}^\alpha \partial_m^\alpha \right\rangle + 2 \left\langle \nabla_{\partial_i}^\alpha \partial_k^\alpha, \nabla_{\partial_j}^\alpha \partial_m^\alpha \right\rangle.$$

By the definition of the Christoffels,

$$\left\langle \nabla_{\partial_i}^\alpha \partial_k^\alpha, \nabla_{\partial_j}^\alpha \partial_m^\alpha \right\rangle = \left\langle \tilde{\Gamma}_{ik}^r \partial_r^\alpha, \tilde{\Gamma}_{jm}^s \partial_s^\alpha \right\rangle \\ = \tilde{\Gamma}_{ik}^r \tilde{\Gamma}_{jm}^s \langle \partial_r^\alpha, \partial_s^\alpha \rangle \\ = \tilde{\Gamma}_{ik}^r \tilde{\Gamma}_{jm}^s \tilde{g}_{rs}, \\ \left\langle \nabla_{\partial_j}^\alpha \partial_k^\alpha, \nabla_{\partial_i}^\alpha \partial_m^\alpha \right\rangle = \left\langle \tilde{\Gamma}_{jk}^r \partial_r^\alpha, \tilde{\Gamma}_{im}^s \partial_s^\alpha \right\rangle \\ = \tilde{\Gamma}_{jk}^r \tilde{\Gamma}_{im}^s \langle \partial_r^\alpha, \partial_s^\alpha \rangle \\ = \tilde{\Gamma}_{jk}^r \tilde{\Gamma}_{im}^s \tilde{g}_{rs},$$

then

$$2\tilde{R}_{ijkm} = \tilde{g}_{jm,ki} + \tilde{g}_{km,ji} - \tilde{g}_{jk,mi} - \tilde{g}_{im,kj} - \tilde{g}_{km,ij} + \tilde{g}_{ik,mj} - 2\tilde{\Gamma}_{jk}^r \tilde{\Gamma}_{im}^s \tilde{g}_{rs} + 2\tilde{\Gamma}_{ik}^r \tilde{\Gamma}_{jm}^s \tilde{g}_{rs}.$$

Remark 3.6. If $\alpha = 1$, then

$$2R_{ijkm} = g_{jm,ki} + g_{km,ji} - g_{jk,mi} - g_{im,kj} - g_{km,ij} + g_{ik,mj} - 2\Gamma_{jk}^r \Gamma_{im}^s g_{rs} + 2\Gamma_{ik}^r \Gamma_{jm}^s g_{rs}.$$

Since $g_{km,ji} = g_{km,ij}$, then

$$2R_{ijkm} = g_{jm,ki} - g_{jk,mi} - g_{im,kj} + g_{ik,mj} - 2\Gamma_{jk}^r \Gamma_{im}^s g_{rs} + 2\Gamma_{ik}^r \Gamma_{jm}^s g_{rs},$$

then

$$2R_{ijkm} = \tilde{g}_{jm,ki} + \tilde{g}_{km,ji} - \tilde{g}_{jk,mi} - \tilde{g}_{im,kj} - \tilde{g}_{km,ij} + \tilde{g}_{ik,mj} - 2\Gamma_{jk}^r \Gamma_{im}^s g_{rs} + 2\Gamma_{ik}^r \Gamma_{jm}^s g_{rs}.$$

Remark 3.7. For any pair of fractional tangent vectors $X^\alpha, Y^\alpha \in T_p^\alpha N$ we shall denote with $\tilde{\Gamma}(X^\alpha, Y^\alpha)$ the following fractional vector in $T_p^\alpha N$:

$$\tilde{\Gamma}(X^\alpha, Y^\alpha) = \tilde{\Gamma}_{ij}^k X_i^\alpha Y_j^\alpha \partial_k^\alpha.$$

Proposition 3.8. The following expressions hold for any pair $X^\alpha, Y^\alpha \in T_p^\alpha N$:

$$\begin{aligned} 2\tilde{R}(X^\alpha, Y^\alpha, Y^\alpha, X^\alpha) &= \partial_i^\alpha (\tilde{g}_{im} g^{il}) (\partial_k^\alpha g_{jl} + \partial_j^\alpha g_{kl} - \partial_l^\alpha g_{jk}) + \tilde{g}_{im} g^{il} (\partial_i^\alpha \partial_k^\alpha g_{jl} + \partial_i^\alpha \partial_j^\alpha g_{kl} - \partial_i^\alpha \partial_l^\alpha g_{jk}) \\ &\quad - \partial_j^\alpha (\tilde{g}_{jm} g^{jl}) (\partial_k^\alpha g_{il} + \partial_i^\alpha g_{kl} - \partial_l^\alpha g_{ik}) - \tilde{g}_{jm} g^{jl} (\partial_j^\alpha \partial_k^\alpha g_{il} + \partial_j^\alpha \partial_i^\alpha g_{kl} - \partial_j^\alpha \partial_l^\alpha g_{ik}) \\ &\quad + 2 \parallel \tilde{\Gamma}(X^\alpha, Y^\alpha) \parallel^2 - 2 \langle \tilde{\Gamma}(X^\alpha, X^\alpha), \tilde{\Gamma}(Y^\alpha, Y^\alpha) \rangle. \end{aligned}$$

Proof. Since

$$\begin{aligned} \tilde{g}_{rs} X_i^\alpha Y_j^\alpha Y_k^\alpha X_m^\alpha \tilde{\Gamma}_{ik}^r \tilde{\Gamma}_{jm}^s &= \langle X_i^\alpha Y_k^\alpha \tilde{\Gamma}_{ik}^r \partial_r^\alpha, Y_j^\alpha X_m^\alpha \tilde{\Gamma}_{jm}^s \partial_s^\alpha \rangle \\ &= \langle \tilde{\Gamma}(X^\alpha, Y^\alpha), \tilde{\Gamma}(X^\alpha, Y^\alpha) \rangle = \parallel \tilde{\Gamma}(X^\alpha, Y^\alpha) \parallel^2, \end{aligned}$$

and

$$\begin{aligned} \tilde{g}_{rs} X_i^\alpha Y_j^\alpha Y_k^\alpha X_m^\alpha \tilde{\Gamma}_{jk}^r \tilde{\Gamma}_{im}^s &= \langle X_i^\alpha X_m^\alpha \tilde{\Gamma}_{im}^s \partial_s^\alpha, Y_j^\alpha Y_k^\alpha \tilde{\Gamma}_{jk}^r \partial_r^\alpha \rangle \\ &= \langle \tilde{\Gamma}(X^\alpha, X^\alpha), \tilde{\Gamma}(Y^\alpha, Y^\alpha) \rangle, \end{aligned}$$

This completes the proof.

Remark 3.9. If $\alpha = 1$, then

$$\begin{aligned} 2R(X, Y, Y, X) &= g_{jm,ki} - g_{jk,mi} - g_{im,kj} + g_{ik,mj} \\ &\quad + 2 \parallel \Gamma(X, Y) \parallel^2 - 2 \langle \Gamma(X, X), \Gamma(Y, Y) \rangle. \end{aligned}$$

REFERENCES RÉFÉRENCES REFERENCIAS

1. D. Albu, M. Neamtu, D. Opris, The geometry of fractional osculator bundle of higher order and applications. Anal. St. Univ."Al.I.Cuza" Iasi, Proceedings of the Conf. on Diff. Geometry: Lagrange and Hamilton spaces, Sept.3-8, Iasi, (2007) 21–31.
2. M. Axtell, M.E. Bise, Fractional Calculus Application in Control Systems. IEEE, Nat, Aerospace and Electronics Conf., New York, (1990) 563–566.

3. Ding, Yong, and Xinfeng Wu, Fractional integrals on product manifolds. *Potential Analysis*, 304 (2009), 371–383.
4. Guy Jumarie. Riemann-Christoffel Tensor in Differential Geometry of Fractional Order Application to Fractal Space-Time. *Fractals*, 211 (2013).
5. Ion Doru Albu and Dumitru Opris. The geometry of fractional tangent bundle and applications. *Balkan Society of Geometers* (2009).
6. A. Kılı, cman, and W. Saleh. Note on the Fractional Mittag-Leffler Functions by Applying the Modified Riemann-Liouville Derivatives. *Boletim da Sociedade Paranaense de Matematica*, textbf40 2022,1-16
7. K. M. Kolwankar. Studies of Fractal structures and Processes Using Methods of Fractional Calculus; <http://arxiv.org/abs/chaodyn/9811008>.
8. Mehdi Delkhosh. Introduction of Derivatives and Integrals of Fractional order and Its Applications. *Applied Mathematics and physics*, 14 (2013), 103–119.
9. K. S. Miller and B. Ross. An Introduction to the Fractional Calculus and Fractional Differential Equations. John Wiley and Sons, Inc.; New York (1993).
10. K. B. Oldham and J. Spanier. The Fractional Calculus. Academic Press; New York (1974).
11. I. Podlubny. Fractional Order System and Fractional Order Controllers. UEF- 03-94, Inst. Exp. Phys, Slovak Acad. Sci., Kosice (1994).
12. B. Ross. In Fractional Calculus and Its Applications: Lecture Notes in Mathematics; Springer- Verlag; New York; 457 (1975).
13. W. Saleh and A. Kılı cman. Some Remarks on Geometry of Fractional Calculus. *Applied Mathematics & Information Sciences (AMIS)*, 9 3(2015), 1265–1275.
14. W. Saleh and A. Kılı, cman. Some Inequalities For Generalized s-Convex Functions, *JP Journal of Geometry and Topology*, 17 (2015), 63–82.
15. I. N. Sneddon. Special functions of mathematical physics and chemistry. Oliver & Boyd, (1956).
16. M. H. Tavassoli, A. Tavassoli and M. R. Ostad Rahimi. The geometric and physical interpretation of fractional order derivatives of polynomial functions. *Differential Geometry-Dynamical Systems*, 15(2013), 93–104.
17. A. Kılı, cman and B. Fisher. Some Results on the Composition of Singular Distribution. *Applied Mathematics & Information Sciences (AMIS)*, 53 (2011), 597–608.
18. A. Kılı, cman and W. Saleh. Notions of Generalized s-Convex Functions on Fractal Sets. *Journal of Inequalities and Applications*, 20151(2015).

19. A. Kılı, cman and W. Saleh. Some Generalized Hermite-Hadamard Type Integral Inequalities for Generalized s-Convex Functions on Fractal Sets. *Advances in Difference Equations*, 20151(2015).
20. B. Fisher, B. Jolevska-Tuneska and A. Kılı, cman. On defining the incomplete gamma function. *Integral Transforms Spec. Funct.* 14 4(2003), 293–299.
21. B. Fisher, A. Kılı, cman and D. Nicholas. On the beta function and the neutrix product of distributions. *Integral Transforms Spec. Funct.*, 7 1-2(1998), 35–42.
22. B. Fisher and A. Kılı, cman. Results on the Gamma function. *Applied Mathematics & Information Sciences (AMIS)*, 62 (2012), 173–176.

