



GLOBAL JOURNAL OF SCIENCE FRONTIER RESEARCH: H
ENVIRONMENT & EARTH SCIENCE
Volume 22 Issue 2 Version 1.0 Year 2022
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Assessment of the Water Quality of Some Sulfur Springs in the Bashtabia Area within the City of Mosul

By Hadeel Bassam Yahya, Yussra Majed Alshaker & Rawaa Mahmood Hamoshi

Abstract- The research aims to conduct a monthly study of water quality by conducting several physical and chemical tests for sulfur spring water in the Bashtabiya Castle area, which is located on the western bank of the Tigris River in the city of Mosul, and for that, monthly samples were collected from three springs for five months during the period from October The second is 2020 until March 2021.

Keywords: *sulfur spring bashtabia area water quality.*

GJSFR-H Classification: DDC Code: 363.739460973 LCC Code: TD223



Strictly as per the compliance and regulations of:



RESEARCH | DIVERSITY | ETHICS

Assessment of the Water Quality of Some Sulfur Springs in the Bashtabia Area within the City of Mosul

Hadeel Bassam Yahya ^α, Yussra Majed Alshaker ^ο & Rawaa Mahmood Hamoshi ^ρ

Abstract- The research aims to conduct a monthly study of water quality by conducting several physical and chemical tests for sulfur spring water in the Bashtabia Castle area, which is located on the western bank of the Tigris River in the city of Mosul, and for that, monthly samples were collected from three springs for five months during the period from October The second is 2020 until March 2021.

Which included temperature, electrical conductivity, dissolved solids, pH, total hardness, total alkalinity, sulfate ions, phosphates, sodium and potassium, oils and greases, as well as biological tests represented by the total count of bacteria, where the temperature rates ranged between (21.74-22.04) C, the electrical conductivity rates were (5158.59-5521.795) micromoz/cm, while the total dissolved solids rates were between (2579.6-2761 mg/L), and the pH ranged between (6.9-7) during the study period as for the total hardness It was (2080-2220) mg/l in terms of calcium carbonate, as the results showed the total alkalinity rates, which were between (452- 649) mg/l, while sulfate ions averaged (2310.8-2365.3) mg/l, phosphate ions ranged between (0.02154-0.04482) mg/L, and for sodium and potassium ions, their rates ranged between (376-447.6) (31.4-36.2) mg/L, respectively. As for oils, the results showed that their rates were (0.1696-0.1848) mg/L. And the total count of bacteria ranged from the stomach During the study period, between (0.5 - 3.25) cells/ml.

Keywords: sulfur spring bashtabia area water quality.

I. INTRODUCTION

Groundwater is the water that moves and collects in the ground within the geological formations and soil layers, which directly affects the properties of this water (Mohammed, 2015). The amount of water drainage from the springs varies, and the different sources of this water pass through the pores of the soil, which filters its contents and dissolves many salts and elements in it, giving it a high percentage of salinity. That is why it is called mineral water and it is also known as hot water because of its high temperature (Al-Zamili and Al-Asadi, 2014).

Spring water flows from under the surface of the earth and is formed naturally in special water tanks that do not mix with surface water and do not need to make changes or add chemicals. The chemistry of mineral water and its volume change from one place to another

according to the conditions of its formation and the type of formation layers in which this water is stored (Matar and others, 2009). A liter of water (Al Taweel, 2017), the layers of gypsum rocks, and the crater located under the surface of the earth provided a suitable environment for the formation of sulfur supported by special types of anaerobic bacteria, as H_2S gas is released, which interacts with the oxygen dissolved in the water to form sulfuric acid H_2SO_4 , thus forming acidic sulfur water. There are two sources of liberated H_2S gas, the first represented by gas mainly located in the depths and its quantity is small compared to the second source, which is due to bacterial activity, and as products of sulfur formation and reduction resulting from sulfates, the produced gas is oxidized in the presence of special types of bacteria to form sulfuric acid H_2SO_4 (Aswad et al., 2018).

II. STUDY AREA

The study area included a number of sulfur springs, which are located on the western bank of the Tigris River in the city of Mosul, which is close to the Bashtabia Castle. Monthly models were taken to study three hot and cold sulfur springs, Table (1). The study area is characterized by a hot, dry climate in summer and cold and rainy in winter, according to the data of the weather station in Mosul.



Table 1: Shows the location of the sulfur springs near the castle of Bashtabia

Spring name and their locations	Locations	
	E	N
Bashtabia	1 43° 12'00"96	36°35'67"51
	2 43° 12'01"60	36°35'67"63
	3 43° 12'00"78	36°35'67"75

III. SAMPLE COLLECTION

Fifteen water samples were collected at a rate of 3 samples per month from three springs near the castle of Bashtabia (3 samples / spring) for a period of five months during the period from November 2020 to March 2021 using sterile polyethylene bottles of 250 ml capacity for biological tests and a capacity of 2 liters for chemical and physical tests By immersing the bottle

after homogenizing it several times gradually so that its direction faces the flow of water, where the sample was taken and its nozzle was re-sealed tightly inside the water and closed with aluminum foil tightly closed and was kept in a box of iced cork until it was transferred to the laboratory for biological, chemical and physical tests.



Image 1: The first location Image (2) the second location Image (3) the third location

IV. METHODS

a) Physical tests

It includes the measurement of temperature using an alcohol thermometer, measured in degrees, the measurement of total dissolved solids using a T.D.S meter, and electrical conductivity using an E.C meter in micromoz/cm.

b) Chemical tests

The measurement of the pH using a pH meter after regulating it with buffer solutions includes the total hardness by scavenging with Na_2EDTA , the total alkalinity by the guide method, and the determination of sulfate and phosphate by the spectrophotometer, as well as the determination of sodium and potassium ions using a photometric spectrometer. The flame photometer expressed the concentration in mg/l, and finally measured the oils and greases.

c) Biological tests

Bacteriological tests were represented by the total count of bacteria by making a dilution of each sample to a dilution (10^{-5}) with a solution of physiological salt (Normal slain), and placing 1 ml of each of the dilution (10^{-3} , 10^{-4} , 10^{-5}) in a dish. A sterile petri, pour the cooled and sterile Nutrient Agar into the dish and move

the plate in the form of No. 8 until the sample is homogeneous and after solidification of the medium is placed upside down in the incubator and incubated aerobically and anaerobic at a temperature of 37 °C for 24-48 hours.

V. RESULT & DISCUSSION

Water quality depends on the different physical, chemical and biological properties of water, as well as the standard specifications used to measure water quality (Al-Lahibi, 2021).

a) Physical tests

i. Temperature

Temperature is one of the most important indicators of water quality that affects the physical and chemical properties of water (Larnier et al. 2010). The average temperature ranged between (21.74-22.04) °C, the highest rate was in site No. 1 or the lowest was in site 3. The reason for the high temperature of this water is attributed to the chemical reactions that emit heat that accompany the dissolution of the mineral materials that make up gypsum rocks and dolomite when the water of Al-Ayoun passes through it (Al-Hamdani, 2020). As shown in Table No. 2 and Figure.




Figure 2: Averages of the temperature of the studied spring water (°C)

ii. *Electric Conductivity*

Electrical conductivity is the ability of water to transmit electric current and depends on the concentration of dissolved ions in the water and temperature, as it is directly proportional to them (Al-Hamdani, 2020). Its rates reached (5158.59- 5521.795

micromoz/cm. The highest rate was in Site 3 and the lowest was in Site 1, as shown in Table 2 and Figure 3. The reason may be due to the rocky nature of the geological formations of the studied areas (Al-Saffawi, 2019).




Figure 3: The electrical conductivity rate of the studied spring water (micromoz/cm)

iii. *Total Dissolved Solid*

Solids in water represent an important indicator to know the amount of substances or molecules in water, and they include different substances, organic and inorganic substances, or they may be in the form of solid substances dissolved in water consisting of negative ions of compounds combined with ions of positive elements (Bee, 2005). The rates of total dissolved solids were between (2579.6- 2761) mg/L, where the highest value was reached in Site 3 and the lowest in Site 1. The reason may be due to the difference in the geological formations of the layers through which the water passes (Al-Hamdani, 2020). As shown in Table 2 and Figure 4.

b) Chemical Tests

i. pH

It is evident from Table (2) and Figure (5) that the rates were between (6.9-7), the highest rate in site 3 and the lowest in site 2. The reason may be due to the neutralization of the acidity of water and soil with carbonates and bicarbonate salts (Al-Safawi et al.,

2018), and the decrease is due to Some values refer to the biological processes of oxidation and decomposition of organic matter when the oxygen concentration drops to critical levels, thus forming a number of carboxylic and acidic compounds (Sharaky & Abdoun, 2020).




Figure 4: Total Dissolved Solids (TDS) of the studied spring water (mg/L)




Figure 5: Averages of the pH function of the studied springs water

ii. Total Alkalinity

The alkalinity is the amount of negative ions present in water and the resistance to hydrogen ion, and thus it is a measure of water's susceptibility to acidity (Moses & Ishaku, 2020); The results showed the total alkalinity rates, which were between (452-649

mg/L), where the highest value was in site 3 and the lowest value in site 2. The reason for the presence of alkalinity in water is the presence of carbonate, bicarbonate and hydroxide ions. Bicarbonate ion due to the pH values not exceeding 8.3 (Fadipe & Oladepo, 2020). As shown in Figure (6) and Table (2).




Figure 6: The total alkalinity of the studied spring water (mg/L)

iii. Total Hardness

Total hardness is a relative indicator to know the total dissolved solids. Calcium, magnesium, and bicarbonate ions make up the majority of the total dissolved solids. Hardness expresses the water's suitability. Less than that, its effect will be limited (Merdhah and Yassin, 2007; Al-Lahibi, 2021). It was

(2080-2220) mg/L in terms of CaCO_3 , the highest rate was in site 2 and the lowest in site 3 as shown in Table 2 and Figure 7. The reason for the presence of total hardness in water is the positive ions, especially calcium and magnesium ions (Verma et. al., 2018; Al-Hamdani, 2020).



Figure 7: The total hardness of the studied water (mg/L)

iv. Sulphate Ions

It is evident from Table 2 and Figure 8 that the sulfate ion rates ranged between (2310.8-2365.3) mg/L, as the highest rate was in Site 1, while the lowest was in Site 3. The reason for the presence of sulfate ion concentrations is due to the nature of the geological rocks through which the water passes. The dissolution of gypsum rocks and weathering processes is the main

reason for the increase in sulfate concentrations in the study area (Egbueri et. al., 2019), as well as the exposure of some sulfur minerals to oxidative conditions in the presence of water and *Thiobacillus ferrooxidans* bacteria that leads to the liberation of ions of some chemical elements and the formation of acid. Dilute sulfuric in sulfuric water (Duruibe et al. 2007).




Figure 8: Sulfate ion rate for the studied spring water (mg/L)

v. Phosphate Ions

The results of the study showed that the phosphate ion rate was between (0.0215- 0.0448) mg/L as shown in Figure 9 and Table 2, where the highest value was in Site 2 and the lowest in Site 1 and the reason for the decrease in these values may be due to

the ability to precipitate phosphate in the form of calcium phosphate $\text{Ca}_3(\text{PO}_4)_2$ as well as its adsorption by some clay minerals and organic compounds, which reduces its transfer to the aquatic environment (Ohwoghere-Asuma et. al., 2020).




Figure 9: Phosphate ion rate of the studied spring water (mg/L)

vi. Sodium and Potassium Ions

Figure 10 and Table 2 show that the highest rate of sodium and potassium ion is in site 3, while the lowest is in site one, as their rates ranged between (376-447) and (31.4-36.2) mg / liter, respectively. The reason for the presence of sodium ion may be attributed to the nature of geological formations For the earth layers

through which water passes, as well as the processes of weathering, when water reaches the sod minerals, sodium ions will be released (Hui, 2020). As for the presence of potassium ion in these low concentrations compared to sodium, it is due to its adsorption by soil particles within the geological layers of the studied areas (Ibrahim & Nofal, 2020).




Figure 10: Average of sodium and potassium ions for the studied spring water (mg/L)

vii. Oil and Grease

The results of the current study indicated the concentration rates of oils and greases in the studied sulfur springs, which ranged between (0.169-0.184) mg/L, and the superiority of site 1 in the concentration of oils and grease and the lowest average concentrations in site 2 that the presence of oils and grease in water may indicate that these The water is coming from layers that contain oil, as evidence of the presence of bitumen, because the bitumen is one of the oil derivatives (Mahmoud et al., 2006), as shown in Table (2) and Figure (11).




Figure 11: Oil rate of the studied springs water (mg/L)

c) *Biological tests*i. *Total Plate Count*

Tests of the total number of bacteria are among the important indicators that include total aerobic, facultative and non-autotrophic bacteria. These numbers are usually approximate, but most of them are due to the inability to provide a nutrient medium and suitable conditions for the growth of all species (Juma, 2017). Through the results, it is clear that the number of

bacteria exceeded the limit allowed by the World Health Organization, which ranged between (0.5-3.25) cells / ml. The highest rate was in site 1 and the lowest in site 2 as shown in Table 3 and Figure 11. The increase in bacterial numbers coincided with the high temperatures, which are suitable for the growth and reproduction of bacteria, and the low temperature in the winter season, their numbers decreased (Al-Omar, 2010).




Figure 11: Average total number of bacteria in the studied spring water (cell/ml)

Table 2: Rate of physical and chemical tests of spring water

Test location	Temp.	pH	E.C	T.D.S	T.Alk.	T.H	SO ₄ -2	PO ₄ -3	Na ⁺	K ⁺	Oil
control	14.34	7.86	435.719	217.8	119	154.8	166.9	0.067	18.8	2.48	0.1788
location 1	22.04	6.92	5158.59	2579.6	648	2120	2365.3	0.02154	376	31.4	0.1848
location 2	22.02	6.9	5423.23	2711.8	452	2220	2322.6	0.04482	440.6	34.8	0.1696
location 3	21.74	7	5521.8	2761	649	2080	2310.8	0.0296	447.8	36.2	0.182

Table 3: Total number of bacteria (TPC) $\times 10^5$ cells.ml⁻¹

TPC						
MON. LOC.	11	12	1	2	3	AVG.
Control	HEAVY	295	280	115	126	204
LOC. 1	2	0	5	4	4	3.25
LOC. 2	1	0	0	2	0	0.5
LOC. 3	5	2	1	5	0	2

VI. CONCLUSION

The water is of a moderate type of acidic water, and this water is important for the recovery of diseases due to its high content of sulfates and its high temperature. at the optimum interest. It is considered unfit for drinking if we take into account the concentration of sulfates without regard to the rest of the characteristics. It contains more than 400 mg / liter in addition to the taste and smell. It is also characterized by being very hard because it contains concentrations higher than 180 mg / liter and with high salinity because it contains calcium and magnesium salts.

REFERENCES RÉFÉRENCES REFERENCIAS

1. Aswad, Khaled Galal; Mahmoud, Hazem Gomaa; Muhammad, Yasser Mayser. (2018). Study of the movement of some heavy metals in sulfur water in Al-Kasr village, northern Iraq. The Ninth Periodic Scientific Conference of Dams and Water Resources Research Center, November 27-28.
2. Al-Hamdani, Nour Ahmed Saadoun (2020). Application of water quality indicators to assess the water quality of a number of wells in the left coast of the city of Mosul / Iraq. Master Thesis, College of Environmental Sciences and Technologies, University of Mosul.
3. Al-Zamili, Ayed Jassem; Al-Asadi, Kamel Hamza Fleifel (2014). Variation of the characteristics of groundwater in the western plateau of Najaf Governorate using geographic information systems. Journal of Geographical Research, College of Education for Girls, University of Kufa, Issue Nineteen, 219-238.
4. Al-Safawi, Abdulaziz Younis Talei and Al-Shannouna, Reem Adnan Abdul Razzaq and Al-Sardar, Nour Maysir Sadiq (2018). Evaluation of water quality characteristics and calculation of the coefficient (WQI) for some water sources in Abu Maria village, Tal Afar District, Nineveh Governorate. Journal of Education and Science for Pure Sciences. 27(3): 81-98.
5. Al-Taweeel, Lama Saleh Jabbar (2017). Study of some characteristics of sulfur spring water in Al-Hasya and Al-Khasf regions / Iraq. Al-Qadisiyah Journal of Agricultural Sciences, Volume VII, Issue Two, 167-183.
6. Al-Omar, Muthanna Abdul-Razzaq (2010). Environmental pollution. First Edition, Dar Wael for Printing and Publishing, Amman/Jordan.
7. Al-Lahibi, Abdullah Mahmoud Ajil (2021). Evaluation of the efficiency of the old and unified Shirqat water liquefaction stations in Shirqat district and their efficiency in filtering drinking water. Master's Thesis, College of Science Department of Life Sciences, Tikrit University.
8. Gomaa, Zuhair Gerges (2017). Evaluation of the performance efficiency of some sewage treatment plants in the city of Kirkuk. Master Thesis, College of Education for Pure Sciences, University of Mosul.
9. Muhammad, Sheno Mustafa Ali (2015). Study of groundwater quality in Kirkuk Governorate. Kirkuk University Journal, Volume IV, Issue X, 177-194.
10. Mahmoud, Sharif's headache; Maaleh, Khaldoun Abbas; Ahmed, Hamid Saeed (2006). The mineral water of the springs of Heet-Kabisa area, central western Iraq. Iraqi Geology and Mining Journal, Volume Two, Issue Two, 9-38.
11. Matar, Iyad Abdel-Razzaq; Zidane, Tahseen Ali; Mahmoud, Bashar Abdel Aziz. (2009). Analytical comparative study of lime and sulfur spring water in Hit. Anbar University Journal of Pure Sciences, Volume Three, Issue One, 45- 56.
12. Al-Saffawi, A. A. Y. (2019). Water quality of Nimrud district wells southeast of Mosul city for drinking and civil purpose using the Canadian model of water quality. *Pakistan Journal of Analytical & Environmental Chemistry*, 20(1):75- 81. <http://dx.doi.org/10.21743/pjaec/2019.06.10>
13. Bee, L. B. (2005). The usage of Domestic water Filtration System in Malaysia. Bachelor of Civil Engineering, pp:23-33.
14. Duruibe, J. O.; Ogwuegbu, M. O. C.; & Egwurugu, J. N. (2007). Heavy metal pollution and human biotoxic effects. International Journal of physical sciences, 2(5):112-118.

15. Egbueri, J. C.; Mgbenu, C. N.; and Chukwu, C. N. (2019). Investigating the hydrogeochemical processes and quality of water resources in Ojoto and environs using integrated classical methods. *Modeling Earth Systems and Environment*, 5(4): 1443-1461. <https://doi.org/10.1007/s40808-019-00613-y>

16. Fadipe, O. O.; & Oladejo, K. T. (2020). Temporal variation of groundwater resources in Ilesa West Local Government, Osun State Nigeria. *Nigerian Journal of Technology*, 39(1): 315-324. <http://dx.doi.org/10.4314/njt.v39i1.35>

17. Hui, T.; Jizhong, D.; Shimin, M.; Zhuang, K.; & Yan, G. (2021). Application of water quality index and multivariate statistical analysis in the hydrogeochemical assessment of shallow groundwater in Hailun, northeast China. *Human and Ecological Risk Assessment: An International Journal*, 27(3): 651-667. <https://doi.org/10.1080/10807039.2020.1749827>

18. Ibrahim, L. A.; & Nofal, E. R. (2020). Quality and hydrogeochemistry appraisal for groundwater in Tenth of Ramadan Area, Egypt. *Water Science*, 34(1): 50-64. <https://doi.org/10.1080/11104929.2020.1749411>

19. Larnier, K.; Roux, H.; Dartus, D.; & Croze, O. (2010). Water temperature modeling in the Garonne River (France). *Knowledge and Management of Aquatic Ecosystems*, (398): 04. <https://doi.org/10.1051/kmae/2010031>

20. Merdhah , A. and Yassin, A. (2007). Scale Formation in water Reservoir During Water Injection at Haigh-Salinity Formation Water; *Journal of Applied Science*. 7(21): 3198-3207.

21. Moses, A. N.; & Ishaku, S. (2020). Evaluation of Physico-Chemical Properties of Well Water Qualities in Selected Villages in Zing Local Government Area of Taraba State, Nigeria. *International Journal of Contemporary Research and Review*, 11(03): 20282-20288. <https://doi.org/10.15520/ijcrr.v11i03.797>

22. Ohwoghere-Asuma, O.; Iserhien-Emekeme, R.; Aweto, K. E.; & Ofomola, M. O. (2020). Geophysical investigation of resistivity and groundwater quality in Ogbe-Ijoh coastal area of the western Niger Delta of Nigeria. *Applied Water Science*, 10(2): 1-9. <https://doi.org/10.1007/s13201-020-1144-0>

23. Sharaky, A. M.; & Abdoun, S. H. (2020). Assessment of groundwater quality in Bahariya Oasis, Western Desert, Egypt. *Environmental Earth Sciences*, 79(6): 1-14. <https://doi.org/10.1007/s12665-020-8823-x>

24. Verma, D. K.; Bhunia, G. S.; Shit, P. K.; & Tiwari, A. K. (2018). Assessment of groundwater quality of the Central Gangetic Plain Area of India using Geospatial and WQI Techniques. *Journal of the Geological Society of India*, 92(6): 743-752. <https://doi.org/10.1007/s12594-018-1097-1>